



# PHOS-LIFE ORGANIC™

**P 10.7%, Ca 24.5%, trace elements**

## Organic liquid suspension of micronised calcium phosphate

A high analysis, plant-available source of micronised guano combined in a free-flowing suspension with fulvic acid.

### KEY PERFORMANCE

- An organic, high analysis source of calcium and phosphorus
- Calcium and phosphorus are the governors of plant health and high brix levels
- Tank compatible with most fertilisers

### GROWTH STAGE

- ✓ Pre-planting
- ✓ Planting
- ✓ Vegetative Growth
- ✓ Pre-flower, Flowering, Fruit Set & Fill

### SIZES:

5 L, 15 L, 200 L, 1000 L


### PRODUCT CODE:

MMSOPHL

**ACO Certification No. 456**

### TYPICAL ANALYSIS W/V:

Calcium (Ca)	24.57%
Silicon (Si)	10.73%
Total Phosphorus (P)	10.71%
Phosphorus (P) (citrate insoluble)	7.56%
Phosphorus (P) (citrate soluble)	3.15%
Phosphorus (P) (water soluble)	20 mg/L
Total Potassium (K) (as organic)	0.15%
Manganese (Mn)	1.66%
Iron (Fe)	0.86%
Magnesium (Mg)	0.34%
Zinc (Zn)	0.14%
Copper (Cu)	162 mg/L
SG	1.5
pH	6.5 – 8.5
Conductivity	1 – 5 mS/cm
Appearance	Thick, mid-brown liquid, citrus odour



# PHOS-LIFE ORGANIC™

## APPLICATION RATES

### Soil Ameliorant or Fertilisation

10 – 30 L/ha as required. Use higher rate for soil amelioration.

### Liquid Inject

2 – 5 L/ha. Use a minimum dilution rate of 1 part product to 3 parts water. Rate will depend on crop, row spacing & soil type. Always consider N & P crop requirements for starters based on reliable soil test data.

### Vegetables

**FOLIAR SPRAY RATE** 1 L per 100 L water up to 5 L/ha equivalent

**FREQUENCY** Apply weekly or as required

### Orchards & Vineyards

**FOLIAR SPRAY RATE** 1 L per 100 L water up to 10 L/ha equivalent

**FREQUENCY** Apply every 2 – 4 weeks or as required

### Ornamentals & Turf

**FOLIAR SPRAY RATE** 1 L per 100 L water up to 7 L/ha equivalent

**FREQUENCY** Apply every 2 – 4 weeks or as required

### Broadacre Crops

**FOLIAR SPRAY RATE** 2 – 3 L/ha in 60 – 100 L water, depending on canopy closure

**FREQUENCY** Apply as required

### Pasture

**FOLIAR SPRAY RATE** 2 – 3 L/ha in 60 – 100 L water

**FREQUENCY** Apply as required

### Spot-Spraying

**FOLIAR SPRAY RATE** 50 mL per 10 L water

**FREQUENCY** Apply as required

### Home Garden Application

Dilute 45 mL per 9 L watering can and apply the diluted mix at a rate of 1 litre per square metre – this diluted mix can also be poured over foliage

**FREQUENCY** Repeat every 2 – 4 weeks or as required

\* Disclaimer: Any recommendations provided by Nutri-Tech Solutions Pty Ltd (NTS) or its Distributors are advice only. As no control can be exercised over storage, handling, mixing, application or use, or weather, plant or soil conditions before, during or after application (all of which may affect the performance of our program), no responsibility for, or liability for any failure in performance, losses, damages, or injuries (consequential or otherwise), arising from such storage, mixing, application, or use will be accepted under any circumstances whatsoever. NTS recommend you contact an Agronomist prior to product application. The Buyer assumes all responsibility for the use of any NTS products.

### Nutri-Tech Solutions P/L (NTS)

7 Harvest Road, Yandina, Queensland 4561, Australia

P +61 7 5472 9900 | E info@nutri-tech.com.au

[www.nutri-tech.com.au](http://www.nutri-tech.com.au)

## INSTRUCTIONS

### Shake or stir well before use and maintain constant agitation.

It is advisable to pre-mix, adding this product slowly while maintaining agitation in spray tank to prevent settling of solids. Will pass through 200 micron filter under ideal conditions, however, a coarse inline filter (500 microns/35 mesh) is recommended to ensure the delivery lines are free from particle contaminants from other sources. Where higher water rates are required for good foliar coverage, do not exceed maximum product rate per hectare.

MMS products will leave a visible residue on leaves and fruit when used as a foliar. Avoid spraying close to harvest if produce is susceptible to staining. Sensitive foliage or plants should be test sprayed prior to full scale application. Frequencies are a guideline only and can be adjusted according to farm practices and/or in conjunction with crop monitoring. Ensure suitable spray conditions (e.g. temperature, humidity, wind-speed), prior to foliar application.

Compatibility and/or performance cannot be guaranteed when combined with other products. Jar test for compatibility and spray test on small area before wide-spread use.

## IMPORTANT

**Note:** Product in 1000 L tanks may weigh over 1 tonne. Forklift with 2 tonne lifting capacity is recommended.

## STORAGE & HANDLING

Store in a cool dry area out of direct sunlight. This product has been tested to store safely down to 5°C. Sedimentation may occur below 5°C. Do not pre-mix or store in diluted form.

MMS products are very heavy and every care should be taken when handling these materials to avoid back injury.

## YOUR LOCAL NTS DISTRIBUTOR: