

SEACHANGE LIQUID KELP™



Cold composted Australian bull kelp

Liquid Australian Bull Kelp *Durvillaea potatorum*, manufactured with a proprietary processing system. The 90-day cold-composting process retains all of the natural attributes of bull kelp, with none of the denaturing features inherent in some other manufacturing procedures that involve heat and alkaline extraction. The attributes include: minerals, amino acids, natural growth promotants, chelating agents and complex sugars. This product has been developed as part of an NTS quest to reduce the costs to the Australian farmer of the core components of biological agriculture.

KEY PERFORMANCE

- Powerful rescue remedy
- Healthy, balanced growth
- Improve root growth

GROWTH STAGE

- ✓ Planting
- ✓ Vegetative Growth
- ✓ Pre-flowering

SIZES:

1L, 5 L, 20 L, 200 L, 1000 L

PRODUCT CODE:

SCLK

ACO Certification No. 456

TYPICAL ANALYSIS W/V:

Total Potassium (K) (as organic)	1.39%
Calcium (Ca)	0.09%
Magnesium (Mg)	0.09%
Iodine (I)	22 mg/L
SG	1.0
pH	3.6 – 4.6
Conductivity	30 – 50 mS/cm
Appearance	Brown Liquid



SEACHANGE LIQUID KELP™

APPLICATION RATES

Vegetables

FOLIAR SPRAY	1 L per 100 L water up to 5 L/ha equivalent
FERTIGATION	10 – 20 L/ha
FREQUENCY	Weekly or as required

Orchards & Vineyards

FOLIAR SPRAY	1 L per 100 L water up to 7 L/ha equivalent
FERTIGATION	10 – 20 L/ha
FREQUENCY	Every 2 – 4 weeks or as required

Ornamentals & Turf

FOLIAR SPRAY	1 L per 100 L water up to 5 L/ha equivalent
FERTIGATION	10 – 20 L/ha
FREQUENCY	Every 2 – 4 weeks or as required

Broadacre Crops

FOLIAR SPRAY	2 – 3 L/ha in 80 – 100 L water, depending on canopy closure
FREQUENCY	As required

Pasture

FOLIAR SPRAY	2 – 3 L/ha in 80 – 100 L water
FREQUENCY	As required

Spot-Spraying

FOLIAR SPRAY	50 mL per 10 L water
FREQUENCY	As required

Home Garden Application

Dilute 30 mL per 9 L watering can and apply the diluted mix at a rate of 1 litre per square metre. Apply fortnightly or as required.

Tree Tubestock or Vegetable Seedling Treatment

Soak roots in 1:150 dilution

*Disclaimer: Any recommendations provided by Nutri-Tech Solutions Pty Ltd (NTS) or its Distributors are advice only. As no control can be exercised over storage, handling, mixing, application or use, or weather, plant or soil conditions before, during or after application (all of which may affect the performance of our program), no responsibility for, or liability for any failure in performance, losses, damages, or injuries (consequential or otherwise), arising from such storage, mixing, application, or use will be accepted under any circumstances whatsoever. NTS recommend you contact an Agronomist prior to product application. The Buyer assumes all responsibility for the use of any NTS products.

Nutri-Tech Solutions P/L (NTS)

7 Harvest Road, Yandina, Queensland 4561, Australia

P +61 7 5472 9900 | E info@nutri-tech.com.au

www.nutri-tech.com.au

INSTRUCTIONS

Shake or stir well before use. Where higher water rates are required for good foliar coverage, do not exceed maximum product rate per hectare. Sensitive foliage or plants should be test sprayed prior to full scale application.

Frequencies are a guideline only and can be adjusted according to farm practices and/or in conjunction with crop monitoring. Ensure suitable spray conditions (e.g. temperature, humidity, wind-speed), prior to foliar application.

Compatibility and/or performance cannot be guaranteed when combined with other products. Jar test for compatibility and spray test on small area before wide-spread use.

IMPORTANT

Contains some larger particles (up to 100 micron) which may block fine spray nozzles.

Some separation may occur over time. Agitate well during application to ensure solution is homogenous.

Avoid spraying close to harvest if produce is susceptible to staining.

Note: Product in 1000 L tanks may weigh over 1 tonne. Forklift with 2 tonne lifting capacity is recommended.

STORAGE & HANDLING

Store in a cool dry area out of direct sunlight. This product has been tested to store safely down to 5°C. Crystallisation or sedimentation may occur below 5°C. Do not pre-mix or store in diluted form. Seal lid/cap immediately after use.

YOUR LOCAL NTS DISTRIBUTOR: