

	Cocaine	Marijuana	Opiates	Amphetamines	PCP
Medical Names	Benzoylecgonine Cocaethylene Norococaine	THCA 11-Nor-delta-9-tetrahydrocannabinol-9-carboxylic acid	Hydrocodone (Vicodin, Lortab, Lorcet, Hycodan, Vicoprofen), Hydromorphone (Dilaudid), Oxycodone (Percocet, Percodan, Tylox, Oxycotin), Oxymorphone (Numorpha), Codine, Morphine	MDMA (Ecstasy) MDA MDEA (Eve) Methamphetamine Amphetamine	Phencyclidine
Street Names	Blow, Coke, Big C, Flake, Snow, Crack, White Girl, C, White Candy, Freebase, Toot, Nose Candy, Rock	Pot, Reefer, Dope, Mary Jane, Ganja, Hash, Blunts, Aunt Mary, Joint, Smoke, Weed	Heroin (Smack, Horse, Junk, China White), Hillbilly Vikes, Dillies, Hydro, Dust, Juice, M, Miss Emma, School Boy, Oxy, OCs, Ox, 40, 40-Bar, Kicker, Cotton, footballs, M2, Dust,	Ecstasy, Eve, E, X, XTC, Adam, Essence, Roll, Love Drug, Uppers, Speed, Meth, Whites, Dexies, Crank, Black Beauties, Benny, Crystal, Jelly Beans, White Cross	Angel Dust, Devil Stick, PCP, Dummy Dust, Wack, Embalming Fluid, Killer Weed, Super Grass
How Used	Snorted, Dissolved in water or injected	Smoked or Eaten	Swallowed or Injected	Pill form Swallowed	Tablets, capsules or colored powder. Snorted, Smoked or eaten.
What to Look For	Glassine envelopes, razors, small spoons, bitter white crystalline powder, granular rocks	Plastic baggies, rolling paper, "roach" clips, odor of burnt rope	Glassine envelopes, needles & syringes, caps or spoons, tourniquets, needle marks on arms	Tablets of varying colors, possible chain smoking.	Liquid capsules, white or brown powder on paper stamps, sugar cubes, cigarettes, joints, injection paraphernalia
Dangers	Shallow breathing, fever, anxiety, tremors, possible death from convulsions or respiratory arrest	Psychological dependence, increased heart rate, impaired short-term memory, anxiety, lung damage, possible psychosis with chronic use	Lethargy, weight loss, hepatitis, slow & shallow breathing, possible death	Disorientation, severe depression, paranoia, possible hallucinations, increase in blood pressure & heart rate, seizures, fatigue	Unpredictable behavior, flashbacks, possible emotional instability & psychosis, hallucinations, seizures, coma, death
Short-Term Effects	Short-lived euphoria changing to depression, irritability, nervousness, tightness of muscles, dilated pupils, increased temperature, heart rate, blood pressure, insomnia, loss of appetite, restless, irritability, anxiety, fatigue	Altered perception, dilated pupils, lack of concentration & coordination, craving for sweets, increased appetite, laughter, memory & learning problems, distorted perceptions (sight, sounds, time, touch) loss of motor coordination, increased heart rate, anxiety, dry mouth/sore throat	Insensitivity to pain, euphoria, sedation, nausea, vomiting, itchiness, watery eyes, running nose, drowsiness, constipation, slowed breathing,	Long periods without rest or sleep, loss of appetite, anxiety, irritability, rapid speech, tremors, mood elevation, feelings of peacefulness, acceptance & empathy, teeth clinching, loss of inhibitions, transfixion on sights & sounds, nausea, blurred vision, chills/sweating, increased stamina & physical energy	Increased pulse & heart rate, blood pressure & temperature, possible mood & perception alteration, hallucinations, paranoia, panic, anxiety, nausea, tremors, suicidal urge, shallow breathing, vomiting, drooling, loss of balance, dizziness, flicking up/down of eyes, sparse/garbled speech
Long-Term Effects	Paranoia, aggressive paranoid behavior, depression, ulceration of mucous membranes, death	Cancer causing chemicals, hallucinations, bronchitis, emphysema, bronchial asthma, anti-motivational syndrome, lung damage, risk to reproductive system	Severe respiratory depression that can lead to death, addiction, muscle/bone pain, insomnia, diarrhea, vomiting, cold flashes w/ goose bumps.	Heart problems, stroke, liver, kidney and lung damage, erectile dysfunction, insomnia, delusions, paranoia, highly addictive, long term cognitive impairments, nerve cell damage, psychiatric disturbances	Very addictive, psychological dependence, craving, memory loss, speech difficulties, depression, weight loss, mood disorders
Time Detectable	<b>Urine:</b> 1-3 days <b>Oral Fluid:</b> 24-36 Hrs <b>Hair:</b> Up to 90 days	<b>Urine:</b> 1-3 days occasional use, up to 30 days habitual/chronic use <b>Oral Fluid:</b> less than 24 hours <b>Hair:</b> Up to 90 days	<b>Urine:</b> 1-3 days <b>Oral fluid:</b> 24-36 hrs <b>Hair:</b> up to 90 days	<b>Urine:</b> 1-3 days <b>Oral Fluid:</b> 24-36 Hrs <b>Hair:</b> Up to 90 days	<b>Urine:</b> 1-5 days occasional use, up to 30 days habitual/chronic use <b>Oral Fluid:</b> less than 24 hours <b>Hair:</b> Up to 90 days