

Appendix Eight

Healthy Baby Growth Charts

Signs of Adequate Nutrition

Chart One—Week One

If you're breast-feeding, monitoring your baby's growth is of vital concern. How do you know if your baby is getting enough food to grow on? There are a number of objective indicators to healthy growth and proper nutrition. Indicators of healthy baby growth provide mom guidance and feedback as to how well she and her baby are doing. The following indicators represent healthy signs of growth during the first week of life:

- 1. Your baby goes to the breast and nurses.
- 2. Your baby is nursing a minimum of eight times in a 24-hour period.
- 3. Your baby is nursing over 15 minutes at each nursing period.
- 4. You can hear your baby swallowing milk.
- 5. Your baby has passed his first stool called meconium. (Make sure you let the nurses know that you are tracking your baby's growth indicators.)
- 6. Your baby's stooling pattern progresses from meconium (greenish black) to brownie batter transition stools, to yellow stools by the fourth or fifth day. This is one of the most positive signs that your baby is getting enough milk.
- 7. Within 24 to 48 hours, your baby starts having wet diapers, (increasing to three to five a day). By the end of the first week wet diapers are becoming more frequent.

Unhealthy growth indicators for the first week:

- 1. Your baby is not showing any desire to nurse or has a very weak suck.
- 2. Your baby fails to nurse eight times in a 24-hour period.
- 3. Your baby tires quickly at the breast and cannot sustain at least 15 minutes of nursing.
- 4. Your baby continually falls asleep at the breast before taking a full feeding.
- 5. You hear a clicking sound accompanied by dimpled checks while baby is nursing.
- 6. Your baby's stooling pattern is not progressing to yellow stools within a week's time.
- 7. Your baby has not had any wet diapers within 48 hours of birth.

Using the chart to keep track of your baby's vital health indicators can make the difference between healthy and unhealthy growth. If you wish, make a copy of the chart and place it in a convenient location (on a refrigerator, above the crib, etc.). Place the appropriate (V) mark or letter designated for each occurrence. For example, if your baby nurses nine times on day two, then place nine checks on that day. If your baby passes his first meconium stool on the second day, then place an "M" on that day. Knowing what to expect and measuring results will get you and your baby off to a great start.

RT: Chart One Week 1	DAY 3 DAY 4 DAY 5 DAY 6 DAY 7					7–10 days: Weightlb./oz. Lengthinches Any two consecutive days of deviation from what is listed as normal should be reported immediately to your pediatrician.
HEALTHY BABY GROWTE	Place a check (V) for each feeding in a 24 hour period. (Minimum of 8 feedings a day.)	Place a check (V) for each nursing period of 15 or more minutes in length.	Place a "M" for the first stool (Meconium) and a "T" for each brownie battered transition stool.	Place a "Y" to record each yellow stool. (Milk stools should appear by the 4th or 5th day.)	Place a check (V) for each wet diaper. (Wet diapers should start to appear by 48 hours or sooner.)	7–10 days: Weight Any two consecutive days of deviation from what is listed as

S © Gary Ezzo & Robert Bucknam

Signs of Adequate Nutrition

Chart Two—Weeks Two through Four

Just because things have gone well in the first week does not mean you can slack off from monitoring your baby's healthy growth signs. After the first week, some of the healthy growth indicators begin to change. This chart represents healthy baby growth indicators to be monitored over the next three weeks. Please note the changes:

Here is the checklist for the next three weeks.

- 1. Your baby is nursing at least eight times a day.
- 2. Your baby over the next three weeks has two to five or more yellow stools daily. (This number will probably decrease after the first month.)
- 3. Your baby should start to have six to eight wet diapers a day, some saturated.
- 4. Your baby's urine is clear, not yellow.
- 5. Your baby has a strong suck, you see milk, and you can hear an audible swallow.
- 6. You're noticing increased signs of alertness during your baby's waketime.
- 7. Your baby is gaining weight and growing in length.

Unhealthy growth indicators are:

- 1. Your baby is not getting eight feedings a day.
- 2. Your baby has small, scant, and infrequent stools.
- 3. Your baby does not have the appropriate number of wet diapers given his age.
- 4. Your baby's urine is concentrated and bright yellow.
- 5. Your baby has a weak or tiring suck and you cannot hear him swallow.
- 6. Your baby is sluggish or slow to respond to stimulus, and does not sleep between feedings.
- 7. Your baby is not gaining weight or growing in length. Your doctor will direct you in the best strategy to correct this problem.

Any two consecutive days of deviation from what is listed above as normal should be reported immediately to your pediatrician. Using the chart to keep track of your baby's vital health indicators can make the difference between healthy and unhealthy growth. If you wish, make copies of the chart and place it in a convenient location (on a refrigerator, above the crib, etc.). For your assurance record the results with a $(\sqrt{})$ mark for each occurrence of each healthy indicator. For example, six wet diapers on Monday should have six checks in the appropriate box. Knowing what to expect and measuring the expected results against the actual will provide you security and confidence as your baby grows.

HEALTHY BABY GROWTH CHART: Chart Two Weeks 2-4	Y GRO		WTH CHART	: Char	t Two	Weeks 2	4-7
Healthy Growth Indicators	DAY 1	DAY 2	DAY 3	Day 4	DAY 5	DAY 6	DAY 7
Place a check (V) for each feeding in a 24 hour period. (Minimum of 8 feedings a day.)							
Place a check (V) for each wet diaper per day with clear urine. (Norm per day: 5 to 7.)							
Place a check (V) for each wet diaper with yellow concentrated urine. (Norm per day: 0.)							
Place a check (V) for each yellow stool. (For the first month, 2 to 5 or more, per day.)							

© Gary Ezzo & Robert Bucknam 225

Any two consecutive days of deviation from what is listed as normal should be reported immediately to your pediatrician.



HEALTHY BABY GROWTH CHART: Chart Two Weeks 2-4 Summary of Each Day Healthy Growth Indicators Blace a check (v) for each keeling in a 24 hour period. (Minimum of 8 feedings a day.) Blace a check (v) for each wet diaper per day with clear urine. (Norm per day: 5 to 7.) Place a check (v) for each wet diaper with yellow concentrated urine. (Norm per day: 0.) Place a check (v) for each yellow stool. (For the first month, 2 to 5 or more, per day.)

© Gary Ezzo & Robert Bucknam

HEALTHY CROWTH INDICATORS DAY 1 DAY 2 DAY 5 DAY 5 DAY 6 DAY 6 DAY 6 DAY 6 DAY 6 DAY 7 DAY 6 DAY 6 DAY 6 DAY 6 DAY 7 DAY 6 DAY 6 DAY 6 DAY 7 DAY 6 Steelings a day.) Place a check (v) for each wet diaper per day: 5 to 7.) Place a check (v) for each wet diaper with yellow concentrated urrine. (Norm per day: 0.) Place a check (v) for each yellow stool (For the first month, 2 to 5 or more, per day:)	HEALTHY BABY GROWTH CHART: Chart Two Weeks 2-4	Y GRO		HART	: Char	t Two	Weeks 2	4-7
Place a check (v) for each feeding in a 24 hour period. (Minimum of 8 feedings a day.) Place a check (v) for each wet diaper per day with clear urine. (Norm per day: 5 to 7.) Place a check (v) for each wet diaper with yellow concentrated diaper with yellow concentrated urine. (Norm per day: 0.) Place a check (v) for each yellow stool. (For the first month, 2 to 5 or more, per day.)	Healthy Growth Indicators	$ m D_{AY}~1$	DAY 2	DAY 3	DAY 4	DAY 5	Day 6	$D_{AY} T$
Place a check (V) for each wet diaper per day with clear urine. (Norm per day: 5 to 7.) Place a check (V) for each wet diaper with yellow concentrated urine. (Norm per day: 0.) Place a check (V) for each yellow stool. (For the first month, 2 to 5 or more, per day.)	Place a check (V) for each feeding in a 24 hour period. (Minimum of 8 feedings a day.)							
Place a check (V) for each wet diaper with yellow concentrated urine. (Norm per day: 0.) Place a check (V) for each yellow stool. (For the first month, 2 to 5 or more, per day.)	Place a check ($$) for each wet diaper per day with clear urine. (Norm per day: 5 to 7.)							
Place a check (V) for each yellow stool. (For the first month, 2 to 5 or more, per day.)	Place a check (V) for each wet diaper with yellow concentrated urine. (Norm per day: 0.)							
	Place a check (V) for each yellow stool. (For the first month, 2 to 5 or more, per day.)							

© Gary Ezzo & Robert Bucknam

Any two consecutive days of deviation from what is listed as normal should be reported immediately to your pediatrician.

Signs of Adequate Nutrition

Chart Three—Weeks Five through Ten

This third chart differs from the second only in the number of stools eliminated. Basically the rest of the chart is the same. Continue to monitor your baby's growth, especially after your baby starts sleeping through the night.

Here is the checklist for the next six weeks:

- 1. Your baby is nursing at least seven to eight times a day.
- 2. Your baby's stooling pattern again changes. Your baby may have several small stools or one large one. He may have several a day or one every couple of days.
- 3. Your baby should have six to eight wet diapers a day, some saturated.
- 4. Your baby's urine is clear, not yellow.
- 5. Your baby has a strong suck, you see milk, and you can hear an audible swallow.
- 6. You are seeing increasing signs of alertness during your baby's waketime.
- 7. Your baby is gaining weight and growing in length.

Unhealthy growth indicators are:

- 1. Your baby is not getting a minimum of seven feedings a day.
- 2. Your baby has small, scant, and infrequent stools.
- 3. Your baby does not have the appropriate number of wet diapers given his age.
- 4. Your baby's urine is concentrated and bright yellow.
- 5. Your baby has a weak or tiring suck and you cannot hear him swallow.
- 6. Your baby is sluggish or slow to respond to stimulus, and does not sleep between feedings.
- 7. Your baby is not gaining weight or growing in length. Your doctor will direct you in the best strategy to correct this problem.

Any two consecutive days of deviation from what is listed above as normal should be reported immediately to your pediatrician. Using the chart to keep track of your baby's vital health indicators can make the difference between healthy and unhealthy growth. If you wish, make copies of the chart and place it in a convenient location (on a refrigerator, above the crib, etc.). For your assurance, record the results with a (\mathbf{v}) mark for each occurrence of each healthy indicator. For example, six wet diapers on Monday should have six checks in the appropriate box. Knowing what to expect and measuring the expected results against the actual will provide you security and confidence as your baby grows.

HEALT	HEALTHY BABY GROWTH CHART: Chart Three Weeks 5-10	GROW		TH CHART:	Chart	Three 1	Weeks &	5-10
HEALTHY GROWTH INDICATORS	i Indicators	$ m D_{AY}~1$	DAY 2	т <u>Басп</u> <i>Ва</i> у	Day 4	Day 5	DAY 6	DAY 7
Place a check (V) for each feeding. (Minimum should be 7-8 in a 24-hour period.)	or each feeding. be 7-8 in a							
Place a check (V) for each wet diaper per day with clear urine. (Norm per day: 5 to 7.)	or each wet clear urine.							
Place a check (V) for each wet diaper with yellow concentrated urine. (Norm per day: 0.)	or each wet concentrated lay: 0.)							
Place a check (V) for each stool per day.	or each stool							
Any two consec	Any two consecutive days of deviation from what is listed as normal should be reported immediately to your pediatrician.	ion from whai	t is listed as no	ormal should	be reported in	nmediately ta	o your pediatı	rician.

© Gary Ezzo & Robert Bucknam



HEALTHY BABY GROWTH CHART: Chart Three Weeks 5-10	WTH CHART:	Chart Three V	Weeks 5-10
Healthy Growth Indicators Day 1	DAY 2 DAY 3	DAY 4 DAY 5	DAY 6 DAY 7
Place a check (v) for each feeding. (Minimum should be 7-8 in a 24-hour period.)			
Place a check (v) for each wet diaper per day with clear urine. (Norm per day: 5 to 7.)			
Place a check (v) for each wet diaper with yellow concentrated urine. (Norm per day: 0.)			
Place a check (v) for each stool per day.			
Any two consecutive days of deviation from what is listed as normal should be reported immediately to your pediatrician.	nat is listed as normal <i>should</i>	l be reported immediately to	your pediatrician.

© Gary Ezzo & Robert Bucknam

HEALTHY BABY GROWTH CHART: Chart Three Weeks 5-10	MO3	TH CI	HART:	Chart	Three	Weeks (5-10
Healthy Growth Indicators Day 1	r 1	DAY 2 DAY 3	r Each Day DAY 3	DAY 4	D_{AY} 5	Day 6	Day 7
Place a check (V) for each feeding. (Minimum should be 7-8 in a 24-hour period.)							
Place a check (V) for each wet diaper per day with clear urine. (Norm per day: \mathfrak{S} to \mathbb{Z} .)							
Place a check (V) for each wet diaper with yellow concentrated urine. (Norm per day: 0.)							
Place a check (V) for each stool per day.							
Any two consecutive days of deviation from what is listed as normal should be reported immediately to your pediatrician.	m what	is listed as no	ormal <i>should</i>	be reported in	nmediately t	o your pediat	rician.

© Gary Ezzo & Robert Bucknam



HEALTHY BABY GROWTH CHART: Chart Three Weeks 5-10	WTH CI	TH CHART: Summary of Each Day	Chart	Three	Weeks 5	-10
Healthy Growth Indicators Day 1	DAY 2	DAY 3	DAY 4	DAY 5	Day 6	DAY 7
Place a check ($$) for each feeding. (Minimum should be 7-8 in a 24-hour period.)						
Place a check ($$) for each wet diaper per day with clear urine. (Norm per day: 5 to 7.)						
Place a check (V) for each wet diaper with yellow concentrated urine. (Norm per day: 0.)						
Place a check (V) for each stool per day.						
Any two consecutive days of deviation from what is listed as normal should be reported immediately to your pediatrician.	nat is listed as no	ormal <i>should</i>	be reported in	mmediately t	o your pediatri	cian.

© Gary Ezzo & Robert Bucknam

HEALTHY BABY GROWTH CHART: Chart Three Weeks 5-10	MO3	TH CI	HART:	Chart	Three	Weeks (5-10
Healthy Growth Indicators Day 1	r 1	DAY 2 DAY 3	r Each Day DAY 3	DAY 4	D_{AY} 5	Day 6	Day 7
Place a check (V) for each feeding. (Minimum should be 7-8 in a 24-hour period.)							
Place a check (V) for each wet diaper per day with clear urine. (Norm per day: \mathfrak{S} to \mathbb{Z} .)							
Place a check (V) for each wet diaper with yellow concentrated urine. (Norm per day: 0.)							
Place a check (V) for each stool per day.							
Any two consecutive days of deviation from what is listed as normal should be reported immediately to your pediatrician.	m what	is listed as no	ormal <i>should</i>	be reported in	nmediately t	o your pediat	rician.

© Gary Ezzo & Robert Bucknam



HEALTHY BABY GROWTH CHART: Chart Three Weeks 5-10	eks 5-10
HEALTHY GROWTH INDICATORS DAY 1 DAY 2 DAY 3 DAY 4 DAY 5 DAY	DAY 6 DAY 7
Place a check (V) for each feeding. (Minimum should be 7-8 in a 24-hour period.)	
Place a check (v) for each wet diaper per day with clear urine. (Norm per day: 5 to 7.)	
Place a check (v) for each wet diaper with yellow concentrated urine. (Norm per day: 0.)	
Place a check (V) for each stool per day.	
Any two consecutive days of deviation from what is listed as normal should be reported immediately to your pediatrician.	r pediatrician.

© Gary Ezzo & Robert Bucknam