

POWDER

# BIOMIN® BORON

5-0-0

Complexed Minerals for Soil and Foliar Application



## Product Information

**BIOMIN BORON** contains nitrogen and complexed boron for soil and/or foliar applications. It is completely bioavailable and non-phytotoxic to plants when applied in accordance to directions. **BIOMIN BORON** is designed to prevent and correct boron deficiency and to boost crops during critical or fast growing periods. **BIOMIN BORON** may be applied with nitrogen fertilizers.

### GUARANTEED ANALYSIS

Total Nitrogen (N).....	5.0%
5% Water Soluble Organic Nitrogen	
Boron (B).....	15.0%

Derived from Sodium Borate, Citric Acid (Chelating agent), and Hydrolyzed Vegetable Protein.

### KEEP OUT OF REACH OF CHILDREN

**CAUTION** - This product contains boron. Use of this product on any crop other than those listed on the label may result in serious crop injury. Do not apply to crops that will be grazed by ruminant animals. Contact your local county agency or field consultant for further information.

**COMPATIBILITY:** Always perform a compatibility test before mixing any products with **BIOMIN BORON**.

### PREVENTS AND CORRECTS BORON DEFICIENCIES

**NET CONTENTS:** 10 Pounds (4.5 kgs)

U.S. Pat.# 5,504,055

Lot #:

## Directions and General Recommendations

### SOIL APPLICATION

**BIOMIN BORON** may be applied before or at planting through irrigation water, as banding or broadcast. The application rate is 1.5-5.0 pounds per acre (1.7-5.6 Kg/Ha) depending on crop, soil conditions and application methods. For maintenance, 1.5 pounds per acre (1.7 Kg/Ha) are usually recommended.

### FOLIAR APPLICATION

**FIELD CROPS:** Apply 1.0-1.5 pounds of **BIOMIN BORON** per acre (1.1-1.7 Kg/Ha) when crops are 6-12 inches (15-30 cm) high or when deficiency symptoms appear. Repeat application 2-4 weeks later.

**VEGETABLES AND POTATOES:** Apply 1.0-1.5 pounds of **BIOMIN BORON** per acre (1.1-1.7 Kg/Ha) at about 2-4 weeks after emergence or if deficiency symptoms appear. Repeat the application 2-3 weeks later.

**GREENHOUSE CROPS:** Apply 1.0-1.5 pounds of **BIOMIN BORON** per acre (1.1-1.7 Kg/Ha) 2-3 weeks after emergence or transplanting, or when boron deficiencies first appear. Repeat the application every one or two weeks.

### LIMITED WARRANTY

Manufacturer or seller makes no warranty, whether expressed or implied, concerning the use of this product other than for the purposes indicated on the label. Neither manufacturer nor seller shall be liable for any injury or damage caused by this product due to misuse, mishandling or any application not specifically described on the label.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>

**FRUIT TREES AND GRAPES:** Apply 1.0-1.5 pounds of **BIOMIN BORON** per acre (1.1-1.7 Kg/Ha) in early spring when new leaves emerge or when deficiency symptoms appear. Repeat the application 2-4 weeks later.

**BIOMIN BORON** is best applied when plants are in an active growing stage, after irrigation or rainfall. For foliar application, it should be diluted with at least 200 times of water. For best results and more specific recommendations, contact your nearest distributor. Always perform a compatibility test before mixing any products with **BIOMIN BORON**.

**BIOMIN BORON** can be applied to ALL commercial crops including all vegetables, potatoes, carrots, onions, all berries and strawberries, all brassicas, all fruit trees, avocados, persimmons, citrus, all nut trees, vineyards, kiwifruit, tropical fruit, lettuce, asparagus, all melons, beans, ginger, ornamentals, cut flowers and pastures.

**JH Biotech Inc.**

MANUFACTURED BY: JH BIOTECH, INC. P.O. Box 3538, Ventura, CA 93006 U.S.A. Phone: 805-650-9933 Website: [www.jhbiotech.com](http://www.jhbiotech.com)

Copyright © 2014 All rights reserved

0407-D-W-1