

BIOMIN[®] COPPER POWDER



Chelated Mineral for Soil and Foliar Application

GUARANTEED ANALYSIS

Copper (Cu).....	17%
As Amino Acid:	
Nitrogen (N).....	6.0%
Copper is present as a glycine chelate. Nitrogen present as an amino acid.	

PRODUCT INFORMATION

BIOMIN COPPER is a chelated mineral which is chelated mainly by glycine and other organic amino acids. **BIOMIN COPPER** was designed for soil and/or foliar applications. Completely bioavailable and non-phytotoxic to plants when applied in accordance to directions. Designed to prevent and correct copper deficiencies and to boost crops during critical or fast growing periods.

BIOMIN COPPER can be applied to ALL crops including all vegetables, potatoes, carrots, onions, all berries and strawberries, all brassicas, all fruit trees, avocados, persimmons, citrus, all nut trees, vineyards, kiwifruit, tropical fruit, lettuce, asparagus, all melons, beans, ginger, ornamentals, cut flowers and pastures.

CAUTION- Do not apply this product after application of copper containing products as this may cause phytotoxicity.

KEEP OUT OF REACH OF CHILDREN

PREVENTS AND CORRECTS MINERAL DEFICIENCIES

COMPATIBILITY: Always perform a compatibility test before mixing any products with **BIOMIN COPPER**.

DIRECTIONS AND GENERAL RECOMMENDATIONS:

SOIL APPLICATION

BIOMIN COPPER may be applied before or at planting through irrigation water. The application rate is 0.5 – 2.0 kilograms per hectare depending on crop, soil conditions and application methods.

FOLIAR APPLICATION

ALL FIELD CROPS: Apply 0.5 – 1.0 kilograms of **BIOMIN COPPER** per hectare when deficiency symptoms appear. Repeat application as needed.

ALL VEGETABLE CROPS :Apply 100 - 500 grams of **BIOMIN COPPER** per

hectare when deficiency symptoms appear. Repeat application as needed.

GREENHOUSE CROPS: Apply 30 - 50 grams of **BIOMIN COPPER** per 100 liters of water 2-3 weeks after emergence or transplanting. Do not apply to tomatoes after fruit set.

APPLES: Apply 0.5 – 2.0 kilograms of **BIOMIN COPPER** per hectare post harvest at 6 weeks before dormancy only.

FRUIT TREES: Apply 0.5 – 2.0 kilograms of **BIOMIN COPPER** per hectare post harvest at 6 weeks before dormancy only.

KIWIFRUIT: Apply 0.5 – 2.0 kilograms of **BIOMIN COPPER** per hectare post harvest at 6 weeks before dormancy only.

GRAPES: Apply 0.5 – 2.0 kilograms of **BIOMIN COPPER** per hectare post harvest at 6 weeks before dormancy.

BIOMIN COPPER is best applied when plants are in an active growing stage, after irrigation or rainfall. For foliar application, it should be diluted with at least 100 times of water.

BIOMIN COPPER should be used strictly at label. For best results and more specific recommendations, contact your nearest distributor.

LIMITED WARRANTY

Manufacturer or seller makes no warranty, whether expressed or implied, concerning the use of this product other than for the purposes indicated on the label. Neither manufacturer nor seller shall be liable for any injury or damage caused by this product due to misuse, mishandling, or any application not specifically described on the label.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>.

MANUFACTURED BY: JH BIOTECH, INC. P.O. Box 3538, Ventura, CA 93006, USA

NET CONTENTS:

1 Kilograms 2 Kilograms 5 Kilograms 10 Kilograms 25 Kilograms

LOT NO.:

JH Biotech, Inc.

0402-I-G-2
Copyright © 2011 JH Biotech, Inc.
All rights reserved.