

Fast and effective weed control of Couch and other grasses

 **Fusilade Forte**[®]

syngenta[®]

FUSILADE[®] FORTE combines the proven grass weed control and crop safety of FUSILADE with Syngenta's patented ISOLINK surfactant technology for improved performance.

Reliable control of grass weeds

FUSILADE FORTE provides powerful, post-emergent control of many annual grasses and certain perennial grasses in dozens of horticultural crops, ornamentals and nursery trees. It features a highly flexible application window and a high level of crop safety.

Unique formulation

FUSILADE FORTE features ISOLINK technology which combines the active ingredient, surfactants and oils into tiny micro-droplets within an emulsifiable concentrate. There is no need to add any surfactants or oils to FUSILADE FORTE.

Active ingredient

FUSILADE FORTE contains 128g/L of fluazifop-P, present as the butyl ester, with 156g/L hydrocarbon solvent in an emulsifiable concentrate formulation. Fluazifop-P is a member of the "fop" group of herbicides. For resistance management purposes, it is a Group A herbicide.

Mode of action

FUSILADE FORTE has the inhibitor of acetyl coA carboxylase mode of action. The active ingredient inhibits the synthesis of fatty acids and lipids, which are needed to maintain cell integrity and store energy. Plant growth stops within two days and susceptible weeds stop competing with the crop for available moisture and nutrients. Symptoms such as discolouration, necrosis and swelling are evident in 7 to 10 days.

Improved mixing and spraying

FUSILADE FORTE forms a stable emulsion when mixed with water, ensuring uniform distribution throughout the spray mixture.

The advanced formulation technology produces more uniform spray droplets, reducing the potential for spray drift or large droplets bouncing off the leaf surface of target weeds.

Improved spreading and uptake

FUSILADE FORTE with ISOLINK technology ensures more active ingredient is deposited on and retained by the leaf surface. The surfactants aid wetting and spreading, while the oils maximise uptake of the active ingredient through the leaf cuticle and throughout the plant.

Rapid absorption

FUSILADE FORTE is rapidly absorbed by leaves and green stems of grass weeds and is then translocated to the growing points of the plant, including the leaf tips, roots, rhizomes and stolons. This attribute means FUSILADE FORTE has good activity against perennial and annual grass weeds.

Superior rainfastness

FUSILADE FORTE is rainfast within one hour, significantly faster than other "fops". Laboratory studies on volunteer wheat and barley have confirmed that light rainfall after spraying has little - if any - effect on performance.




Figure 1: Irrigation (2.5mm) applied 30 minutes after application to simulate rain washing.

Couch grass control

FUSILADE FORTE delivers thorough and cost-effective control of Couch Grass in tree crops and grape vines when applied at 3.3 to 6.6 L/ha as a single application in spring or autumn. Use the higher rate for well established infestations or where greater control is required. Apply in a minimum of 200L/ha of water.

Commercial trials have demonstrated that FUSILADE FORTE delivers superior long-term regrowth control compared to Verdict*. For best results, FUSILADE FORTE should be applied to stands that have a large amount of vegetative shoots present at the same stage of growth.

“Double Knock” program to control Couch Grass

In stands that contain a high level of advanced shoots and therefore less actively-growing vegetative shoots, consider a split application of FUSILADE FORTE at 3.3 L/ha in spring and then 3.3L/ha in autumn, or a “double-knock” program in conjunction with SPRAY.SEED.

This strategy involves the application of SPRAY.SEED during early summer to remove the top foliage, followed by an application of FUSILADE FORTE 50 to 70 days later to control any regrowth from the stolons and rhizomes. In general, the second application should be made once the regrowth is about 5 cm high, although timing is less critical with a split application or “double-knock” program.

Application guidelines

- Good coverage is essential for optimal results.
- Apply in 100 to 300L/ha water
- Apply using properly calibrated and maintained equipment
- Use flat fan nozzles to produce fine-medium spray droplets (150 to 300 microns)
- Do not apply under conditions of high temperature, low humidity and when weeds are stressed
- Spray in cool and/or humid conditions (ie Delta T <8°C)
- Avoid drift onto susceptible crops
- Avoid contact with foliage when applying as a directed spray to tree crops.

Application timing

Apply FUSILADE FORTE to young, actively-growing weeds. For best results, spray between the three-leaf weed stage and early tillering (three tillers). Use higher application rates once tillering occurs. Spray before late tillering if a complete kill is desired. Do not treat weeds under poor growth or dormant conditions.



Couch Grass (*Cynodon dactylon*)

Withholding Periods

Avocados, bananas, citrus, pome fruit, stone fruit	Nil
Custard apples, kiwifruit, lychees, mangoes, passionfruit, pawpaw, persimmons, pineapples	14 days
Cucumbers, gherkins, squash, zucchini	21 days
Blueberries, blackcurrants, grapes, lettuce, strawberries, tomatoes	4 weeks
Honeydew, pumpkins, watermelons, onions, broadbeans, green beans, rockmelons	5 weeks
Brussel sprouts, brocc oli, cabbage, cauliflower, peanuts	6 weeks
Peas, carrots	7 weeks
Celery	8 weeks
Potato	10 weeks
Capsicums	11 weeks

Refer to the label directions for other crops and withholding periods for grazing.

 **Fusilade Forte**[®]

syngenta[®]

VISIT WWW.SYNGENTA.COM.AU

For further information please call the Syngenta Technical Product Advice Line on 1800 067 108 or visit www.syngenta.com.au. Product labels and usage directions should be followed for the application of any product referred to in this publication. The information contained in this brochure is believed to be accurate. No responsibility or liability is accepted in respect of this information and those non-excludable conditions implied by and Federal or State legislation or law of a Territory. © Registered trademarks of Syngenta Australia. TN17/071