



200 Pharmacy & Home Medical Supplies
 9133 S. Stony Island Ave.
 Chicago, IL 60617
 Phone: 773-374-4550 Fax: 773-374-4660

Fitting Notes: Patient fittings are between the hours of 9 am – 10:30 am Monday through Saturday. Patient should sleep with legs elevated the night before to ensure proper fit.

Patient:	D.O.B.:
Address: _____ _____	
Phone #: _____	
Diagnosis:	

Number of Pairs:	Number of Refills:
-------------------------	---------------------------

Type of Stocking (Check One)					
Compression:	Calf	Thigh	Pantyhose	Maternity	Thigh w/waist
15-20mmHg	<input type="checkbox"/> L <input type="checkbox"/> R				
20-30mmHg	<input type="checkbox"/> L <input type="checkbox"/> R				
30-40mmHg	<input type="checkbox"/> L <input type="checkbox"/> R				
Other: <input type="checkbox"/> Custom Strength: _____ Type: _____					

Additional Items: <input type="checkbox"/> Latex Free Donning Gloves <input type="checkbox"/> Rubber Gloves <input type="checkbox"/> Donning Device

Physician Name:

Signature:

NPI #:	Phone #:
--------	----------

Compression Level At the ankle	Indications
15-20mmHg	<ul style="list-style-type: none"> Tired, achy legs Occupational edema Leg discomfort from long hours of standing or sitting Prophylaxis of varicosities and swelling during pregnancy Reduce swelling during travel Prophylaxis of DVT during travel
20-30mmHg	<ul style="list-style-type: none"> Heavy, fatigued, tired aching legs Mild edema in lower extremities Mild varicosities with minimal edema Mild varicosities during pregnancy Minimal edema upper extremities Post-procedure of small veins Prevention or management of DVT
30-40mmHg	<ul style="list-style-type: none"> Moderate varicosities with mild to moderate edema (also during pregnancy) Post-procedure of larger veins to maintain treatment Prevention, management or treatment of DVT or Post-Thrombotic Syndrome Superficial phlebitis Mild lymphedema after decongestant therapy to maintain reduction Orthostatic/ postural hypotension Post-fracture, post-traumatic edema
Contraindications	
Contraindications	<ul style="list-style-type: none"> Arterial insufficiency, intermittent claudication, ischemia Uncontrolled congestive heart failure Acute dermatitis, weeping dermatosis, cutaneous sepsis
Cautions	Signs of infections; Skin sensitivities or allergies; Neuropathy, Diabetes; Confinement to bed or non-ambulatory use otherwise prescribed by the physician

MEASURE CIRCUMFERENCE: Ankle: Right _____ Left _____ Calf: Right _____ Left _____ Thigh: Right _____ Left _____ Circle: Inch/ CM