



CHILL PILLS

FOR THE SYMPTOMATIC RELIEF OF PAIN CAUSED BY NAUGHTY STUDENTS, COMPLAINING PARENTS AND ALL OTHER AILMENTS ASSOCIATED WITH TEACHING. EAT ONE AS REQUIRED FOR INSTANT RELIEF. MOST EFFECTIVE WHEN TAKEN WITH ALCOHOL.

**WARNING: HIGHLY ADDICTIVE
KEEP OUT OF REACH OF CHILDREN.**



CHILL PILLS

FOR THE SYMPTOMATIC RELIEF OF PAIN CAUSED BY NAUGHTY STUDENTS, COMPLAINING PARENTS AND ALL OTHER AILMENTS ASSOCIATED WITH TEACHING. EAT ONE AS REQUIRED FOR INSTANT RELIEF. MOST EFFECTIVE WHEN TAKEN WITH ALCOHOL.

**WARNING: HIGHLY ADDICTIVE
KEEP OUT OF REACH OF CHILDREN.**



CHILL PILLS

FOR THE SYMPTOMATIC RELIEF OF PAIN CAUSED BY NAUGHTY STUDENTS, COMPLAINING PARENTS AND ALL OTHER AILMENTS ASSOCIATED WITH TEACHING. EAT ONE AS REQUIRED FOR INSTANT RELIEF. MOST EFFECTIVE WHEN TAKEN WITH ALCOHOL.

**WARNING: HIGHLY ADDICTIVE
KEEP OUT OF REACH OF CHILDREN.**



CHILL PILLS

FOR THE SYMPTOMATIC RELIEF OF PAIN CAUSED BY NAUGHTY STUDENTS, COMPLAINING PARENTS AND ALL OTHER AILMENTS ASSOCIATED WITH TEACHING. EAT ONE AS REQUIRED FOR INSTANT RELIEF. MOST EFFECTIVE WHEN TAKEN WITH ALCOHOL.

**WARNING: HIGHLY ADDICTIVE
KEEP OUT OF REACH OF CHILDREN.**



CHILL PILLS

FOR THE SYMPTOMATIC RELIEF OF PAIN CAUSED BY NAUGHTY STUDENTS, COMPLAINING PARENTS AND ALL OTHER AILMENTS ASSOCIATED WITH TEACHING. EAT ONE AS REQUIRED FOR INSTANT RELIEF. MOST EFFECTIVE WHEN TAKEN WITH ALCOHOL.

**WARNING: HIGHLY ADDICTIVE
KEEP OUT OF REACH OF CHILDREN.**



CHILL PILLS

FOR THE SYMPTOMATIC RELIEF OF PAIN CAUSED BY NAUGHTY STUDENTS, COMPLAINING PARENTS AND ALL OTHER AILMENTS ASSOCIATED WITH TEACHING. EAT ONE AS REQUIRED FOR INSTANT RELIEF. MOST EFFECTIVE WHEN TAKEN WITH ALCOHOL.

**WARNING: HIGHLY ADDICTIVE
KEEP OUT OF REACH OF CHILDREN.**



CHILL PILLS

FOR THE SYMPTOMATIC RELIEF OF PAIN CAUSED BY NAUGHTY STUDENTS, COMPLAINING PARENTS AND ALL OTHER AILMENTS ASSOCIATED WITH TEACHING. EAT ONE AS REQUIRED FOR INSTANT RELIEF. MOST EFFECTIVE WHEN TAKEN WITH ALCOHOL.

**WARNING: HIGHLY ADDICTIVE
KEEP OUT OF REACH OF CHILDREN.**



CHILL PILLS

FOR THE SYMPTOMATIC RELIEF OF PAIN CAUSED BY NAUGHTY STUDENTS, COMPLAINING PARENTS AND ALL OTHER AILMENTS ASSOCIATED WITH TEACHING. EAT ONE AS REQUIRED FOR INSTANT RELIEF. MOST EFFECTIVE WHEN TAKEN WITH ALCOHOL.

**WARNING: HIGHLY ADDICTIVE
KEEP OUT OF REACH OF CHILDREN.**



CHILL PILLS

FOR THE SYMPTOMATIC RELIEF OF PAIN CAUSED BY NAUGHTY STUDENTS, COMPLAINING PARENTS AND ALL OTHER AILMENTS ASSOCIATED WITH TEACHING. EAT ONE AS REQUIRED FOR INSTANT RELIEF. MOST EFFECTIVE WHEN TAKEN WITH ALCOHOL.

**WARNING: HIGHLY ADDICTIVE
KEEP OUT OF REACH OF CHILDREN.**