

Try out [PMC Labs](#) and tell us what you think. [Learn More.](#)



Neuropsychopharmacology

[Neuropsychopharmacology](#). 2013 Dec; 38(13): 2708–2716.

PMCID: PMC3828542

Published online 2013 Aug 21. Prepublished online 2013 Aug 1.

PMID: [23903032](#)

doi: [10.1038/npp.2013.183](#)

Acute Effects of *Scelletium tortuosum* (Zembrin), a Dual 5-HT Reuptake and PDE4 Inhibitor, in the Human Amygdala and its Connection to the Hypothalamus

[David Terburg](#),^{1,2,5,*} [Supriya Syal](#),^{2,3,5} [Lisa A Rosenberger](#),¹ [Sarah Heany](#),² [Nicole Phillips](#),² [Nigel Gericke](#),⁴
[Dan J Stein](#),² and [Jack van Honk](#)^{1,2}

¹Department of Psychology, Utrecht University, Utrecht, The Netherlands

²Department of Psychiatry and Mental Health, University of Cape Town, Cape Town, South Africa

³Department of Psychology, University of Toronto, Toronto, ON, Canada

⁴HG&H Pharmaceuticals (Pty) Ltd, Bryanston, South Africa

^{*}Department of Psychology, Utrecht University, Heidelberglaan 2, Utrecht 3584 CS, The Netherlands, Tel: +31 30 253 3043, Fax: +31 30 253 4511, E-mail: d.terburg@uu.nl

⁵These authors contributed equally to this work.

Received 2013 Feb 26; Revised 2013 Jul 22; Accepted 2013 Jul 22.

[Copyright](#) © 2013 American College of Neuropsychopharmacology

This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivs 3.0 Unported License. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-nd/3.0/>

Abstract

The South African endemic plant *Scelletium tortuosum* has a long history of traditional use as a masticatory and medicine by San and Khoikhoi people and subsequently by European colonial farmers as a psychotropic in tincture form. Over the past decade, the plant has attracted increasing attention for its possible applications in promoting a sense of wellbeing and relieving stress in healthy individuals and for treating clinical anxiety and depression. The pharmacological actions of a standardized extract of the plant (Zembrin) have been reported to be dual PDE4 inhibition and 5-HT reuptake inhibition, a combination that has been argued to offer potential therapeutic advantages. Here we tested the acute effects of Zembrin administration in a pharmacofMRI study focused on anxiety-related activity in the amygdala and its connected neurocircuitry. In a double-blind, placebo-controlled, cross-over design, 16 healthy participants were scanned during performance in a perceptual-load and an emotion-matching task. Amygdala reactivity to fearful faces under low perceptual load conditions was attenuated after a single 25 mg dose of Zembrin. Follow-up connectivity analysis on the emotion-matching task showed that amygdala–hypothalamus coupling was also reduced. These results demonstrate, for the first time, the attenuating effects of *S. tortuosum* on the threat circuitry of the human brain and provide supporting

evidence that the dual 5-HT reuptake inhibition and PDE4 inhibition of this extract might have anxiolytic potential by attenuating subcortical threat responsivity.

Keywords: anxiety, amygdala, hypothalamus, phosphodiesterase, serotonin, *sceletium*

INTRODUCTION

The South African endemic plant *Sceletium tortuosum* (L.) N.E. Br. (synonym *Mesembryanthemum tortuosum* L.), of the succulent family Mesembryanthemaceae, has a long history of traditional use by San and Khoikhoi people as a masticatory and medicine ([Smith et al, 1996](#)) and later by colonial farmers as a psychotropic in tincture form ([Pappe, 1868](#)). Over the past 15 years, the plant has attracted increasing attention for its hypothesized applications in promoting a sense of wellbeing and relieving stress in healthy individuals and for treating anxiety and depression in clinically anxious and depressed patients ([Gericke and Viljoen, 2008](#)). A recent *in vivo* study in rats demonstrated a positive effect of an extract of *S. tortuosum* on restraint-induced anxiety ([Smith, 2011](#)), and a small series of case reports described preliminary evidence for antidepressant and anxiolytic activity in patients suffering from major depression who were treated with tablets of milled *S. tortuosum* raw material ([Gericke, 2001](#)).

The mechanisms of action on the central nervous system (CNS) of a standardized extract of *S. tortuosum* (Zembrin) were recently identified as comprising blockade of the serotonin (5-HT) transporter and selective inhibition of the phosphodiesterase-4 (PDE4) enzyme ([Harvey et al, 2011](#)). 5-HT reuptake inhibitors (SSRIs) are widely used for the treatment of anxiety disorders and depression ([Pringle et al, 2011](#)). However, a combination of SSRIs with a PDE4 inhibitor has been argued to have synergistic therapeutic potential in CNS disorders by providing greater symptomatic efficacy and broader therapeutic utility than either as a drug on its own. In particular, repeated treatment with SSRIs can upregulate PDE4 ([Ye et al, 2000](#)), which in turn reduces the sensitivity to SSRIs in response to long-term treatment, suggesting that dual treatment with SSRIs and PDE4 inhibitors may be a promising approach ([Cashman et al, 2009](#)).

Cyclic nucleotide PDEs comprise a diverse group of enzymes that are important regulators of signal transduction. PDEs are classified into 11 families based on sequence homology, substrates and regulation by modulators. Enzymes in the PDE4 family catalyze the hydrolysis of cyclic AMP (cAMP) and have a critical role in controlling the intracellular concentration of cAMP and increasing phosphorylation of cAMP-response element-binding protein ([Li et al, 2009](#)). PDE4s are found throughout the brain ([Perez-Torres et al, 2000](#)), but their levels are decreased in unmedicated depressed individuals ([Fujita et al, 2012](#)). This reflects a downregulation of the cAMP cascade that can potentially be restored using PDE4 inhibitors ([Duman et al, 1999](#)), an idea which is supported by an increasing number of animal studies that indicate the potential of PDE4 inhibitors for novel treatments of anxiety and depression ([Halene and Siegel, 2007](#); [Li et al, 2009](#)). Indeed, the prototypical PDE4 inhibitor rolipram has been shown in both animal ([Saccomano et al, 1991](#)) and clinical ([Fleischhacker et al, 1992](#)) studies to have antidepressant activity.

A consistent challenge in translating these pre-clinical studies of PDE4s into clinical practice has been the side effects ([Duman et al, 1999](#)); in particular nausea and vomiting ([Rock et al, 2009](#)). The combination of SSRIs and PDE4 inhibitors might therefore not only have synergistic therapeutic potential but may allow for lower PDE4 doses that are better tolerated. Zembrin has been found to be safe and well-tolerated at doses of 8 and 25 mg taken orally once a day in a randomized, double-blind, parallel-group, placebo-controlled clinical trial, supporting the ethnobotanical record of safe use of *S. tortuosum* ([Nell et al, 2013](#)). Here, we report a functional magnetic resonance imaging (fMRI) study that is the first to test the activity of *S. tortuosum* (Zembrin) in the human brain and is thereby also the first study on the effects of a dual PDE4 and 5-HT reuptake inhibitor in humans. The present study was

designed to test the effects of a single administration of Zembrin (25 mg) on anxiety-related activity in the human brain. A reliable and often used index of fear responsivity and anxiety in the human brain is the blood oxygenation level-dependent (BOLD) response to facial threat in the bilateral amygdala as measured with fMRI (Freitas-Ferrari *et al.*, 2010). The amygdala can be considered a hub in the brain's threat system involved in threat detection and subsequent promotion and regulation of defensive reactions in the subcortical threat circuit, particularly the hypothalamus, midbrain, and brainstem (Davis and Whalen, 2001; Walker *et al.*, 2003). Moreover, the amygdala relays threatening information to the cortex and orchestrates, together with cortical structures, the regulation of fear and anxiety (Kim *et al.*, 2011; Terburg *et al.*, 2012). Given this diverse role in threat processing, we used two fMRI experiments to independently assess anxiety-related amygdala activation (Bishop *et al.*, 2007) and functional connectivity of the amygdala with the threat circuit of the brain (Fisher *et al.*, 2011; Hariri *et al.*, 2002; Kirsch *et al.*, 2005; van Wingen *et al.*, 2010; van Wingen *et al.*, 2008). As the amygdala is involved in the regulation of fear responsivity in the subcortical threat circuit (hypothalamus and midbrain), as well as with controlling mechanisms in the prefrontal cortex (PFC), we specifically focused the latter analysis on these areas. We hypothesized that administration of Zembrin would reduce anxiety-related amygdala activity as well as functional connectivity of the amygdala with the cortical and/or subcortical threat circuit.

MATERIALS AND METHODS

Investigational Product

The active ingredient of Zembrin is a standardized and characterized aqueous ethanolic (purified water 30% V/V and ethanol 70% V/V, spray-dried onto a maltodextrin carrier) extract of the above ground material of a cultivated traditionally used selection of the South African plant *S. tortuosum*. Figure 1 shows the chemical structures of the four active alkaloids (mesembrenone, mesembrenol, mesembranol, and mesembrine), which were quantified by high pressure liquid chromatography (HPLC) analysis against validated analytical reference standards (Harvey *et al.*, 2011). See Supplementary Information for a detailed description.




Figure 1

Structures of the four main mesembrine alkaloids that are quantified by HPLC to define the alkaloid content and composition of the extract of *Sceletium tortuosum* (Zembrin).

Each active opaque white size-1 hardshell gelatin capsule, lot number NG022, contained 25 mg of Zembrin together with inert excipients, equivalent to 50 mg of dry raw above-ground *S. tortuosum* plant material. Placebo capsules, lot number NG009, contained only inert excipients. Both placebo and active capsules were identical in size, color, opacity, and odor and were manufactured to GMP by Archon Vitamin Corporation, New Jersey, USA. This study was approved by the Clinical Trials Committee of the Medicines Control Council of South Africa.

Participants and Procedure

Sixteen (eight female) right-handed, healthy undergraduate students (aged 18–21 years, mean=19.2 years) from the University of Cape Town participated in this study. All participants were right-handed, medication-free, had no history of neurological disease or psychopathology (screened using Mini-International Neuropsychiatric Interviews ([Sheehan et al, 1998](#))), had normal or corrected-to-normal vision, and received transportation and remuneration for research time (research credit or ZAR200).

Each participant was scanned once after Zembrin and once after placebo administration, separated by 5–9 days (mean=6.8 days), and counterbalanced across participants. Researchers remained blind to administration conditions. Each scan session took place promptly at 2 h post administration. Participants were monitored for the entire duration between administration and scanning and were asked to report any mood changes or side effects (see [Supplementary Table S1](#)). After the final session, participants were asked to identify the session that involved administration of the investigational product, which was correctly identified by nine participants, which is not significantly (one-tailed binomial test; $P=0.30$) above chance level. Response to this question, as well as gender, were entered as covariates in all fMRI analyses but did not change the results.

Perceptual-Load Task (PLT)

The PLT was adapted from [Bishop et al \(2007\)](#). In each trial, participants saw a string of six letters superimposed on a task-irrelevant unfamiliar face for 200 ms. Participants were required to decide whether the letter string contained an 'X' or 'N'. The letter string comprised six Xs or Ns in the low-load condition, and a single target letter and five non-target letters (HKMWZ) arranged in a semi-random order in the high-load condition. Target letters appeared in each position equally often. Participants responded by a single key press using button boxes held in each hand. 'N' and 'X' responses were counterbalanced for hand across participants. The faces were of eight different individuals with fearful and neutral expressions ([Ekman and Friesen, 1976](#)). Data were acquired for a total of 12 blocks in the high-load and 12 blocks in the low-load condition wherein each block consisted of 4 trials. Load stayed constant across all trials within a block, but emotion varied on a trial-by-trial basis such that each block contained two fearful and two neutral faces presented in a random order. Within blocks, the interstimulus interval was randomly jittered with a mean of 4.5 s and a minimum of 3 s.

Previous research has shown that the PLT can assess state-anxiety-related amygdala activity. Fearful compared with neutral faces induce threat responses, which under high-load conditions are inhibited in favor of task performance ([Bishop, 2008](#); [Lavie, 1995](#)). In low-load conditions, however, inhibition of the threat response is not necessary for correct performance, and amygdala reactivity to the fearful compared with neutral faces correlates with state anxiety ([Bishop et al, 2007](#)).

Emotion-Matching Task (EMT)

The EMT was adapted from [Hariri et al \(2002\)](#). Four blocks of an emotion-matching task were interleaved with five blocks of a sensorimotor-control task involving shape matching. During each 5-s trial in the emotion condition, participants saw a target face on top and two probe faces at the bottom.

Participants' task was to match the emotion of the face on top with one of the other faces using a left or a right key press (corresponding to the position of the face). During the sensorimotor-control shape-matching trials, participants saw oval shapes in a configuration analogous to the faces and were instructed to match the orientation of the oval (vertical or horizontal ellipses) on top to the ovals on the bottom. The emotional faces expressed anger or fear ([Tottenham et al, 2009](#)), and luminosity of the oval shapes was matched to that of the faces. Each block consisted of six trials, and blocks were interspersed by a 2-s presentation of instructions for the next block (ie 'match emotion' or 'match shape').

The EMT contrasts emotion with shape-matching blocks and is thereby a reliable amygdala activator ([Hariri et al, 2002](#)), which is stable over multiple sessions ([Manuck et al, 2007](#)). Recent evidence also indicates that the EMT provides a valid measure of pharmacological and state-related changes in connectivity patterns of the amygdala with cortical ([van Wingen et al, 2010](#); [van Wingen et al, 2008](#)) as well as subcortical structures ([Kirsch et al, 2005](#)).

MRI Acquisition

Participants were scanned in a 3-T Allegra Siemens head scanner (VA25 platform, Siemens Medical Systems GmbH, Erlangen, Germany), using a four-channel receiver. A high-resolution T1-weighted anatomical volume (magnetization-prepared rapid gradient echo) was collected in between the two functional runs using the following parameters: TR/TE=2.53 s/6.6 ms, FA=7°, FOV=256 × 256 × 171, and voxel-size=1.0 × 1.0 × 1.3 mm. Functional volumes (echo-planar imaging) were obtained with 36 interleaved-ascending axial slices, EPI factor=64, TR/TE=2 s/27 ms, flip angle=70°, FOV=224 × 224 × 145 mm, and voxel-size=3.5 mm isotropic. Before the start of each task, five volumes were acquired and discarded to allow for stabilization of the MR signal, and 255 (PLT) and 150 (EMT) functional volumes were acquired for data-analysis. Task-stimuli were back projected onto a screen visible via an angled mirror attached to the helmet coil. Each task was explained to the participants in detail 15 min before the scan, and they completed practice rounds in the scanner before the start of each task.

MRI Analysis

All MR images were processed and analyzed using SPM8 (www.fil.ion.ucl.ac.uk/spm; Wellcome Department of Imaging Neuroscience, London, UK). Functional volumes were realigned to the first image in the series. The anatomical volume was coregistered to the mean realigned image and transformed to standard space (Montreal Neurological Institute; MNI) using the unified model as implemented in SPM8. As the PLT is an event-related design, these images were slice-time corrected with the 19th slice as reference. Functional volumes were transformed to MNI space using the parameters from the normalization of the anatomical volume, resampled to 2 mm isotropic, and smoothed with an 8-mm full-width half-maximum Gaussian smoothing kernel.

For both tasks, we applied a general linear model (GLM) to the single-subject data using a canonical hemodynamic response function, with the realignment parameters and a vector with the timing of the button presses convolved with a hemodynamic function as regressors of no interest, and a high-pass filter cutoff of 128 s. For both tasks, mean BOLD signal change from the bilateral amygdala were subsequently extracted for all individual contrast maps using the MarsBar toolbox (<http://marsbar.sourceforge.net>) and tested in repeated measures ANOVAs. In the GLM for the PLT, each trial was modeled with a box-car function for the duration of the stimulus and separate contrast maps were computed for each condition: LOAD (high, low), and EMOTION (fear, neutral). Contrast maps were entered in a full-factorial design and tested for effects of LOAD (high>low), EMOTION (fear>neutral), DRUG (Zembrin *versus* placebo), and their interactions (whole-brain family wise error (FWE) corrected $P<0.05$, and extend-threshold 20 voxels).


In the GLM for the EMT, each block was modeled with a 30-s box-car function and separate contrast maps were computed for emotion and shape matching. A full-factorial design was used to test for main effects of CONDITION (emotion>shape), DRUG (Zembrin *versus* placebo) and their interaction (whole-brain FWE-corrected $P<0.05$, extend-threshold 20 voxels). Furthermore, we performed a region of interest (ROI) analysis to assess activity in the threat circuit (ventro-medial PFC/orbitofrontal cortex (vmPFC/OFC), amygdala, hypothalamus, and midbrain) with a ROI volume FWE-corrected $P<0.05$. ROIs were based on the Wake Forest University pickatlas (<http://www.fmri.wfubmc.edu/download.htm>), except for the hypothalamus, which was constructed by drawing an 8-mm sphere around the central coordinate (MNI: 0–2–13) of hypothalamus subregions recently described by Baroncini *et al* (2012).

The peak voxels of the contrast of interest indexing anxiety-related amygdala reactivity in the PLT were used for connectivity analysis in the EMT targeting the threat circuit. For each participant, time course (first eigenvariate) was extracted from the spheres of 6 mm around these peak voxels, and mean corrected for general task effects. These were used to assess connectivity for the whole task and both conditions separately by entering them in the original GLM, as well as for psycho-physiological interaction (PPI) analysis as implemented in SPM8 to assess differences between conditions. The resulting contrast maps were entered in the separate full-factorial models to test for DRUG effects in functional connectivity of the bilateral amygdala with the threat circuit (ROI-corrected).

RESULTS

PLT

One participant performed below chance level (47% correct), which was >3 SDs below average (85%, $SD=4$), and was therefore excluded from analysis. A $2 \times 2 \times 2$ repeated-measures ANOVA on the error rates with DRUG, LOAD, and EMOTION as within-subject factors revealed a significant effect of LOAD ($F(1,14)=194.2$, $P<0.00001$) but with no other main effects or interactions (all $P_s>0.27$). Reaction times revealed the same pattern: a significant effect of LOAD ($F(1,14)=57.4$, $P<0.00001$) but with no other main effects or interactions (all $P_s>0.28$; see [Figure 2a](#)).



[Open in a separate window](#)

Figure 2

(a) Mean error rates and reaction times on the PLT (error bars represent SEM). (b) Increased BOLD signal in the high-load compared with low-load condition of the PLT plotted in a glass brain (whole-brain FWE-corrected, $P < 0.05$).

In line with earlier studies with the PLT, whole-brain FWE-corrected positive main effects of LOAD (high>low) were observed bilaterally in the parietal and occipital cortices, anterior insula, precentral,

and midfrontal PFC, and dorsal anterior cingulate cortex (see [Table 1](#) and [Figure 2b](#)). No other main effects or interactions were observed using this statistical threshold.

Table 1**Significant Clusters from the fMRI Analyses**

[Open in a separate window](#)


Abbreviations: ACC, anterior cingulate cortex; OFC, orbitofrontal cortex; PFC, prefrontal cortex; SMC, supplementary motor cortex.

All effects are whole-brain FWE-corrected with an extent threshold of 20 voxels unless stated otherwise. Coordinates refer to MNI space.

^aFWE-corrected for ROI volume.

^bFWE-corrected cluster-threshold with uncorrected cluster-defining threshold $P < 0.001$.

A $2 \times 2 \times 2$ repeated-measures ANOVA on the extracted mean signal in the bilateral amygdala, with DRUG, LOAD, and EMOTION as within-subject factors, showed a significant three-way interaction ($F(1,14)=6.81$, $P=0.028$, partial $\eta^2=0.30$), as well as a main effect of LOAD ($F(1,14)=13.02$, $P=0.003$, partial- $\eta^2=0.48$). As hypothesized, the three-way interaction was due to a significant attenuation of amygdala activity after Zembrin administration on the contrast fear>neutral in the low-load condition ($F(1,14)=6.90$, $P=0.020$, partial $\eta^2=0.33$, see [Figure 3a](#) for the associated contrast estimates), which was non-significant in the high-load condition ($F(1,14)=0.003$, $P=0.956$).



[Open in a separate window](#)

Figure 3

(a) Mean extracted BOLD signal change from the bilateral amygdala on low-load fearful and neutral trials during the PLT (error bars represent SEM). (b) Increased BOLD signal for placebo compared with Zembrin on low-load fearful compared with neutral trials during the PLT in the bilateral amygdala plotted on a template brain ($P < 0.01$, uncorrected for visualization).

Finally, to determine the peak voxels of this effect for further connectivity analyses we performed an ROI analysis on this particular contrast, which confirmed the significantly reduced amygdala reactivity for the left (58 voxels, $P = 0.008$, peak voxel $-26\ 0, -18$, $P = 0.004$) and right (13 voxels, $P = 0.025$, peak voxel $24\ 4, -20$, $P = 0.010$) amygdala (see [Table 1](#) and [Figure 3b](#)).


Finally, a more liberal whole-brain analysis (FWE-corrected cluster threshold $P < 0.05$, cluster-defining threshold $P < 0.001$ uncorrected) showed a three-way interaction within the dorsal striatum, precuneus, and bilateral postcentral gyrus. Follow-up contrasts established that the effects in the dorsal striatum and

precuneus emerged from the same contrast as the effect in the amygdala (placebo>Zembrin (low load: fear>neutral), see [Table 1](#)), but the effect in the bilateral postcentral gyrus was not significant in any of the simpler contrasts.

EMT

The participant excluded from the PLT also scored near to chance level (56% correct) on the EMT, thus was also excluded here. No differences were found in performance for CONDITION or DRUG (emotion matching: 92% correct, SD=9, shape matching: 93%, SD=13).

Whole-brain FWE-corrected positive main effects of CONDITION (emotion>shape) were observed bilaterally in the fusiform and temporal cortices, cerebellum, frontal inferior PFC, amygdala, and hippocampus (see [Table 1](#) and [Figure 4a](#)). No main effect of DRUG or interaction was observed using this or a more liberal (FWE-corrected cluster threshold $P<0.05$, cluster-defining threshold $P<0.001$ uncorrected) statistical threshold. ROI analysis on the threat circuit shows reliable activation in the midbrain, hypothalamus, amygdala, and vmPFC/OFC on the emotion>shape-matching contrast (see [Table 1](#)), again with no main or interaction effects of DRUG. A 2×2 repeated-measures ANOVA on the extracted mean signal in the bilateral amygdala, with DRUG and CONDITION as within-subject factors, revealed a marginally significant interaction effect ($F(1,14)=3.13$, $P=0.099$, partial $\eta^2=0.18$) indicating marginally stronger amygdala activity on the emotion>shape-matching contrast in the Zembrin (percentage of signal change; 0.61, SD=0.53) compared with placebo (0.38, SD=0.23) condition.



[Open in a separate window](#)

Figure 4

(a) Increased BOLD signal in the emotion-matching compared with shape-matching blocks during the EMT plotted in a glass brain (whole-brain *FWE*-corrected, $P < 0.05$). (b) Stronger functional connectivity of the bilateral amygdala (seed region) with the hypothalamus for placebo compared with Zemrin during the EMT plotted on a template brain ($P < 0.01$, uncorrected for visualization), and parameter estimates of the mean regression coefficient for this cluster, showing that the positive correlation between amygdala and hypothalamus in the placebo condition was reduced after Zemrin administration (error bars represent SEM).

Finally, time course analysis based on the peak voxels in the amygdala as identified in the PLT analysis revealed that the amygdala showed overall decreased functional connectivity with the hypothalamus (8 voxels, $P=0.034$) after Zembrin compared with placebo administration (see [Figure 4b](#)) but no effects in midbrain or vmPFC/OFC. Further specification of this effect revealed that the reduced amygdala–hypothalamus connectivity was marginally larger in the emotion compared with the shape-matching condition (PPI analysis, 13 voxels, $P=0.068$). This was further confirmed by a significant hypothalamus cluster in the emotion-matching condition (47 voxels, $P=0.014$), which did not reach significance in the shape-matching condition (peak voxel, uncorrected $P=0.246$). As ROI correction in larger ROIs such as vmPFC/OFC and midbrain might obscure local effects, we also used a more liberal uncorrected threshold of $P<0.001$ for these analyses, but this did not reveal any other significant clusters.

DISCUSSION

In this pharmaco-fMRI study, we tested the effects of a single administration of Zembrin, a *S. tortuosum* extract that can be characterized as a dual PDE4 and 5-HT reuptake inhibitor ([Harvey et al, 2011](#)). Using two independent fMRI designs, we show that Zembrin compared with placebo administration reduces anxiety-related ([Bishop et al, 2007](#)) amygdala reactivity and attenuates amygdala–hypothalamus coupling. These results not only have importance as the first evidence on the dampening effects of Zembrin on the brain's threat system but support further work on the clinical applicability of dual PDE4 and 5-HT reuptake inhibitors for the treatment of anxiety disorders and depression.

Anxiety is characterized by hyper-responsivity to mild threats. The task we used to assess amygdala reactivity directly taps into this mechanism as mild threats, ie fearful faces, had to be ignored in favor of a cognitive task, ie letter detection. Although clinically anxious individuals show such threat interference in general ([Bar-Haim et al, 2007](#)), healthy individuals can successfully ignore task-irrelevant stimuli under high task demands ([Lavie, 1995](#)). In line with this, the amygdala is activated by unattended facial fear, particularly under low perceptual load conditions ([Silvert et al, 2007](#)), which is further increased by state anxiety ([Bishop et al, 2007](#)). In the present study, this amygdala reactivity to facial fear under low-load conditions was significantly attenuated after Zembrin administration. Although we did not observe general subjective mood effects, the effects noted here arguably reflect a reduction of responsivity of the threat system related to the anxiolytic properties of *S. tortuosum* ([Gericke, 2001](#); [Gericke and Viljoen, 2008](#); [Smith et al, 1996](#)). It must furthermore be noted that in contrast to the previous study with the PLT, the behavioral data did not show the expected EMOTION \times LOAD interaction. Most likely this is due to a ceiling effect in performance in the low-load condition, as the error rate in the current study ($\sim 3\%$) was only half of the error rate in the earlier study ($\sim 6\%$). Crucially, this does not affect our interpretation of the fMRI effects as behavioral performance was previously shown to be independent from state anxiety and its relation to the amygdala's BOLD signal ([Bishop et al, 2007](#)).

Attenuation of threat responsivity, as measured with the BOLD signal, can result from reduced basal threat responding, as well as increased cognitive control over these threat responses. Our results are in line with the former as during the matching of threatening facial expressions a decrease in functional connectivity of the amygdala and hypothalamus was observed after Zembrin administration. This effect was marginally stronger when contrasted with the shape-matching control condition, whereas cortical areas showed no such main or interaction effects even when using a less stringent statistical threshold. The amygdala and hypothalamus are connected through the bed nucleus of the stria terminalis, which is heavily involved in anxiety-related behaviors ([Walker et al, 2003](#)), thus arguably, this effect reflects anxiolytic actions of Zembrin on basal threat responding in this subcortical threat circuit.

Combined, these results provide support for the anxiolytic potential of dual 5-HT reuptake and PDE4

inhibitors (Harvey *et al*, 2011). This combination has previously been argued to have synergistic potential in the treatment of anxiety and depression (Cashman *et al*, 2009), with the added value that lower doses can be used to reduce side effects (Duman *et al*, 1999; Rock *et al*, 2009). Although we cannot separate these pharmacological actions in the present results, the effects seem to compare well with the literature on both the mechanisms (Harmer *et al*, 2009) (Harmer *et al*, 2009). Acute SSRI administration has been shown to increase amygdala reactivity in tasks that required explicit evaluation of threatening facial expression (Bigos *et al*, 2008) but generally reduces amygdala reactivity to unattended facial expressions of threat (Del-Ben *et al*, 2005; Murphy *et al*, 2009). The first effect is marginally supported in the present data and is usually interpreted as an effect of increased agitation and increased sensitivity for social information in general after first use of SSRIs (Anderson *et al*, 2008; Harmer *et al*, 2009), which normally disappears after long-term administration (Pringle *et al*, 2011). The second effect has been interpreted as the underlying mechanism for the anxiolytic properties of SSRIs in long-term administrations, which are often described in terms of functional coupling of prefrontal areas with the amygdala (Fisher *et al*, 2011; McCabe and Mishor, 2011; Pringle *et al*, 2011). In the present data, this is further supported as the reduction of amygdala reactivity was particularly observed in a contrast that has previously been linked to state anxiety (Bishop *et al*, 2007). Finally, although we did not find evidence for the effects on PFC–amygdala coupling in the present study, these effects might be observed in future studies of long-term Zembrin administration.

FMRI studies on PDE4 inhibitors are, to our knowledge, currently not available, but PDE4 inhibitors can be specifically linked to anxiolytic functions in the subcortical threat circuit. Although PDE4s can be found in cortical as well as subcortical areas (Korff *et al*, 2009), especially the sub-types PDE4B and PDE4D (Perez-Torres *et al*, 2000), which are particular the targets of Zembrin (Harvey *et al*, 2011), can be found in the olfactory nucleus, ventral striatum, hypothalamus, and midbrain of mice (Cherry and Davis, 1999). Moreover, PDE4B has been argued to be associated with acute effects on the noradrenergic stress system (Cherry and Davis, 1999), has been linked to panic disorder (Otowa *et al*, 2011), and the prototypical PDE4 inhibitor rolipram has been shown to affect the amygdala (Rutten *et al*, 2009; Werenicz *et al*, 2012), possibly through its enhancing effect on Fos-like immunoreactivity (Bureau *et al*, 2006). Future fMRI research should establish how these effects are reflected in amygdala reactivity to threat, but taken together these data do suggest that PDE4 inhibitors target the subcortical threat circuit, which is further supported by our results.

In sum, we show here that the dual 5-HT reuptake and PDE4 inhibitor Zembrin reduces amygdala reactivity to unattended facial fear and also decouples amygdala–hypothalamus connectivity. These results support the potential anxiolytic actions of Zembrin and provide a foundation for exploring the clinical potential of dual PDE4 and 5-HT reuptake inhibitors for the treatment of anxiety disorders and depression.

FUNDING AND DISCLOSURE

This study was made possible by funds from H.L. Hall and Sons, Ltd, South Africa and the Netherlands Organization of Scientific Research (Brain and Cognition Grant 056-24-010). NG is the Director, Medical and Scientific Affairs, of HG&H Pharmaceuticals (Pty) Ltd, the company that has developed the extract of *Sceletium tortuosum* (Zembrin). DJS has received research grants and/or consultancy honoraria from Abbott, Astrazeneca, Eli-Lilly, GlaxoSmithKline, Jazz Pharmaceuticals, Johnson & Johnson, Lundbeck, Orion, Pfizer, Pharmacia, Roche, Servier, Solvay, Sumitomo, Takeda, Tikvah, and Wyeth. The other authors declare no conflict of interest.

Footnotes

[Supplementary Information](#) accompanies the paper on the Neuropsychopharmacology website

(<http://www.nature.com/npp>)

Supplementary Material

Supplementary Information

[Click here for additional data file.](#) (235K, pdf)

References

- Anderson IM, McKie S, Elliott R, Williams SR, Deakin JF. Assessing human 5-HT function *in vivo* with pharmacMRI. *Neuropharmacology*. 2008;**55**:1029–1037. [[PubMed](#)] [[Google Scholar](#)]
- Bar-Haim Y, Lamy D, Pergamin L, Bakermans-Kranenburg MJ, van IJzendoorn MH. Threat-related attentional bias in anxious and nonanxious individuals: a meta-analytic study. *Psychol Bull*. 2007;**133**:1–24. [[PubMed](#)] [[Google Scholar](#)]
- Baroncini M, Jissendi P, Balland E, Besson P, Pruvo J-P, Francke J-P, et al. MRI atlas of the human hypothalamus. *Neuroimage*. 2012;**59**:168–180. [[PubMed](#)] [[Google Scholar](#)]
- Bigos KL, Pollock BG, Aizenstein HJ, Fisher PM, Bies RR, Hariri AR. Acute 5-HT reuptake blockade potentiates human amygdala reactivity. *Neuropsychopharmacology*. 2008;**33**:3221–3225. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
- Bishop SJ. Neural mechanisms underlying selective attention to threat. *Ann NY Acad Sci*. 2008;**1129**:141–152. [[PubMed](#)] [[Google Scholar](#)]
- Bishop SJ, Jenkins R, Lawrence AD. Neural processing of fearful faces: effects of anxiety are gated by perceptual capacity limitations. *Cereb Cortex*. 2007;**17**:1595–1603. [[PubMed](#)] [[Google Scholar](#)]
- Bureau Y, Handa M, Zhu Y, Laliberte F, Moore CS, Liu S, et al. Neuroanatomical and pharmacological assessment of Fos expression induced in the rat brain by the phosphodiesterase-4 inhibitor 6-(4-pyridylmethyl)-8-(3-nitrophenyl) quinoline. *Neuropharmacology*. 2006;**51**:974–985. [[PubMed](#)] [[Google Scholar](#)]
- Cashman JR, Voelker T, Johnson R, Janowsky A. Stereoselective inhibition of serotonin re-uptake and phosphodiesterase by dual inhibitors as potential agents for depression. *Bioorg Med Chem*. 2009;**17**:337–343. [[PubMed](#)] [[Google Scholar](#)]
- Cherry JA, Davis RL. Cyclic AMP phosphodiesterases are localized in regions of the mouse brain associated with reinforcement, movement, and affect. *J Comp Neurol*. 1999;**407**:287–301. [[PubMed](#)] [[Google Scholar](#)]
- Davis M, Whalen PJ. The amygdala: vigilance and emotion. *Mol Psychiatry*. 2001;**6**:13–34. [[PubMed](#)] [[Google Scholar](#)]
- Del-Ben CM, Deakin JF, McKie S, Delvai NA, Williams SR, Elliott R, et al. The effect of citalopram pretreatment on neuronal responses to neuropsychological tasks in normal volunteers: an fMRI study. *Neuropsychopharmacology*. 2005;**30**:1724–1734. [[PubMed](#)] [[Google Scholar](#)]
- Duman RS, Malberg J, Thome J. Neural plasticity to stress and antidepressant treatment. *Biol Psychiatry*. 1999;**46**:1181–1191. [[PubMed](#)] [[Google Scholar](#)]
- Ekman P, Friesen W. *Pictures of Facial Affect*. Consulting Psychologist Press: Palo Alto, CA, USA; 1976. [[Google Scholar](#)]
- Fisher PM, Price JC, Meltzer CC, Moses-Kolko EL, Becker C, Berga SL, et al. Medial prefrontal cortex serotonin 1A and 2A receptor binding interacts to predict threat-related amygdala reactivity. *Biol Mood Anxiety Disord*. 2011;**1**:2. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
- Fleischhacker WW, Hinterhuber H, Bauer H, Pflug B, Berner P, Simhandl C, et al. A multicenter double-blind study of three different doses of the new cAMP-phosphodiesterase inhibitor rolipram

- in patients with major depressive disorder. *Neuropsychobiology*. 1992;**26**:59–64. [[PubMed](#)] [[Google Scholar](#)]
- Freitas-Ferrari MC, Hallak JE, Trzesniak C, Filho AS, Machado-de-Sousa JP, Chagas MH, et al. Neuroimaging in social anxiety disorder: a systematic review of the literature. *Prog Neuropsychopharmacol Bol Psychiatry*. 2010;**34**:565–580. [[PubMed](#)] [[Google Scholar](#)]
- Fujita M, Hines CS, Zoghbi SS, Mallinger AG, Dickstein LP, Liow JS, et al. Downregulation of brain phosphodiesterase type IV measured with 11C-(R)-rolipram positron emission tomography in major depressive disorder. *Biol Psychiatry*. 2012;**72**:548–554. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
- Gericke N. Clinical application of selected South African medicinal plants. *Aust J Med Herbalism*. 2001;**13**:3. [[Google Scholar](#)]
- Gericke N, Viljoen AM. *Sceletium*—a review update. *J Ethnopharmacol*. 2008;**119**:653–663. [[PubMed](#)] [[Google Scholar](#)]
- Halene TB, Siegel SJ. PDE inhibitors in psychiatry—future options for dementia, depression and schizophrenia. *Drug Discov Today*. 2007;**12**:870–878. [[PubMed](#)] [[Google Scholar](#)]
- Hariri AR, Mattay VS, Tessitore A, Kolachana B, Fera F, Goldman D, et al. Serotonin transporter genetic variation and the response of the human amygdala. *Science*. 2002;**297**:400–403. [[PubMed](#)] [[Google Scholar](#)]
- Harmer CJ, Goodwin GM, Cowen PJ. Why do antidepressants take so long to work? A cognitive neuropsychological model of antidepressant drug action. *Br J Psychiatry*. 2009;**195**:102–108. [[PubMed](#)] [[Google Scholar](#)]
- Harvey AL, Young LC, Viljoen AM, Gericke NP. Pharmacological actions of the South African medicinal and functional food plant *Sceletium tortuosum* and its principal alkaloids. *J Ethnopharmacol*. 2011;**137**:1124–1129. [[PubMed](#)] [[Google Scholar](#)]
- Kim MJ, Loucks RA, Palmer AL, Brown AC, Solomon KM, Marchante AN, et al. The structural and functional connectivity of the amygdala: from normal emotion to pathological anxiety. *Behav Brain Res*. 2011;**223**:403–410. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
- Kirsch P, Esslinger C, Chen Q, Mier D, Lis S, Siddhanti S, et al. Oxytocin modulates neural circuitry for social cognition and fear in humans. *J Neurosci*. 2005;**25**:11489–11493. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
- Korff S, Stein DJ, Harvey BH. Cortico-striatal cyclic AMP-phosphodiesterase-4 signalling and stereotypy in the deer mouse: attenuation after chronic fluoxetine treatment. *Pharmacol Biochem Behav*. 2009;**92**:514–520. [[PubMed](#)] [[Google Scholar](#)]
- Lavie N. Perceptual load as a necessary condition for selective attention. *J Exp Psychol Hum Percept Perform*. 1995;**21**:451–468. [[PubMed](#)] [[Google Scholar](#)]
- Li YF, Huang Y, Amsdell SL, Xiao L, O'Donnell JM, Zhang HT. Antidepressant- and anxiolytic-like effects of the phosphodiesterase-4 inhibitor rolipram on behavior depend on cyclic AMP response element binding protein-mediated neurogenesis in the hippocampus. *Neuropsychopharmacology*. 2009;**34**:2404–2419. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
- Manuck SB, Brown SM, Forbes EE, Hariri AR. Temporal stability of individual differences in amygdala reactivity. *AJ Psychiatry*. 2007;**164**:1613–1614. [[PubMed](#)] [[Google Scholar](#)]
- McCabe C, Mishor Z. Antidepressant medications reduce subcortical-cortical resting-state functional connectivity in healthy volunteers. *Neuroimage*. 2011;**57**:1317–1323. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
- Murphy SE, Norbury R, O'Sullivan U, Cowen PJ, Harmer CJ. Effect of a single dose of citalopram on amygdala response to emotional faces. *Br J Psychiatry*. 2009;**194**:535–540. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
- Nell H, Siebert M, Chellan P, Gericke N. 2013A **randomized, double-blind, parallel-group, placebo-**

- controlled trial of extract *Sceletium tortuosum* (Zembrin®) in healthy adults** *J Altern Complement Med*(in press). [[PubMed](#)]
- Otowa T, Kawamura Y, Sugaya N, Yoshida E, Shimada T, Liu X, et al. Association study of PDE4B with panic disorder in the Japanese population. *Prog Neuropsychopharmacol Biol Psychiatry*. 2011;**35**:545–549. [[PubMed](#)] [[Google Scholar](#)]
- Pappe L. 1868 *Florae Capensis Medicae Prodromus. An Enumeration of South African Indigenous Plants used as Remedies by the Colonists of the Cape of Good Hope* 3rd edn. W Brittain: Cape Town, South Africa [[Google Scholar](#)]
- Perez-Torres S, Miro X, Palacios JM, Cortes R, Puigdomenech P, Mengod G. Phosphodiesterase type 4 isozymes expression in human brain examined by in situ hybridization histochemistry and [³H]rolipram binding autoradiography. Comparison with monkey and rat brain. *J Chem Neuroanat*. 2000;**20**:349–374. [[PubMed](#)] [[Google Scholar](#)]
- Pringle A, Browning M, Cowen PJ, Harmer CJ. A cognitive neuropsychological model of antidepressant drug action. *Prog Neuropsychopharmacol Biol Psychiatry*. 2011;**35**:1586–1592. [[PubMed](#)] [[Google Scholar](#)]
- Rock EM, Benzaquen J, Limebeer CL, Parker LA. Potential of the rat model of conditioned gaping to detect nausea produced by rolipram, a phosphodiesterase-4 (PDE4) inhibitor. *Pharmacol Biochem Behav*. 2009;**91**:537–541. [[PubMed](#)] [[Google Scholar](#)]
- Rutten K, Van Donkelaar EL, Ferrington L, Blokland A, Bollen E, Steinbusch HWM, et al. Phosphodiesterase inhibitors enhance object memory independent of cerebral blood flow and glucose utilization in rats. *Neuropsychopharmacology*. 2009;**34**:1914–1925. [[PubMed](#)] [[Google Scholar](#)]
- Saccomano NA, Vinick FJ, Koe BK, Nielsen JA, Whalen WM, Meltz M, et al. Calcium-independent phosphodiesterase inhibitors as putative antidepressants: [3-(bicycloalkoxy)-4-methoxyphenyl]-2-imidazolidinones. *J Med Chem*. 1991;**34**:291–298. [[PubMed](#)] [[Google Scholar](#)]
- Sheehan DV, Lecrubier Y, Sheehan KH, Amorim P, Janavs J, Weiller E, et al. The Mini-International Neuropsychiatric Interview (M.I.N.I.): the development and validation of a structured diagnostic psychiatric interview for DSM-IV and ICD-10. *J Clin Psychiatry*. 1998;**59** (Suppl 20):22–33. [[PubMed](#)] [[Google Scholar](#)]
- Silvert L, Lepsien J, Fragopanagos N, Goolsby B, Kiss M, Taylor JG, et al. Influence of attentional demands on the processing of emotional facial expressions in the amygdala. *Neuroimage*. 2007;**38**:357–366. [[PubMed](#)] [[Google Scholar](#)]
- Smith C. The effects of *Sceletium tortuosum* in an *in vivo* model of psychological stress. *J Ethnopharmacol*. 2011;**133**:31–36. [[PubMed](#)] [[Google Scholar](#)]
- Smith MT, Crouch NR, Gericke N, Hirst M. Psychoactive constituents of the genus *Sceletium* N.E.Br. and other Mesembryanthemaceae: a review. *J Ethnopharmacol*. 1996;**50**:119–130. [[PubMed](#)] [[Google Scholar](#)]
- Terburg D, Morgan BE, Montoya ER, Hooge IT, Thornton HB, Hariri AR, et al. Hyper-vigilance for fear after basolateral amygdala damage in humans. *Translational Psychiatry*. 2012;**2**:e115. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
- Tottenham N, Tanaka JW, Leon AC, McCarry T, Nurse M, Hare TA, et al. The NimStim set of facial expressions: judgments from untrained research participants. *Psychiatry Res*. 2009;**168**:242–249. [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
- van Wingen G, Mattern C, Verkes RJ, Buitelaar J, Fernandez G. Testosterone reduces amygdala-orbitofrontal cortex coupling. *Psychoneuroendocrinology*. 2010;**35**:105–113. [[PubMed](#)] [[Google Scholar](#)]
- van Wingen GA, van Broekhoven F, Verkes RJ, Petersson KM, Bäckström T, Buitelaar JK, et al. Progesterone selectively increases amygdala reactivity in women. *Mol Psychiatry*.

2008;**13**:325–333. [[PubMed](#)] [[Google Scholar](#)]

Walker DL, Toufexis DJ, Davis M. Role of the bed nucleus of the stria terminalis versus the amygdala in fear, stress, and anxiety. *Eur J Pharmacol.* 2003;**463**:199–216. [[PubMed](#)] [[Google Scholar](#)]

Werenicz A, Christoff RR, Blank M, Jobim PF, Pedroso TR, Reolon GK, et al. Administration of the phosphodiesterase type 4 inhibitor rolipram into the amygdala at a specific time interval after learning increases recognition memory persistence. *Learn Mem.* 2012;**19**:495–498. [[PubMed](#)] [[Google Scholar](#)]

Ye Y, Jackson K, O'Donnell JM. Effects of repeated antidepressant treatment of type 4A phosphodiesterase (PDE4A) in rat brain. *J Neurochem.* 2000;**74**:1257–1262. [[PubMed](#)] [[Google Scholar](#)]

Articles from *Neuropsychopharmacology* are provided here courtesy of **Nature Publishing Group**
