

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/49704819>

Antimicrobial potential of some plant extracts against *Candida* species

Article in *Brazilian journal of biology = Revista brasleira de biologia* · November 2010

DOI: 10.1590/S1519-69842010000500022 · Source: PubMed

CITATIONS

112

READS

229

7 authors, including:



José Francisco Höfling

University of Campinas

143 PUBLICATIONS 2,543 CITATIONS

[SEE PROFILE](#)



Paula Cristina Anibal

University of Campinas

18 PUBLICATIONS 282 CITATIONS

[SEE PROFILE](#)



Gustavo Obando-Pereda

Universidad Católica de Santa María

23 PUBLICATIONS 270 CITATIONS

[SEE PROFILE](#)



Vivian Fernandes Furetti

Centro Universitário Herminio Ometto (UNIARARAS)

14 PUBLICATIONS 574 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Efeitos antimicrobianos de óleos essenciais sobre microrganismos iniciadores do biofilme dental [View project](#)



Effect of Five Common Hospital Decontamination Agents and Disinfectants on Common Hospital Acquired Infections Causing Bacterial Species [View project](#)

Antimicrobial potential of some plant extracts against *Candida* species

Höfling, JF.^{a,*}, Anibal, PC.^{a,b}, Obando-Pereda, GA.^a,
Peixoto, IAT.^{a,c}, Furletti, VF.^{a,c}, Foglio, MA.^b and Gonçalves, RB.^a

^aÁrea de Microbiologia e Imunologia, Departamento de Diagnóstico Oral, Faculdade de Odontologia de Piracicaba, Universidade Estadual de Campinas – Unicamp, Av. Limeira, 901, CP 052, CEP 13414-900, Piracicaba, SP, Brazil

^bCentro Pluridisciplinar de Pesquisas Químicas, Biológicas e Agrícolas, Departamento de Fitoquímica, Universidade Estadual de Campinas – Unicamp, Campinas, SP, Brazil

^cCentro Pluridisciplinar de Pesquisas Químicas, Biológicas e Agrícolas, Departamento de Microbiologia, Universidade Estadual de Campinas – Unicamp, Campinas, SP, Brazil

*e-mail: hofling@fop.unicamp.br

Received April 16, 2009 – Accepted January 12, 2010 – Distributed November 30, 2010

Abstract

The increase in the resistance to antimicrobial drugs in use has attracted the attention of the scientific community, and medicinal plants have been extensively studied as alternative agents for the prevention of infections. The *Candida* genus yeast can become an opportunistic pathogen causing disease in immunosuppressive hosts. The purpose of this study was to evaluate dichloromethane and methanol extracts from *Mentha piperita*, *Rosmarinus officinalis*, *Arrabidaea chica*, *Tabebuia avellanedae*, *Punica granatum* and *Syzygium cumini* against *Candida* species through the analysis of Minimum Inhibitory Concentration (MIC). Results presented activity of these extracts against *Candida* species, especially the methanol extract.

Keywords: *Candida*, plant extracts activity, antifungal activity, antimicrobial agents.

Potencial antimicrobiano de extratos de plantas na inibição de leveduras do gênero *Candida*

Resumo

Devido ao aumento da resistência aos antimicrobianos em uso, as plantas medicinais têm sido intensamente estudadas como agentes alternativos para a prevenção de doenças e infecções. A levedura do gênero *Candida*, por ser um patógeno oportunista, tem sua virulência aumentada ao adquirir resistência aos antifúngicos, desencadeando doenças, principalmente em hospedeiros imunossuprimidos. O propósito deste trabalho foi avaliar os extratos diclorometano e metanol das plantas *Mentha piperita*, *Rosmarinus officinalis*, *Arrabidaea chica*, *Tabebuia avellanedae*, *Punica granatum* e *Syzygium cumini* contra linhagens do gênero *Candida* através dos testes de Concentração Inibitória Mínima (CIM). Os resultados demonstraram atividade dos extratos sobre as espécies de *Candida*, particularmente o extrato metanol.

Palavras-chave: *Candida*, atividade de extratos de plantas, atividade antifúngica, agentes antimicrobianos.

1. Introduction

Medicinal plants and corresponding preparations have been used for a wide range of purposes and for many centuries people have been trying to treat diseases as well as alleviate symptoms by using different plant extracts and formulations (Cowan, 1999). Plants such as *Mentha piperita* (Iskan et al., 2002), *Rosmarinus officinalis* (Nascimento et al., 2000), *Arrabidaea chica* (Taylor, 1998), *Tabebuia avellanedae* (Machado et al., 2003), *Punica granatum* (Nascimento et al., 2000; Duraipandiyan et al., 2006) and *Syzygium cumini* (Chandrasekaram and Venkatesalu, 2004) have been used due to their antimicrobial properties.

In the 2000-2006 period, approximately 50% of new chemical molecules extracted from natural products demonstrated their importance for the development of drugs

in the treatment of infectious diseases (Newman and Cragg, 2007). The choice of an appropriate antifungal treatment is important, though limited to a few licensed agents (Provine and Hadley, 2000). The increasing resistance of microorganisms to antimicrobial drugs in use has attracted the attention of the scientific community regarding the search for new cost-effective drugs of natural or synthetic origin (Pai et al., 2004).

Members of the *Candida* genus are commensal aerobic microorganisms, which are part of the indigenous microbial flora in humans and can be found in the oral cavity and the digestive as well as vaginal tracts (Odds, 1988). They may become opportunistic pathogens causing a number of diseases in immunosuppressive hosts (De Repentigny et al., 2004),

mainly HIV-positive patients (Erköse and Erturan, 2007). A particular characteristic of *Candida* is its ability to invade oral mucosa tissues by the development of hyphae, which adhere to tissue surfaces and lead to inflammation (Ellepola and Samaranyake, 2001).

Although the colonisation by *Candida albicans* is common and causes severe injuries in immunocompromised patients, other *Candida* species have been isolated from such patients, healthy patients and children with Down's syndrome, such as *C. glabrata*, *C. krusei*, *C. tropicalis*, *C. lusitaniae*, *C. parapsilosis*, *C. guilliermondii* (Erköse and Erturan, 2007; Höfling et al., 2001; Ribeiro et al., 2006; Rodrigues et al., 2004) and *C. dubliniensis*. These yeasts were recognised as the source of mucosal infection in HIV-positive patients and regarded as a significant cause of infections in humans, such as abdominal infections and fungemia (Erköse and Erturan, 2007).

The purpose of this study was to evaluate the potential activity of extracts from six selected plants against ten *Candida* species.

2. Material and Methods

Selected plants: the species *Mentha piperita* L. (leaves, voucher no. UEC 1253), *Arrabidaea chica* (Bonpl.) B. Verl. (leaves, voucher no. UEC 1254), *Rosmarinus officinalis* L. (leaves, voucher no. UEC 1264), *Syzygium cumini* L. (seeds, voucher no. UEC 143724), *Punica granatum* L. (fruit, voucher no. UEC 143723) and *Tabebuia avellanedae* Lor. ex Griseb (bark, voucher no. UEC 1256) were collected from the experimental field of the Research Centre for Chemistry, Biology and Agriculture, State University of Campinas (CPQBA/UNICAMP), Brazil. Voucher specimens were deposited at the Herbarium of the State University of Campinas (UEC) and were identified by Professor Jorge Yoshio Tamashiro, Ph.D.

Preparation of extracts: Each selected plant (5 g) was extracted with 600 mL of dichloromethane (Labsynth PA) with a Dispersive Extratur (Quimis® Q-252-28 model) and

filtered thereafter. The vegetal residue was reextracted with methanol (Labsynth PA). Solvents were evaporated under reduced pressure and dried using a rotary evaporator (Buchi® R-200 model). Crude extracts were monitored by chromatography in Thin Layer in silicagel chromatoplaques (60 F254 Merck 1.05554). The dried plant extracts were dissolved in Tween20 (Labsynth) and sterile distilled water, filtered through a 0.22-µm membrane filter (TPP) and stored at 4 °C until further use. The extracts were diluted at the moment of use with the concentration ranging from 1 to 0.001 mg/mL.

Microorganisms: The test organisms used were *Candida albicans* CBS-562, *C. dubliniensis* CBS-7987, *C. parapsilosis* CBS-604, *C. tropicalis* CBS-94, *C. guilliermondii* CBS-566, *C. utilis* CBS-5609, *C. krusei* CBS-573, *C. lusitaniae* B-06, *C. glabrata* B-07, *C. rugosa* B-12, proceeding from the Microbiology and Immunology Laboratory, at the School of Dentistry of Piracicaba (FOP/UNICAMP).

Screening for antimicrobial activities: Preparation of inoculum for susceptibility tests was carried out by microdilution as set forth by the CLSI's M27-A2 recommendation protocol (CLSI, 2002). The yeasts were grown overnight at 37 °C in Sabouraud Dextrose Agar (Merck) plates, and inocula for the assays were prepared by diluting scrape cell mass in 0.85% NaCl solution, adjusted to 0.5 Mc Farland scale and confirmed by spectrophotometric reading at 625 nm. Cell suspensions were finally diluted to 5.0×10^3 CFU/mL. 50 µL of diluted extract were added in 50 µL of RPMI-1640 in microplates (96 wells) + 100 µL of microorganisms. After that, the samples were incubated at 37 °C for 24-48 hours, in duplicate. Fluconazol was used as control standard in concentrations ranging from 64-0.125 µg/mL. The microplates were incubated at 37 °C for 48 hours.

3. Results

All *Candida* species in the in vitro test presented sensitivity to the plant extracts in use, though not for all extracts (Table 1).

Table 1. In vitro antifungal activity of dichloromethane and methanol extracts (mg/mL).

Microorganisms	Dichloromethane extracts						Methanol extracts					
	AC	MP	SC	TA	RO	PG	AC	MP	SC	TA	RO	PG
<i>C. albicans</i>	0.015	R	0.03	R	0.007	0.001	R	0.007	0.001	0.003	0.001	0.003
<i>C. dubliniensis</i>	0.03	R	0.06	R	0.015	0.001	R	0.007	0.001	0.007	0.007	0.003
<i>C. parapsilosis</i>	0.015	R	0.007	R	0.03	0.001	R	0.003	0.001	0.003	0.003	0.003
<i>C. tropicalis</i>	0.015	R	0.03	R	0.015	0.001	R	0.003	0.001	0.015	0.003	0.001
<i>C. guilliermondii</i>	0.015	R	0.007	R	0.007	0.001	R	0.001	0.001	0.001	0.003	0.001
<i>C. utilis</i>	R	R	0.001	R	R	0.001	R	0.007	0.001	0.003	R	0.003
<i>C. krusei</i>	0.007	R	0.03	0.06	0.003	0.001	R	0.007	0.001	0.007	0.001	0.001
<i>C. lusitaniae</i>	0.007	R	0.03	R	0.007	0.001	R	0.007	0.001	0.003	0.001	0.001
<i>C. glabrata</i>	R	R	0.001	R	R	0.001	R	0.003	0.001	0.001	R	0.003
<i>C. rugosa</i>	0.007	R	0.03	R	0.001	0.001	R	0.003	0.001	0.003	0.001	0.003

AC, *Arrabidaea chica*; MP, *Mentha piperita*; SC, *Syzygium cumini*; TA, *Tabebuia avellanedae*; RO, *Rosmarinus officinalis*; PG, *Punica granatum*; R, Resistant.

4. Discussion

Because of the increasing development of drug resistance to human pathogens and the appearance of undesirable effects of certain antifungal agents, the search for new antimicrobial agents is of great concern today (Phongpaichit et al., 2005). A multidisciplinary approach to drug discovery, involving the generation of truly novel molecular diversity from natural product sources combined with total and combinatorial synthetic methodologies, and including the manipulation of biosynthetic pathways, provides the best solution to the current productivity crisis facing the scientific community engaged in drug discovery and development (Newman and Cragg, 2007).

Our results revealed a strong activity of *Punica granatum*, *Syzygium cumini* and *Rosmarinus officinalis* (dichloromethane and methanol extracts), *Arrabidaea chica* (dichloromethane extract), *Mentha piperita* and *Tabebuia avellanedae* (methanol extract), with MIC varying from 0.06 to 0.001 mg/mL; and no or less activity of *Arrabidaea chica* (methanol extract), *Mentha piperita* and *Tabebuia avellanedae* (dichloromethane extract). The methanol extract of *A. chica* showed activity against yeasts within 24 hours (data not shown). However, these yeasts showed resistance to this extract after 48 hours. Among the 10 yeasts used in this study, the most resistant were *C. glabrata* and *C. utilis*, and *C. krusei* and *C. guilliermondii* were the most sensitive strains to the tested extracts.

Data obtained from other studies demonstrated positive results for these plants as well. Duraipandiyar et al. (2006) observed inhibitory effects of *P. granatum*, while Nascimento et al. (2000) showed activity of extracts of *P. granatum*, *S. cumini* and *R. officinalis* against *Candida albicans*. Chandrasekaram and Venkatesalu (2004) reported effectiveness of methanol extract of *S. cumini* and Portillo et al. (2001) found positive results for the dichloromethane extract of *T. avellanedae* against yeast *C. albicans*, thus validating our results. On the other hand, Duarte et al. (2005) obtained moderate activity testing *M. piperita* oil against *C. albicans* while Duraipandiyar et al. (2006), Portillo et al. (2001) and Dulger and Gonuz (2004) observed resistance of the extracts of *S. cumini*, *T. avellanedae* (methanol extract) and *R. officinalis* (ethanolic extract), respectively. Although these results are not in accordance with ours, there may be many other contributing factors, such as the seasonal period when plants are collected.

However, some compounds like tannins (*P. granatum*, *A. chica*, *S. cumini* and *R. officinalis*), anthocyanins (*A. chica*), flavonoids (*A. chica* and *R. officinalis*), naphthoquinones (*T. avellanedae*), menthol and menthone (*M. piperita*) are known to have antimicrobial properties against microorganisms (Alcerito et al., 2002; Cordeiro et al., 2006; Gershon, 1975; Hussein et al., 1997; Iscan et al., 2002; Wagner et al., 1989). Components like tannins found in plants may act on the cell membrane and precipitin proteins (Nawwar et al., 1994), which turns out to be a target for studies.

The findings presented in this paper indicate that the extracts obtained from the selected plants had anti-*Candida*

activity. *Punica granatum* and *Syzygium cumini* extracts exerted strong antifungal activity and can be a source for the development of new therapeutic agents, as they inhibited the growth of *Candida*. Subsequently, bioguided fractionation shall be conducted on *Punica granatum* to identify the active compounds against *Candida* genus species.

Acknowledgements — This study was supported by the Research Foundation for the State of São Paulo (Fundação de Apoio à Pesquisa do Estado de São Paulo – FAPESP) and the Fund for Education, Research and Extension (Fundo de Apoio ao Ensino, Pesquisa e Extensão – FAEPEX).

References

- ALCERITO, T., BARBO, FE., NEGRI, G., SANTOS, DYAC., MEDA, CI., YOUNG, MCM., CHÁVEZ, D. and BLATT, CTT., 2002. Folicular epicuticular wax of *Arrabidaea brachypoda*: flavonoids and antifungal activity. *Biochemical Systematics and Ecology*, vol. 30, p. 677-683.
- CHANDRASEKARAN, M. and VENKATESALU, V., 2004. Antibacterial and antifungal activity of *Syzygium jambolonum* seeds. *Journal of Ethnopharmacology*, vol. 91, p. 105-108.
- Clinical and Laboratory Standards Institute - CLSI, 2002. *Método de Referência para Testes de Diluição em Caldo para Determinação da Sensibilidade a Terapia Antifúngica das Leveduras*: Norma Aprovada. 2. ed. Wayne, Pa: National Committee for Clinical Laboratory Standards. Norma M27-A2.
- CORDEIRO, CHG., SACRAMENTO, LVS., CORRÊA, MA., PIZZOLITTO, AC. and BAUAB, TM., 2006. Análise farmacognóstica e atividade antibacteriana de extratos vegetais empregados em formulação para a higiene bucal. *Revista Brasileira de Ciências Farmacêuticas*, vol. 42., no. 3, p. 395-404.
- COWAN, MM., 1999. Plant product as antimicrobial agents. *Clinical Microbiology Reviews*, vol. 12, no. 4, p. 564-582.
- DE REPENTIGNY, L., LEWANDOWSKI, D. and JOLICOEUR, P., 2004. Immunopathogenesis of oropharyngeal candidiasis in human immunodeficiency virus infection. *Clinical Microbiology Reviews*, vol. 17, p. 729-759.
- DUARTE, MCT., FIGUEIRA, GM., SARTORATTO, A., REHDER, VLG. and DELARMELENA, C., 2005. Anti-*Candida* activity of Brazilian medicinal plants. *Journal of Ethnopharmacology*, vol. 97, p. 305-311.
- DULGER, B. and GONUZ, A., 2004. Antimicrobial activity of certain plants used in Turkish traditional Medicine. *Asian Journal of Plant Sciences*, vol. 3, no. 1, p. 104-107.
- DURAI PANDIYAN, V., AYYANAR, M. and IGNACIMTHU, S., 2006. Antimicrobial activity of some ethnomedicinal plants used by Paliyar tribe from Tamil Nadu, India. *BMC Complementary and Alternative Medicine*, vol. 6, no. 35, p. 1-7.
- ELLEPOLA, ANB. and SAMARANAYAKE, LP., 2001. Adjunctive use of chlorhexidine in oral candidoses: a review. *Oral Diseases*, vol. 7, p. 11-17.
- ERKÖSE, G. and ERTURAN, Z., 2007. Oral *Candida* colonization of human immunodeficiency virus infected subjects in Turkey and its relation with viral load and CD4⁺ T-lymphocyte count. *Mycoses*, vol. 50, p. 485-490.

- GERSHON, H., 1975. Fungitoxicity of 1,4-naphthoquinones to *Candida albicans*. *Canadian Journal of Microbiology*, vol. 21, p. 1317-21.
- HÖFLING, JF., ROSA, EAR., PEREIRA, CV., BORIOLLO, FG. and RODRIGUES, JAO., 2001. Differentiation and numerical analysis of oral yeasts based on SDS-Page profiles. Influence of the culture media on the whole-cell protein extracts. *Brazilian Journal of Biology*, vol. 61, no. 3, p. 507-516.
- HUSSEIN, SAM., BARAKAT, HH., MERFORT, I. and NAWWAR, MAM., 1997. Tannins from the leaves of *Punica granatum*. *Phytochemistry*, vol. 45, p. 819-23.
- ISCAN, G., KIRIMER, N., KÜRKCÜOĞLU, M., BASER, KHC. and DEMIRCI, F., 2002. Antimicrobial screening of *Mentha piperita* essential oils. *Journal of Agricultural and Food Chemistry*, vol. 50, p. 3943-3946.
- MACHADO, TB., PINTO, AV., PINTO, MCFR., LEAL, ICR., SILVA, MG., AMARAL, ACF., KUSTER, RM. and NETTO-DOSSANTOS, KR., 2003. In vitro activity of Brazilian medicinal plants, naturally occurring naphthoquinones and their analogues, against methicillin-resistant *Staphylococcus aureus*. *International Journal of Antimicrobial Agents*, vol. 21, p. 279-284.
- NASCIMENTO, GGF., LOCATELLI, J., and FREITAS, PC. and SILVA, GL., 2000. Antibacterial activity of plant extracts and phytochemicals on antibiotic-resistant bacteria. *Brazilian Journal of Microbiology*, vol. 31, p. 247-256.
- NAWWAR, MAN., HUSSEIN, SAM. and MERFORT, I., 1994. NMR spectral analysis of polyphenols from *Punica granatum*. *Phytochemistry*, vol. 36, p. 793-798.
- NEWMAN, DJ. and CRAGG, GM., 2007. Natural products as source of new drugs over the last 25 years. *Journal of Natural Products*, vol. 70, p. 461-477.
- ODDS, FC., 1988. *Candida and candidoses: a review and bibliography*. 2nd ed. London: Basilière Tindall.
- PAI, MR., ACHARYA, LD. and UDUPA, N., 2004. Evaluation of antiplaque activity of *Azadirachta indica* leaf extract gel – a 6-week clinical study. *Journal of Ethnopharmacology*, vol. 90, p. 99-103.
- PHONGPAICHIT, S., SUBHADHIRASAKUL, S. and WATTANAPIROMSAKUL, C., 2005. Antifungal activities of extracts from Thai medicinal plants against opportunistic fungal pathogens associated with AIDS patients. *Mycoses*, vol. 48, p. 333-338.
- PORTILLO, A., VILA, R., FREIXA, B., ADZET, T. and CAÑIGUERAL, S., 2001. Antifungal activity of Paraguayan plants used in traditional medicine. *Journal of Ethnopharmacology*, vol. 76, p. 93-98.
- PROVINE, H. and HADLEY, S., 2000. Preliminary evaluation of a semisolid agar antifungal susceptibility test for yeasts and molds. *Journal of Clinical Microbiology*, vol. 2, p. 537-541.
- RIBEIRO, EL., SCROFERNEKER, ML., CAVALHAES, MS., CAMPOS, CC., NAGATO, GM., SOUZA, NA., FERREIRA, WM., CARDOSO, CG., DIAS, SMS., PIMENTA, FC. and TOLEDO, AO., 2006. Phenotypic aspects of oral strains of *Candida albicans* in children with Dow's syndrome. *Brazilian Journal of Biology*, vol. 66, no. 3, p. 939-944.
- RODRIGUES, JAO., HÖFLING, JF., TAVARES, FCA., DUARTE, KMR., GONÇALVES, RB. and AZEVEDO, RA., 2004. Evaluation of biochemical and serological methods to identify and clustering yeast cells of oral *Candida* species by CHROMagar test, SDS-Page and ELISA. *Brazilian Journal of Biology*, vol. 64, no. 2, p. 317-326.
- TAYLOR, L., 1998. *Herbal secrets of the rainforest*. Rocklin: Prima Publishing, Inc.
- WAGNER, H., KREHER, B., LOTTER, H., HAMBURGER, MO. and CORDELL, GA., 1989. Structure determination of new isomeric naphtha [2,3-b] furan-4,9-diones from *Tabebuia avellanedae* by the selective-INEPT technique. *Helvetica Chimica Acta*, vol. 72, p. 659-67.