


# Melatonin interferes with COVID-19 at several distinct ROS-related steps



ELSEVIER



## Review Article

# Melatonin interferes with COVID-19 at several distinct ROS-related steps

## Highlights

- SARS-CoV-2 infection generates overwhelming levels of neutrophil myeloperoxidase
- Hypochlorous acid and other reactive oxygen species destroy tetrapyrrole rings
- This causes nitric oxide, oxygen, and vitamin B12 deficiencies; markers of COVID-19
- Melatonin inhibits myeloperoxidase activity and scavenges reactive oxygen

species

•

Melatonin supplements can prevent pathophysiological consequences of COVID-19 disease

## Abstract

Recent studies have shown a correlation between COVID-19, caused by [severe acute respiratory syndrome coronavirus 2](#) (SARS-CoV-2) infection, and the distinct, exaggerated immune response titled “cytokine storm”. This immune response leads to excessive production and accumulation of [reactive oxygen species](#) (ROS) that cause clinical signs characteristic of COVID-19 such as decreased oxygen [saturation](#), alteration of hemoglobin properties, decreased nitric oxide (NO) bioavailability, [vasoconstriction](#), elevated cytokines, cardiac and/or renal injury, enhanced D-dimer, leukocytosis, and an increased neutrophil to lymphocyte ratio. Particularly, neutrophil [myeloperoxidase](#) (MPO) is thought to be especially abundant and, as a result, contributes substantially to [oxidative stress](#) and the pathophysiology of COVID-19. Conversely, [melatonin](#), a potent MPO inhibitor, has been noted for its anti-inflammatory, anti-oxidative, anti-apoptotic, and neuroprotective actions. Melatonin has been proposed as a safe therapeutic agent for COVID-19 recently, having been given with a US Food and Drug Administration emergency authorized cocktail, REGEN-COV2, for management of COVID-19 progression. This review distinctly highlights both how the destructive interactions of HOCl with [tetrapyrrole](#) rings may contribute to oxygen deficiency and hypoxia, [vitamin B12 deficiency](#), NO deficiency, increased oxidative stress, and sleep disturbance, as well as how melatonin acts to prevent these events, thereby improving COVID-19 prognosis.

- [< Previous](#)
- [Next >](#)

## Keywords

COVID-19

Melatonin

Mammalian peroxidase

Reactive oxygen species

Inflammation

Hemoprotein

Sleep

## Abbreviations

ROS

reactive oxygen species

NETs

neutrophil extracellular traps

iNOS

inducible nitric oxide synthase


MMP

matrix metalloproteinase

## 1. Introduction

In the severe stages of COVID-19 caused by [severe acute respiratory syndrome coronavirus 2](#) (SARS-COV-2) infection, signs of hypoxemia accompanied by decreased response to oxygen therapy supplementation are present, regardless of the overall preservation of [lung mechanics](#) [1,2]. Certain reactions of the immune system during COVID-19 are associated with what has been characterized as a “cytokine storm” in which overwhelming amounts of [reactive oxygen species](#) (ROS) including [superoxide](#) ( $O_2^{\bullet-}$ ), hydrogen peroxide ( $H_2O_2$ ), [hydroxyl radical](#) ( $\bullet OH$ ), and [peroxynitrite](#) ( $ONOO^-$ ), as seen in other inflammatory disorders, are produced ([Fig. 1](#)). Excessive activation of neutrophils caused by the overactive immune response generates [myeloperoxidase](#) (MPO) and contributes to the formation of neutrophil extracellular traps (NETs). NETs are mesh-like DNA fibers that are cast out from neutrophils in response to stimuli including microorganisms, microbial products, and [chemokines](#) [3]. The formation of NETs is highly dependent on, and most commonly triggered by, the protein kinase C (PKC) activator [phorbol myristate acetate](#) (PMA), a potent neutrophil stimulus that activates  $O_2^{\bullet-}$  anion producing NADPH-oxidase [4]. PKC activation by PMA triggers the production of intragranular ROS. MPO activity through the generation of HOCl, along with other ROS, works to destroy assaulting pathogens, but may lead to tissue damage when HOCl is produced in excess [[5], [6], [7]] ([Fig. 1](#)). HOCl exists in approximate equilibrium with [hypochlorite](#) ion ( $^-OCl$ ) at normal body pH 7.4 (pKa 7.59); however only the uncharged molecule can easily penetrate the cell membrane, most likely by [passive diffusion](#) through [porins](#), proteins on the cell membrane that form channels [8,9]. MPO can be attached to extracellular NETs, be released at the plasma membrane due to inappropriate trafficking control, bind to the [glycocalyx](#) of several cell types and further increase neutrophil recruitment, or otherwise leak from the phagolysosome. All of these events can cause additional catalysis of HOCl [10]. In addition, the production of excessive amounts of ROS can interfere with the catalytic site of many [hemoprotein](#) model compounds, ultimately contributing to malfunction and/or destruction of hemoglobin and red blood cells, [nitric oxide synthase](#) (NOS), [catalase](#), [corrin](#) rings such as in [vitamin B12](#), zinc-Cys clusters, and sulfur-sulfur bonds, deficiencies of which are characteristic of COVID-19 [8,[11], [12], [13], [14], [15], [16]]. HOCl might also modulate other important pathological mechanisms in COVID-19 including the release of free iron (Fe(II)) from HOCl mediated hemoprotein destruction, which reacts with HOCl directly or through the

[Fenton reaction](#) generating  $\cdot\text{OH}$ . Additionally, direct NO consumption by the near [diffusion rate](#) reaction of nitric oxide (NO) with  $\text{O}_2^{\cdot-}$  generates  $\text{ONOO}^-$  and leads to [vasoconstriction](#) [17,18]. Understanding these mechanisms will help the development of therapeutic strategies to combat SARS-CoV-2 and other related inflammatory disorders.



1. [Download : Download high-res image \(418KB\)](#)
2. [Download : Download full-size image](#)

Fig. 1. The proposed pathway of SARS-CoV-2 infection associated with uncontrolled immune response and [cytokine storm](#) generating [myeloperoxidase](#) (MPO) and reactive oxygen species (ROS) resulting in decreased NO and [vasoconstriction](#), decreased  $\text{O}_2$  and hypoxia, and [vitamin B12 deficiency](#) through [tetrapyrrole](#) ring destruction mechanism, and the prevention by [melatonin](#).

Therapeutic treatment with [melatonin](#) has shown satisfactory outcomes in COVID-19 through potential preventive mechanisms, anticipated reduction of symptom severity, and possible reduction of immune pathology [[19], [20], [21]]. Ramlall et al., found that exposure to melatonin in intubated COVID-19 patients was associated with a more positive disease outcome [22]. It is thought that melatonin may not directly promote the host defense system against the virus but rather increase the tolerance of the host to the virus such as by suppression of inflammatory or apoptotic species e.g. nuclear factor kappa-light-chain-enhancer of activated B cells (NF- $\kappa$ B) p65 or tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ) [19,20]. Furthermore, melatonin may be able to suppress [cytokine storm](#) effects on activation of monocyte/macrophage and inflammatory [cytokine release](#), as well as inhibition of mammalian peroxidase chlorinating activity (Fig. 1) [21,23,24]. In addition, it has also been suggested that melatonin has preventive effects against SARS-CoV-2 infection by acting as an indirect inhibitor of angiotensin-converting enzyme 2 (ACE2) receptor coupling to SARS-CoV-2. Finally, this report highlights that melatonin can be an adjuvant to enhance the effectiveness of anti-SARS-CoV-2 vaccines [25]. The beneficial effects of melatonin in ameliorating COVID-19

symptoms have been attributed to its multiple roles as an [antioxidant](#), anti-inflammatory, and immunoregulatory agent, as well as its ability to counteract circadian disruption, all of which allow melatonin to combat multiple diseases including metabolic syndrome, cancer, diabetes, infertility, and cardiovascular diseases [[20,26](#)].

Melatonin is produced in the mitochondria of all organs in addition to its synthesis in the pineal gland; although, it is thought that the initial function was as an antioxidant during evolution with circadian rhythm effects evolving thereafter [[27](#)]. With the preventive benefits as stated above, melatonin is a stable and inexpensive over-the-counter drug, generally safe and effective for both short- and long-term use, displays a very high safety profile, can be easily synthesized when needed in large quantities, and can be easily self-administered [[10](#), [20](#), [28](#), [29](#), [30](#)]. Due to this convenience, Reiter et al., have suggested that melatonin treatment should be considered for prophylactic use, treatment alone, or treatment in combination with other drugs to fight SARS-CoV-2 infection [[20](#)]. The recommended pharmacological oral dose is 100–400 mg as an adjunct once a day immediately after contact with an infected SARS-CoV-2 individual or at the start of experiencing symptoms [[20](#)].

This review is an extension and continuation of our previous work [[1,31](#)], and differentiates from other reports on the subject, in that it discusses the ability of HOCl, the final product of MPO, to mediate the destruction of [tetrapyrrole](#) rings and the subsequent effects such as oxygen deficiency and hypoxia, [vitamin B12 deficiency](#), NO deficiency, and the release of corresponding free metal. Also, this review unveils the mechanisms by which melatonin's actions as an antioxidant can be beneficial in preventing these effects thereby reducing dire outcomes through prevention of the pathophysiological consequences of SARS-CoV-2.

## 2. HOCl-associated hemoprotein heme destruction

As an important part of the first immune response, the activation of neutrophils results in the production of ROS; however, when ROS release is excessive due to excess stimulation of neutrophil activity, the result is often physiological damage such as cellular mitochondria poisoning, [oxidative phosphorylation uncoupling](#), and [lipid peroxidation](#). Of commonly appearing ROS, HOCl and H<sub>2</sub>O<sub>2</sub> are long-lived compared to O<sub>2</sub><sup>•-</sup> and •OH, and they can further generate •OH upon reaction with free [metal ions](#), a process that may be enhanced by O<sub>2</sub><sup>•-</sup>. HOCl can also directly react with O<sub>2</sub><sup>•-</sup> to generate •OH [[32](#)]. In turn, •OH can react with organic molecules at rates that approach diffusion-limited and contribute to [hemoprotein](#) heme modification [[33](#)].

[MPO](#) catalyzes H<sub>2</sub>O<sub>2</sub>-dependent [oxidation](#) of Cl<sup>-</sup> through the formation of a ferryl [π cation radical](#) (E – Fe(IV) = O<sup>•+π</sup>) intermediate, compound I, to form HOCl [[5,6](#)]. Alternatively, Compound I can oxidize several organic and inorganic substrates (e.g., NO and nitrite (NO<sub>2</sub><sup>-</sup>) to form nitrosonium cation (NO<sup>+</sup>) and [nitrogen dioxide](#) (NO<sub>2</sub>), respectively) during which the heme undergoes two sequential one e<sup>-</sup> reduction steps producing compound II (MPO-Fe(IV) = O) and MPO-Fe(III), respectively [[34,35](#)]. Normal MPO levels range from 18 to 39 ng/ml in human

plasma, with a significant increase up to 55 ng/ml and 287 ng/ml in inflammatory diseases, making MPO an efficient [oxidative stress](#) biomarker. HOCl accumulated by neutrophils have been reported at concentrations up to 25–50 mM/h, but is difficult to measure due to the amount of neutrophils present, MPO released, and H<sub>2</sub>O<sub>2</sub> availability [[10,36](#)]. Enhanced levels of HOCl are sufficient to substantially damage biomolecules through processes such as thiol oxidation, [chloramine](#) formation, aromatic [chlorination](#), and [aldehyde](#) generation [[37](#)]. Furthermore, HOCl can alter the biological function of hemoproteins through oxidation or destruction of heme (iron-protoporphyrin IX), an essential co-factor involved in multiple [biological processes](#) such as oxygen transport and storage, [electron transfer](#), drug and [steroid metabolism](#), [signal transduction](#), and [microRNA](#) processing [[1,11,12,15,38](#)]. The destructive actions of accumulated HOCl, or other ROS, on hemoprotein heme moieties and other biomolecules could explain some features of COVID-19 and other similar respiratory disorders such as acute lung injury (ALI)/ acute respiratory distress syndrome (ARDS) [[1,39](#)], [[40](#)], [[41](#)]]. These features include substantial hypoxemia not sufficiently explained by alveolar-parenchymal pathology, [zinc deficiency](#), [vitamin B12](#) deficiency, microvascular injury, thromboembolism, pulmonary hypertension, and damaged hemoglobin and red blood cell function accompanied by relative unresponsiveness to O<sub>2</sub> supplementation.

In the absence of an HOCl scavenger, there is a complex and interdependent relationship between levels of self-generated HOCl and MPO [catalytic activity](#) during steady-state catalysis. Enhanced self-generated HOCl in the milieu regulates MPO catalytic activity by heme degradation and subsequent free iron release, a process that is attenuated by HOCl scavengers (e.g. methionine) [[38](#)]. These studies clearly showed that self-generated HOCl may serve as a ligand for MPO, leading to catalytic inhibition and formation of a MPO–Fe(III)–OCl complex, which subsequently sets the stage for MPO heme destruction and free iron release. The direct reaction between HOCl with MPO–Fe(III) is fast and occurs with a second-order rate constant of  $2 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$  (pH 7) [[42](#)]. Key targets for HOCl-induced protein damage are various protein side chains, particularly those that are sulfur-containing. Rate constants reported by Davies and Pattison for reaction of HOCl and different amino acids range from  $26 \text{ M}^{-1} \text{ s}^{-1}$  to  $3.6 \times 10^8 \text{ M}^{-1} \text{ s}^{-1}$ , a rate comparable to that of the direct reaction of HOCl with MPO [[10,43,44](#)]. Regardless of the variation in the rate constants among different amino acids, the final products for these reactions have been detected even for the lowest rate reaction, i.e. halogenated [tyrosine](#) has been proposed as a biomarker for detection of HOCl-induced protein damage [[45](#)]. Furthermore, an initial rate of  $8 \times 10^3 \text{ M}^{-1} \text{ s}^{-1}$  was found for the binding of HOCl to hemoglobin, further supporting the possibility of HOCl-mediated hemoglobin damage in a biological setting [[12,43](#)]. Similarly, an initial rate constant of  $2 \times 10^3 \text{ M}^{-1} \text{ s}^{-1}$  was found for the binding of HOCl to a vitamin B12 derivative [[8,14](#)]. In a previous study, 200  $\mu\text{M}$  HOCl was found to react with and destroy the heme of hemoglobin in red blood cells [[12](#)]. This is consistent with recent work by Elahi et al., in which the harmful effect of COVID-19 on red blood cells is demonstrated [[46](#)].

As COVID-19 is known to be a disease of vascular endothelial inflammation, recruited neutrophils will likely be in relative proximity to passing erythrocytes [[47](#)]. HOCl and its adducts have been demonstrated to induce lysis of erythrocytes, exposing the erythrocytes' cytoplasmic hemoglobin to further oxidative damage [[[48](#)], [[49](#)], [[50](#)]]. Monochloramines derived from HOCl can also directly oxidize

hemoglobin within erythrocytes without hemolysis [51]. Therefore, high concentrations of HOCl can mediate heme destruction of several hemoproteins and [corrin](#) ring compounds including hemoglobin and other related hemoprotein model compounds such as [lactoperoxidase](#), myeloperoxidase, [catalase](#), [eosinophil peroxidase](#), and [cobalamin](#) derivatives, and more importantly can be prevented in the presence of 1:1 or 1:2 [melatonin](#) concentration [11,15,[52], [53], [54], [55]].

Given the toxicity of overwhelming production of HOCl and downstream products such as free iron, HOCl is physiologically regulated. MPO protection against self-inactivation is a necessary process that prevents HOCl mediated MPO heme destruction, therefore allowing the enzyme to function at full capacity in generating HOCl [53]. One potential pathway of preventing MPO auto-inactivation is the rapid consumption of HOCl through rapid reaction with [human serum albumin](#) (HSA) and other proteins found in blood and other bodily fluids [56]. However, during COVID-19-associated inflammation, HSA and other HOCl scavengers suffer from decreased effectiveness due to competitive binding of SARS-COV-2, and HOCl production already exacerbated by inflammatory mediators is further unchecked [57,58]. However, HOCl consumption by reaction with these substances does not account for the complete loss of HOCl-dependent signaling and viral killing, suggesting that alternative pathways exist for HOCl depletion. [In vitro studies](#) demonstrate that HOCl can react with a variety of hemoprotein model compounds at various oxidation states (e.g., ferric, ferrous, ferrous-dioxy, and ferryl) [11,12,59]. Reactions of HOCl with hemoprotein intermediates that catalyze electron transfer reactions are another potential pathway for HOCl consumption.

Excessive HOCl levels may adversely affect both the oxidation state of the hemoglobin (Hb) heme iron as well as heme stability. For example, the initial step of the reaction of oxyHb, the main O<sub>2</sub> transporter, with HOCl involves oxidation of heme iron from ferrous to ferric state and subsequent release of O<sub>2</sub> ([Fig. 1](#)) [12]. Hb then enters a state that cannot bind oxygen, forming an Fe(III)-OCl complex by binding OCl<sup>-</sup>. In the absence of a high O<sub>2</sub> concentration, the Fe(III)-OCl complex very quickly transitions to a hemoprotein ferryl π-cation complex, Compound I, via heterolytic cleavage of the O—Cl bond. Compound I is unstable and shortly decays into a ferryl complex, Compound II. In most cases of compounds with hemoprotein moieties, like in the case of hemoglobin, Compounds I and II populations depend on their formation and decay rates during steady state catalysis, and Compound II is known to be the most stable intermediate of hemoproteins. In vitro studies have shown that Compound II is also the predominant species formed prior to heme destruction, an irreversible step leading to depletion of the hemoprotein at hand [12].

Another family of essential enzymes that may be affected by increased HOCl production in COVID-19 are the three isoforms of [NO synthases](#): inducible (iNOS), neuronal (nNOS), and endothelial (eNOS). These enzymes have many roles such as vascular smooth muscle control, inhibition of [platelet aggregation](#), and vascular inflammation, whose dysfunction has been implicated in COVID-19 [47,60]. Reduction of NOS-Fe(III) by electrons provided by the [reductase](#) domain of the enzyme is the first step in the physiological reaction mechanism. Ferrous heme then combines with oxygen to form the corresponding Fe—O<sub>2</sub> complex. Further reduction of this intermediate by the cofactor H<sub>4</sub>B leads to the formation of a ferric-peroxo complex ([Fig. 1](#)). Subsequent breakage of the O—O bond leads to the formation of highly valent ferryl [porphyrin](#) π- cation radical complex, compound I,

which finally oxidizes the substrate, L-Arg, to form the intermediate N-hydroxyarginine (NOHA). Conversion of NOHA into [citrulline](#) and NO needs a second episode of oxygen activation, in which H<sub>4</sub>B is thought to act as the electron provider. Destabilization of the NOS-Fe(II)-O<sub>2</sub> and/or Compound I intermediates by HOCl during NO synthase steady-state catalysis may disturb NO production, leading to [vasoconstriction](#) in COVID-19 patients [[61](#), [62](#), [63](#)]. Pathological HOCl levels may also diminish NO generation by mediating NOS dimer subunit dissociation and [protein unfolding](#) through mechanisms that involve heme destruction and disturbance of the zinc-Cys cluster that controls H<sub>4</sub>B binding and the [enzyme stability](#) (Camp et al., unpublished results).

Catalase (CAT) and mammalian peroxidases (MPO, EPO, and LPO) are other examples of heme-containing enzymes that can be affected by overproduction of HOCl [[11,64](#)]. These enzymes have been previously proposed to be involved in the etiology of various lung diseases ranging from [cystic fibrosis](#) to acute hypoxemic respiratory failure, and now possibly COVID-19 [[65,66](#)]. CAT and MPO have been shown to interact through diffusion of the substrate H<sub>2</sub>O<sub>2</sub> and the product HOCl, regardless of the localization of CAT and MPO [[64](#)]. The interactions of OCl<sup>-</sup> with both mammalian peroxidases and CAT include its ability to form the Fe(III)-OCl complex and subsequently generate compounds I and II ([Fig. 1](#)). Therefore, HOCl overproduction could lead to the imbalance of MPO and CAT through protein modification and/or heme destruction [[64](#)]. Inactivation of both enzymes leads to H<sub>2</sub>O<sub>2</sub> accumulation, which, when combined with free iron, causes further oxidative damage to the host by generation of •OH. Therefore, the catalase/mammalian peroxidase heme destruction mediated by HOCl inhibits normal enzyme function, enhances iron (Fe) levels, and decreases levels of [antioxidants](#). The severity of this process depends mainly on the bioavailability of HOCl [[64](#)].

Not only is heme destruction irreversible and depletes essential enzymes in the fight against SARS-CoV-2, but it also releases toxic products such as free iron which subsequently generates [free radicals](#) along with various protein aggregates. [[1,12,15](#)] In particular, free iron acts as the [Fenton reagent](#), generating even more short-lived •OH from longer-lived species such as HOCl and H<sub>2</sub>O<sub>2</sub> [[1,32](#)]. Free iron can also damage blood vessels and increase [vascular permeability](#) through [vasodilation](#) [[65](#), [66](#), [67](#)]. This vasodilation not only leads to hypotension, but also metabolic acidosis. [Iron overload](#) may explain increases in the risk of ischemic cardiovascular events, accelerated [thrombus formation](#), impaired vasoreactivity, and enhance the production of ROS in patients infected with SARS-CoV-2, found by some authors [[1](#)]. Therefore, in COVID-19, despite O<sub>2</sub>/NO supplementation, the desired physiological response may be absent as essential heme-containing enzymes and proteins, such as Hb and NOS, have been disturbed biochemically through increases in HOCl and other ROS.

### **3. HOCl mediates corrin ring (vitamin B12) destruction**

The association between COVID-19 and [Vitamin B12 deficiency](#) leading to worse outcomes of respiratory viral infections has been previously established [[68](#)]. Furthermore, COVID-19 and [vitamin B12](#) deficiency have shared symptoms, such as increased [oxidative stress](#), increased [dehydrogenase](#), hyperhomocysteinemia,



hypercoagulation, and [vasoconstriction](#), particularly of the renal and pulmonary systems [69]. Consistent with this, recent investigation has shown that [methylcobalamin](#) supplements not only have the potential to reduce COVID-19-related organ damage, but also reduce other symptoms [68,69]. A clinical study has also shown reduced COVID-19 symptom severity in patients who received vitamin B12 (500 µg), [vitamin D](#) (1000 IU), and magnesium supplements, displaying significant reduction in the need for O<sub>2</sub> and intensive care support [69,70].

Using a variety of biochemical, physiological, and kinetic techniques, it has been shown that [melatonin](#) ceases HOCl-mediated vitamin B12 derivative [corrin](#) destruction [8,14]. Vitamin B12 deficiency has been associated with several conditions related to memory loss, immune system disorders, aging, heart disease, male infertility, diabetes, sleep disorders, depression, mental disorders, and inflammation [71]. Patients with one or more of these conditions might be more susceptible to severe SARS-CoV-2 infection [72]. In addition to ROS generated by the [cytokine storm](#), SARS-CoV-2 infection may also interfere with vitamin B12 metabolism, therefore contributing to the pathogenesis of respiratory, gastrointestinal, and central nervous systems infections [69].

The two derivatives most commonly used for treatment of vitamin B12 deficiency are [hydroxocobalamin](#) and cyanocobalamin, with cyanocobalamin being preferred in the United States. However, outside of the U.S., such as in the United Kingdom, hydroxocobalamin is preferred due to its ability to firmly bind to [plasma proteins](#) allowing it to remain in the body longer [73]. Although the amount of scientific evidence is limited, the safety of cyanocobalamin supplementation for COVID-19 patients, versus other vitamin B12 derivatives, may be a concern. The [biosynthesis](#) of the B12 coenzyme from cyanocobalamin results in the release of CN<sup>-</sup>, which could lead to acute cyanide poisoning, causing unwanted inflammation. Furthermore, when HOCl levels are high, HOCl-mediated reactions may cause the destruction of the corrin ring of vitamin B12, releasing [cyanogen chloride](#) (CNCl), free active cobalt (Co), and other corrin [degradation products](#). Indeed, our previously published results suggest that the degradation of cyanocobalamin mediated by HOCl is largely modulated by the concentration of HOCl in the reaction milieu. Thus, any dysregulation of neutrophil/macrophage derived HOCl contributes to inflammation and, similarly, reduction in HOCl consumption could manifest as increased cyanocobalamin destruction and CNCl generation [14]. Therefore, it is of enormous therapeutic and pharmacologic importance to prevent HOCl-mediated damage, especially in chronic inflammation where a higher rate of infiltration of monocytes/macrophages over a longer period leads to pathologic alterations.

## **4. Mechanism of HOCl mediated tetrapyrrole ring destruction (Heme in hemoprotein and corrin ring in vitamin B12)**

In [hemoprotein](#), Fe is attached to the nitrogen of four pyrrole rings and a fifth proximal attachment to either the nitrogen of histidine such as in hemoglobin, [catalase](#), and mammalian peroxidases or to the sulfur of cysteine, as seen in [NOS](#) and [cytochrome P450](#). The sixth axial site of Fe can accommodate a variety of diatomic ligands such as O<sub>2</sub>, NO, CO, CN<sup>-</sup> and <sup>-</sup>OCl. The excessive production of ROS in the [cytokine storm](#) can result in modification of the heme [prosthetic group](#)

inhibiting the protein function, whereas in [vitamin B12](#) derivatives, the structure is based on a [corrin](#) ring containing four pyrrole rings attached to a center [Co atom](#), distinguished by two directly attached pyrrole rings. The six coordination sites of Co in these compounds are the four pyrrole [nitrogen atoms](#) of the corrin ring, the nitrogen of the 5,6-dimethylbenzimidazole group at the lower (or  $\alpha$ -) axial ligand, and naturally occurs with either a cyano-, hydroxo-, aquo-, methyl-, or adenosyl-group at the upper/ $\beta$ -axial ligand site [74]. Here, the overproduction of ROS and HOCl from the cytokine storm nonenzymatically mediates corrin ring destruction and the generation of free Co [14].

Pyrrole ring destruction by HOCl can occur as a result of direct attack on any of the carbon-methylene bridges between the rings, forming chlorinated adducts [12,14]. These chlorinated intermediates are unstable, quickly releasing  $\text{Cl}^-$  and decaying to an [epoxide](#) or [aminal](#). The epoxide implements the buildup of a hydroxylated compound, with the  $\cdot\text{OH}$  group being connected to the carbon-methylene bridge of the [porphyrin](#) ring where the initial attack by HOCl takes place [12,14]. Attack by a second [hydroxyl](#) functional group creates vicinal diol, which can be split by either [hemolytic](#) cleavage or a transition metal mediated process (Fe, Co) by the formation of [dioxetane](#) intermediate through a heterolytic  $2e^-$  process; both processes result in a pair of [carbonyl compounds](#). The aldehydes that are produced may be further oxidized by HOCl to create carboxylic acid through a mechanism previously described [75]. Cleavage of the  $\text{C}=\text{C}$  bond can take place at both the carbon-methylene bridge and the terminal  $\text{C}=\text{C}$  bond, leading to the generation of [formaldehyde](#) [12]. This single carbon [aldehyde](#) can be oxidized to [formic acid](#) by the [electrophilic addition](#) of HOCl. As recently reported, HOCl can alter the [tetrapyrrole](#) ring through a mechanism that involves disrupting the axial coordination of the Fe (hemoproteins) or Co (vitamin B12 derivatives) atom, causing ring destruction [8,14]. These alterations of the tetrapyrrole ring geometry might therefore make the ring a more eligible target to HOCl-attack and ring breakage, which is associated with significant Fe/Co release.

## 5. Other effects of ROS

Generation of ROS and [reactive nitrogen species](#) (RNS) by the [cytokine storm](#) in COVID-19 can also result from increased [matrix metalloproteinase](#) (MMP) expression [76]. MMPs are zinc- and calcium-dependent enzymes associated with extracellular matrix remodeling that can also enhance activation of [chemokines](#) and cytokines from leukocytes [77,78]. In turn, generation of ROS from activated macrophages further activates and increases the expression of MMPs. Conversely, [melatonin](#) has been demonstrated to downregulate MMP expression, MMP protein level, and upregulate tissue inhibitors of MMPs [79,80]. These modulatory actions can prevent extracellular matrix remodeling associated with the inflammatory [signaling molecules](#), especially interleukin- $1\beta$ . ROS can also disrupt the function of [metalloproteins](#) by causing disturbance in their structural components and releasing free metals. Of note, HOCl can disturb the zinc-tetrathiolate cluster of [iNOS](#) leading to the irreversible release of zinc [81]. Other examples of ROS mediated metalloprotein disruption include the release of free iron from ferritin by  $\text{O}_2^{\cdot-}$ , as well as rapid inactivation of enzymes of the dehydratase-lyase family by  $\text{O}_2^{\cdot-}$  followed by the [oxidation](#) of the iron-sulfur cluster releasing Fe(II) [82].

## 6. Melatonin is a potent inhibitor of MPO

Studies have shown the severity of disease is correlated with the level of inflammatory immune response caused from the pro-inflammatory cytokines released in the [cytokine storm](#), with an exceptionally heightened response in those with more severe cases. Neutrophil activity, [MPO](#) activity, and ROS generation play an important role in the inflammatory immune response that contributes to the overwhelming production of HOCl; thus, MPO inhibition and elimination of unwanted ROS are undoubtedly important treatment targets in patients with COVID-19. It has been shown that triggering NET formation is dependent upon ROS formation and processing by MPO; therefore, inhibiting MPO could block the formation of NETs [4,83]. [Melatonin](#) is not only a reversible inhibitor and important regulator of MPO activity, but also plays an important role in detoxifying ROS. These phenomena position the indole as a powerful supplement to fight against early stages and severe SARS-CoV-2 infection. Melatonin-engaging therapies could reduce the stage of virus-associated pathology by controlling the host immune response to viral infection, mainly produced by [alveolar macrophages](#) and neutrophil MPO activity and unwanted ROS overproduction. Melatonin inhibits MPO chlorinating activity in a dual mechanism, which includes allosteric binding to the entrance of the MPO heme pocket and accelerating the formation and the decay of MPO compound II, which inhibits the chlorinating activity and slows down the [peroxidation](#) activity of the enzyme [23,24]. This allosteric binding to the MPO heme pocket entrance is enhanced by  $\text{Cl}^-$  binding to the halide-binding site, where HOCl is generated, and blocks  $\text{H}_2\text{O}_2$  from the catalytic site. Thus, melatonin competes with  $\text{H}_2\text{O}_2$  and switches the reaction from free to melatonin-bound enzyme (active to inactive form), and melatonin also competes with the co-substrate,  $\text{Cl}^-$ , and switches the reaction from a  $2e^-$  to a  $1e^-$  [oxidation](#) mechanism [23]. Studies have shown that rapid mixing of MPO preincubated with melatonin and  $\text{Cl}^-$  against the same volume of  $\text{H}_2\text{O}_2$  solution caused immediate buildup of a transient intermediate, compound II, which then decays to MPO – Fe(III) through oxidation of another melatonin molecule, thereby closing the peroxidation cycle [23]. The ability of melatonin to compete with  $\text{H}_2\text{O}_2$  and  $\text{Cl}^-$  on the active sites of MPO – Fe(III) and MPO compound I, respectively, is a key feature that drives the enzyme to alter its function to peroxidase activity. This mechanism of competitive inhibition has been shown to occur with tryptophan and several phenolic and aromatic amines [84].

More recently, kinetic evidence for an allosteric coupling among melatonin, the MPO entrance binding (regulatory) site, and the catalytic center that induces detectable inhibition of MPO has been provided [23,53]. Consequently, oxidation of melatonin is required before it diffuses from the binding site of the enzyme and restores the [catalytic activity](#) of MPO [23]. Thus, inactivation of MPO and its catalytic duration can be controlled effectively by melatonin supplementation. Indeed, when the melatonin concentration is less than twice the  $\text{H}_2\text{O}_2$  concentration,  $\text{H}_2\text{O}_2$  consumption proceeds in a slower and linear manner, and MPO decays rapidly to MPO – Fe(III) after melatonin oxidation. This behavior clearly demonstrates that MPO is capable of restoring its catalytic activity and rejoining the peroxidase cycle after melatonin exhaustion [23,53]. On the other hand, when the melatonin concentration is greater than twice the  $\text{H}_2\text{O}_2$  concentration, the initial slow phase of  $\text{H}_2\text{O}_2$  consumption remains at the same rate through the progression of the reaction and ceases when  $\text{H}_2\text{O}_2$  is completely consumed [23,53]. The oxidation of melatonin is supported by the control studies that show no  $\text{H}_2\text{O}_2$  consumption in the absence of MPO. When functioning as a  $1e^-$

substrate, melatonin competes with  $\text{Cl}^-$  and promotes alterations in substrate switching influencing the distribution of peroxidase intermediates capable of executing 1e- versus 2e- oxidation reactions. Understanding the circumstances that lead to MPO inhibition can provide important information and a general framework of understanding the role of MPO in certain diseases [85].

Computational modeling of melatonin has shown the potential docking of melatonin to MPO-Fe(III), compound I, and compound II forms, although the crystal structure of MPO-melatonin complex remains unresolved. As melatonin is a relatively bulky molecule, the indole ring must dock parallel to the plane of the heme, with the side chain oriented towards the outside of the distal cavity [86]. Under these circumstances, the indole ring is close to the MPO D pyrrole with a distance average of the closest indole atom to the center of the D ring is ranging between 3.3 and 3.5 Å. Comparing MPO – Fe(III), compound I, and compound II modeling pointed that melatonin was deviated ~1 Å from the ferryl oxygen in compounds I and II. Therefore, the ability of melatonin to bind to MPO intermediates makes it an effective inhibitor of MPO activity.

## 7. Melatonin is a potent scavenger of ROS

ROS induced by viral infection are necessary for eliminating viruses phagocytosed by immune cells, and take part in [signal transduction](#) between various immune cells [[87], [88], [89]]. However, the inhibitory effects controlled by the immune system may be affected by aging and diseases (e.g. diabetes, cancer, and heart problems), causing some groups of individuals to be more vulnerable to SARS-CoV-2 infection [72]. Similarly, the inhibitory effects are also genetically controlled and thus may fluctuate with race and ethnicities [72,90]. Thus, the pathogenicity of new variants of SARS-CoV-2, varying immune response, imbalanced and uncontrolled ROS production associated with the [cytokine storm](#), exuberant endothelial inflammatory reactions, vascular thrombosis, and bioavailability of enzymatic and non-enzymatic [antioxidant](#) machinery, in combination, may all become the determining factors that enable the virus to initiate infection [1].

Through melatonin supplementation, damaging conditions such as protein oxidation, [lipid peroxidation](#), and DNA damage are diminished by reduction of ROS and free [metal ion](#) production [53,[91], [92], [93], [94], [95]]. Mass spectrometric and high-performance liquid chromatography studies have identified melatonin metabolites together with HOCl, intact [tetrapyrrole](#) rings, and [hemoprotein](#), indicating that supplemented melatonin is likely consumed in exertion of its protective effects [54]. Furthermore, these melatonin metabolites are not known to have biological consequences, nor do they seem to effect [MPO](#) activity [23]. Other studies have highlighted melatonin's antioxidant ability in comparison with other known HOCl scavengers such as [lycopene](#), methionine, taurine, [cystine](#), cysteine, and [uric acid](#) [24,75,96].

[Melatonin](#) can also inhibit the activity of MMPs through binding of melatonin to the active site, as shown in [molecular docking simulations](#) [97]. As previously explained, melatonin can further suppress deleterious effects of excessive MMPs by directly scavenging the ROS that increase [MMP](#) expression as well as protect against inflammatory lung pathologies, such as ARDS and COVID-19, through suppression of inflammation, [oxidative stress](#), DNA damage, and cell apoptosis. [98,99].

## 8. Vitamin B12, melatonin, myeloperoxidase, and sleep

[Vitamin B12](#) is essential for neural function, and a significant portion of the elderly population is deficient in this water-soluble vitamin [100]. Vitamin B12 accompanies [melatonin](#) as an [antioxidant](#) and a regulatory agent in circadian rhythm. By acting on the pineal gland, vitamin B12 takes part in rearrangement of the sleep-wake cycle through activation of the direct earlier release of melatonin at dark, as well as through causing melatonin to drift off faster by alerting the body to morning light [101]. Due to these properties, vitamin B12 derivatives (cyano-, hydroxo-, aquo-, methyl-, or adenosyl-) have been successfully used as therapeutic drugs for the treatment of serious sleep-wake disorders associated with COVID-19 patients [101].

Melatonin and peroxidase activities are apparently coupled through complex and interdependent pathways. It appears that the affinity of MPO for melatonin is very high under physiological conditions [23]. MPO may be acting as a chelator for melatonin and limiting its bioavailability and function. This may partly explain the harmful effects of sleep loss on various body functions. In addition, melatonin may have an important role in the inhibition of MPO activity in various tissues during inflammation. Hence, it is possible to speculate that increased inflammation and decreased immunity associated with lack of sleep are the result of a lack of inhibitory protective effects of melatonin on MPO during prolonged waking hours and chronic sleep loss. On the other hand, the pathogenesis of various disease processes can be very complicated and involves multiple inflammatory mediators and oxidizing agents. The biological consequences of melatonin–peroxidase interactions may have broad implications in the regulation of sleep, inflammation, infectious, cardiovascular events in vivo, and COVID-19.

## 9. Preserving lung mechanics through melatonin supplementation

Melatonin potentially has several other benefits against COVID-19 in addition to what has already been listed. As discussed by Zhang et al., in the case of COVID-19, there is high risk of developing ALI/ARDS and other complications such as respiratory failure, heart failure, sepsis, and even sudden cardiac arrest, a progression similar to that of SARS- and MERS- induced pneumonia [21]. Melatonin can act against several elements of ALI/ARDS processes. It mediates activity of sirtuins such as sirtuin-1, which can downregulate transition of macrophages towards pro-inflammation through inhibition of [high mobility group box chromosomal protein 1](#) and participate in other antiviral processes [[102], [103], [104]]. Melatonin can also suppress the activation of NF- $\kappa$ B, a pro-inflammatory mediator in ALI/ARDS, both in [T cells](#) and lung tissue [105,106]. Nuclear factor erythroid 2-related factor 2 (NRF2), a pulmonoprotective factor, has been shown to be enhanced by [melatonin](#) and has been postulated to be involved in CoV-associated ALI/ARDS. Melatonin also upregulates anti-oxidative enzymes and downregulates pro-oxidative enzymes and has been shown to directly enhance the immune response through improvement of proliferation and maturation of natural killing cells, T and B lymphocytes, [granulocytes](#), and [monocytes](#) [107]. Melatonin has also been shown to decrease the level of circulating cytokines [21]. These anti-inflammatory and pulmonoprotective effects of melatonin further suggest its

possible suitability for treatment of COVID-19.

## 10. Conclusion

With the discovery of multiple COVID-19 variants spreading around the world, the question remaining is can the current vaccines keep up, or will they become less effective against SARS-CoV-2 mutations? There is a possibility that the cost and time needed to develop a vaccine targeted at new strains may not be met before another spike occurs. In this case, recognition of treatment options is of utmost importance. [Melatonin](#) is particularly notable, as its supplementation is safe for use with other treatments, as it has been previously given to COVID-19 patients in combination with the US Food and Drug Administration emergency authorized cocktail, REGEN-COV2 [[20,21](#)].

Many of the known clinical signs of COVID-19 could potentially be associated with defects in the function of several key [tetrapyrrole](#) and heme proteins, notably hemoglobin, [cyanocobalamin](#), and [NO synthases](#). Here, we have outlined the destructive effects of ROS, especially HOCl, on these biomolecules. As one of the major sources of HOCl is through overactivity of [MPO](#) provoked by the [cytokine storm](#) of COVID-19, inhibition of MPO and/or elimination of HOCl may play a beneficial role in diverse [biological processes](#) by reducing the metal release mediated by HOCl and associated increases in other ROS. Related studies from our lab have shown that MPO and its effects can be inhibited at three points: 1) through heme reduction that causes collapse or narrowing in heme pocket geometry that prevents the access of the substrate to the catalytic site of the enzyme (e.g. ascorbate) [[108](#)]; 2) switching the MPO catalytic cycle from [peroxidation](#) to catalase-like activity (e.g. melatonin, tryptophan, tryptophan analogs) [[23,84,109](#)]; or 3) direct scavenging of HOCl (e.g. lycopene) [[75](#)].

The ability of melatonin to act as an [antioxidant](#), anti-inflammatory, and immunoregulatory agent allows the unique opportunity for its use in a variety of therapeutic approaches, and it has shown promising benefits in cancer, diabetes, infertility, and inflammatory diseases. The outstanding benefits of melatonin provide an accessible, relatively inexpensive, and safe treatment that may ease COVID-19 symptoms commonly seen with the cytokine storm and overactive immune response, leading to improved patient outcomes and reduced suffering. In addition to use alone, promising evidence is emerging to establish melatonin as a therapeutic treatment that could potentially be used in combination with current [antiviral agents](#). Further clinical and experimental studies are required to confirm the applied benefits of melatonin supplementation as a therapeutic agent for treatment of patients infected with SARS-COV2.

## Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

## References

[1]

P.T. Goud, D. Bai, H.M. Abu-Soud

Int. J. Biol. Sci., 17 (2021), pp. 62-72

[2]

A.M. Dondorp, M. Hayat, D. Aryal, A. Beane, M.J. Schultz

Am. J. Trop. Med. Hyg., 102 (2020), pp. 1191-1197

[3]

H. Parker, C.C. Winterbourn

Front. Immunol., 3 (2012), p. 424

[4]

H. Bjornsdottir, A. Welin, E. Michaelsson, V. Osla, S. Berg, K. Christenson, M. Sundqvist, C. Dahlgren, A. Karlsson, J. Bylund

Free Radic. Biol. Med., 89 (2015), pp. 1024-1035

[5]

E.A. Podrez, H.M. Abu-Soud, S.L. Hazen

Free Radic. Biol. Med., 28 (2000), pp. 1717-1725

[6]

M.J. Davies, C.L. Hawkins, D.I. Pattison, M.D. Rees

Antioxid. Redox Signal., 10 (2008), pp. 1199-1234

[7]

J.M. Pullar, M.C. Vissers, C.C. Winterbourn

IUBMB Life, 50 (2000), pp. 259-266

[8]

D. Maitra, I. Ali, R.M. Abdulridha, F. Shaeib, S.N. Khan, G.M. Saed, S. Pennathur, H.M. Abu-Soud

PLoS One, 9 (2014)

[9]

A.J. Kettle, A.M. Albrett, A.L. Chapman, N. Dickerhof, L.V. Forbes, I. Khalilova, R. Turner

Biochim. Biophys. Acta, 1840 (2014), pp. 781-793

[10]

A. Ulfig, L.I. Leichert

Cell. Mol. Life Sci., 78 (2021), pp. 385-414

[11]

M.G. Bonini, A.G. Siraki, B.S. Atanassov, R.P. Mason

Free Radic. Biol. Med., 42 (2007), pp. 530-540

[12]

D. Maitra, J. Byun, P.R. Andreana, I. Abdulhamid, M.P. Diamond, G.M. Saed, S. Pennathur, H.M. Abu-Soud

Free Radic. Biol. Med., 51 (2011), pp. 374-386

[13]

D. Maitra, J. Byun, P.R. Andreana, I. Abdulhamid, G.M. Saed, M.P. Diamond, S. Pennathur, H.M. Abu-Soud

Free Radic. Biol. Med., 51 (2011), pp. 364-373

[14]

H.M. Abu-Soud, D. Maitra, J. Byun, C.E.A. Souza, J. Banerjee, G.M. Saed, M.P. Diamond, P.R. Andreana, S. Pennathur

Free Radic. Biol. Med., 52 (2012), pp. 616-625

[15]

C.E. Souza, D. Maitra, G.M. Saed, M.P. Diamond, A.A. Moura, S. Pennathur, H.M. Abu-Soud

PLoS One, 6 (2011)

[16]

M.H. Zou, C. Shi, R.A. Cohen

J. Clin. Invest., 109 (2002), pp. 817-826

[17]

J.S. Beckman, J. Chen, H. Ischiropoulos, J.P. Crow

Methods Enzymol., 233 (1994), pp. 229-240

[18]

B. Halliwell

Br. J. Exp. Pathol., 70 (1989), pp. 737-757



[19]

Y. Zhou, Y. Hou, J. Shen, R. Mehra, A. Kallianpur, D.A. Culver, M.U. Gack, S. Farha, J. Zein, S. Comhair, C. Fiocchi, T. Stappenbeck, T. Chan, C. Eng, J.U. Jung, L. Jehi, S. Erzurum, F. Cheng

PLoS Biol., 18 (2020)

[20]

R.J. Reiter, P. Abreu-Gonzalez, P.E. Marik, A. Dominguez-Rodriguez

Front. Med. (Lausanne), 7 (2020), p. 226

[21]

R. Zhang, X. Wang, L. Ni, X. Di, B. Ma, S. Niu, C. Liu, R.J. Reiter

Life Sci., 250 (2020), p. 117583

[22]

V. Ramlall, J. Zucker, N. Tatonetti

medRxiv (2020)

[23]

S. Galijasevic, I. Abdulhamid, H.M. Abu-Soud

Biochemistry, 47 (2008), pp. 2668-2677

[24]

T. Lu, S. Galijasevic, I. Abdulhamid, H.M. Abu-Soud

Br. J. Pharmacol., 154 (2008), pp. 1308-1317

[25]

D.P. Cardinali, G.M. Brown, S.R. Pandi-Perumal

Diseases, 8 (2020)

[26]

D. Bonnefont-Rousselot, F. Collin

Toxicology, 278 (2010), pp. 55-67

[27]

A. Galano, D.X. Tan, R.J. Reiter

J. Pineal Res., 51 (2011), pp. 1-16

[28]

R.J. Reiter, J.C. Mayo, D.X. Tan, R.M. Sainz, M. Alatorre-Jimenez, L. Qin  
J. Pineal Res., 61 (2016), pp. 253-278

[29]

A.G. Wade, I. Ford, G. Crawford, A. McConnachie, T. Nir, M. Laudon, N.  
Zisapel

BMC Med., 8 (2010), p. 51

[30]

L.P. Andersen, I. Gogenur, J. Rosenberg, R.J. Reiter

Clin. Drug Investig., 36 (2016), pp. 169-175

[31]

R. Sethuram, D. Bai, H.M. Abu-Soud

**Potential role of zinc in the COVID-19 disease process and its  
probable impact on reproduction**

[32]

F. Shaeib, J. Banerjee, D. Maitra, M.P. Diamond, H.M. Abu-Soud

Free Radic. Biol. Med., 58 (2013), pp. 154-159

[33]

F. Collin

Int. J. Mol. Sci., 20 (2019)

[34]

H.M. Abu-Soud, S.L. Hazen

J. Biol. Chem., 275 (2000), pp. 37524-37532

[35]

U. Burner, P.G. Furtmuller, A.J. Kettle, W.H. Koppenol, C. Obinger

J. Biol. Chem., 275 (2000), pp. 20597-20601

[36]

F.A. Summers, P.E. Morgan, M.J. Davies, C.L. Hawkins

Chem. Res. Toxicol., 21 (2008), pp. 1832-1840

[37]

R. Stocker, J.F. Keaney Jr.

Physiol. Rev., 84 (2004), pp. 1381-1478

[38]

D. Maitra, F. Shaeib, I. Abdulhamid, R.M. Abdulridha, G.M. Saed, M.P. Diamond, S. Pennathur, H.M. Abu-Soud

Free Radic. Biol. Med., 63 (2013), pp. 90-98

[39]

M.Z. Tay, C.M. Poh, L. Renia, P.A. MacAry, L.F.P. Ng

Nat. Rev. Immunol., 20 (2020), pp. 363-374

[40]

O.A. Khomich, S.N. Kochetkov, B. Bartosch, A.V. Ivanov

Viruses, 10 (2018)

[41]

Q. Liu, Y.H. Zhou, Z.Q. Yang

Cell. Mol. Immunol., 13 (2016), pp. 3-10

[42]

R. Floris, R. Wever

Eur. J. Biochem., 207 (1992), pp. 697-702

[43]

D.I. Pattison, M.J. Davies

Chem. Res. Toxicol., 14 (2001), pp. 1453-1464

[44]

P.G. Furtmuller, U. Burner, W. Jantschko, G. Regelsberger, C. Obinger

Redox Rep., 5 (2000), pp. 173-178

[45]

T. Nybo, M.J. Davies, A. Rogowska-Wrzesinska

Redox Biol., 26 (2019), p. 101236

[46]

S. Shahbaz, L. Xu, M. Osman, W. Sligl, J. Shields, M. Joyce, D.L. Tyrrell, O. Oyegbami, S. Elahi

Stem Cell Reports, 16 (2021), pp. 1165-1181

[47]

J. Zhang, K.M. Tecson, P.A. McCullough

Rev. Cardiovasc. Med., 21 (2020), pp. 315-319

[48]

C.L. Hawkins, B.E. Brown, M.J. Davies

Arch. Biochem. Biophys., 395 (2001), pp. 137-145

[49]

M.C. Vissers, A. Stern, F. Kuypers, J. van den Berg, C.C. Winterbourn

Free Radic. Biol. Med., 16 (1994), pp. 703-712

[50]

I.B. Zavodnik, E.A. Lapshina, L.B. Zavodnik, M. Soszynski, G. Bartosz, M. Bryszewska

Bioelectrochemistry, 58 (2002), pp. 127-135

[51]

M.B. Grisham, M.M. Jefferson, E.L. Thomas

J. Biol. Chem., 259 (1984), pp. 6757-6765

[52]

D. Maitra, I. Abdulhamid, M.P. Diamond, G.M. Saed, H.M. Abu-Soud

J. Pineal Res., 53 (2012), pp. 198-205

[53]

F. Shaeib, S.N. Khan, I. Ali, T. Najafi, D. Maitra, I. Abdulhamid, G.M. Saed, S. Pennathur, H.M. Abu-Soud

PLoS One, 10 (2015)

[54]

R. Jeelani, D. Maitra, C. Chatzicharalampous, S. Najeemuddin, R.T. Morris, H.M. Abu-Soud

J. Pineal Res., 64 (2018)

[55]

H.M. Abu-Soud, D. Maitra, F. Shaeib, S.N. Khan, J. Byun, I. Abdulhamid, Z. Yang, G.M. Saed, M.P. Diamond, P.R. Andreana, S. Pennathur

J. Inorg. Biochem., 140 (2014), pp. 245-254

[56]

A. Ulfig, V. Bader, M. Varatnitskaya, N. Lupilov, K.F. Winklhofer, L.I. Leichert

Redox Biol., 43 (2021), p. 101981

[57]

A.S. Johnson, R. Fatemi, W. Winlow

Front. Cardiovasc. Med., 7 (2020), p. 153

[58]

V. Arroyo, R. Garcia-Martinez, X. Salvatella

J. Hepatol., 61 (2014), pp. 396-407

[59]

L. Gebicka, E. Banasiak

Toxicol. in Vitro, 26 (2012), pp. 924-929

[60]

U. Forstermann, W.C. Sessa

Eur. Heart J., 33 (2012), pp. 829-837

[61]

J.R. Klinger, S.H. Abman, M.T. Gladwin

Am. J. Respir. Crit. Care Med., 188 (2013), pp. 639-646

[62]

W. Fang, J. Jiang, L. Su, T. Shu, H. Liu, S. Lai, R.A. Ghiladi, J. Wang

Free Radic. Biol. Med., 163 (2021), pp. 153-162

[63]

J.C. Pieretti, O. Rubilar, R.B. Weller, G.R. Tortella, A.B. Seabra

Virus Res., 291 (2021), p. 198202

[64]

I. Ali, S.N. Khan, C. Chatzicharalampous, D. Bai, H.M. Abu-Soud

J. Inorg. Biochem., 197 (2019), p. 110706

[65]

W.Y. Ong, B. Halliwell

Ann. N. Y. Acad. Sci., 1012 (2004), pp. 51-64

[66]

D. Trinder, C. Fox, G. Vautier, J.K. Olynyk

Gut, 51 (2002), pp. 290-295

[67]

Y. Kohgo, K. Ikuta, T. Ohtake, Y. Torimoto, J. Kato

Int. J. Hematol., 88 (2008), pp. 7-15

[68]

T.H. Jovic, S.R. Ali, N. Ibrahim, Z.M. Jessop, S.P. Tarassoli, T.D. Dobbs, P. Holford, C.A. Thornton, I.S. Whitaker

Nutrients, 12 (2020)

[69]

H. Shakoor, J. Feehan, K. Mikkelsen, A.S. Al Dhaheri, H.I. Ali, C. Platat, L.C. Ismail, L. Stojanovska, V. Apostolopoulos

Maturitas, 144 (2021), pp. 108-111

[70]

C.W. Tan, L.P. Ho, S. Kalimuddin, B.P.Z. Cherng, Y.E. Teh, S.Y. Thien, H.M. Wong, P.J.W. Tern, M. Chandran, J.W.M. Chay, C. Nagarajan, R. Sultana, J.G.H. Low, H.J. Ng

Nutrition, 79–80 (2020)

[71]

F. O'Leary, S. Samman

Nutrients, 2 (2010), pp. 299-316

[72]

E.J. Williamson, A.J. Walker, K. Bhaskaran, S. Bacon, C. Bates, C.E. Morton, H.J. Curtis, A. Mehrkar, D. Evans, P. Inglesby, J. Cockburn, H.I. McDonald, B. MacKenna, L. Tomlinson, I.J. Douglas, C.T. Rentsch, R. Mathur, A.Y.S. Wong, R. Grieve, D. Harrison, H. Forbes, A. Schultze, R. Croker, J. Parry, F. Hester,

S. Harper, R. Perera, S.J.W. Evans, L. Smeeth, B. Goldacre

Nature, 584 (2020), pp. 430-436

[73]

J. Minigh

**XPharm: The Comprehensive Pharmacology Reference**

Elsevier (2007), pp. 1-6

[74]

R. Banerjee, S.W. Ragsdale

Annu. Rev. Biochem., 72 (2003), pp. 209-247

[75]

S. Pennathur, D. Maitra, J. Byun, I. Sliskovic, I. Abdulhamid, G.M. Saed, M.P. Diamond, H.M. Abu-Soud

Free Radic. Biol. Med., 49 (2010), pp. 205-213

[76]

T. Ueland, J.C. Holter, A.R. Holten, K.E. Muller, A. Lind, G.K. Bekken, S. Dudman, P. Aukrust, A.M. Dyrhol-Riise, L. Heggelund

J. Inf. Secur., 81 (2020), pp. e41-e43

[77]

D.A. Siwik, W.S. Colucci

Heart Fail. Rev., 9 (2004), pp. 43-51

[78]

R.T. de Pinho, W.S. da Silva, L.M. de Castro Cortes, P. da Silva Vasconcelos Sousa, R.O. de Araujo Soares, C.R. Alves

Exp. Parasitol., 147 (2014), pp. 72-80

[79]

W. Qin, J. Li, R. Zhu, S. Gao, J. Fan, M. Xia, R.C. Zhao, J. Zhang

Aging (Albany NY), 11 (2019), pp. 11391-11415

[80]

Y. Zhang, F. He, Z. Chen, Q. Su, M. Yan, Q. Zhang, J. Tan, L. Qian, Y. Han

Aging (Albany NY), 11 (2019), pp. 10499-10512

[81]

H. Li, C.S. Raman, C.B. Glaser, E. Blasko, T.A. Young, J.F. Parkinson, M. Whitlow, T.L. Poulos

J. Biol. Chem., 274 (1999), pp. 21276-21284

[82]

K. Jomova, M. Valko

Toxicology, 283 (2011), pp. 65-87

[83]

G. Schonrich, M.J. Raftery, Y. Samstag

Adv. Biol. Regul., 77 (2020)

[84]

S. Galijasevic, I. Abdulhamid, H.M. Abu-Soud

Free Radic. Biol. Med., 44 (2008), pp. 1570-1577

[85]

V.F. Ximenes, S.O. Silva, M.R. Rodrigues, L.H. Catalani, G.J. Maghzal, A.J. Kettle, A. Campa

J. Biol. Chem., 280 (2005), pp. 38160-38169

[86]

H.R. Hallingback, R.R. Gabdoulline, R.C. Wade

Biochemistry, 45 (2006), pp. 2940-2950

[87]

M. Seyoum, B. Enawgaw, M. Melku

Thromb. J., 16 (2018), p. 16

[88]

C.G. Molteni, N. Principi, S. Esposito

Free Radic. Res., 48 (2014), pp. 1163-1169

[89]

D.G. Franchina, C. Dostert, D. Brenner

Trends Immunol., 39 (2018), pp. 489-502



[90]

C.O. Jacob

Clin. Immunol., 220 (2020), p. 108591

[91]

R.J. Reiter, A. Korkmaz, S. Ma, S. Rosales-Corral, D.X. Tan

Mutat. Res. Rev. Mutat. Res., 751 (2012), pp. 7-14

[92]

F.J. Gomez, J. Raba, S. Cerutti, M.F. Silva

J. Pineal Res., 52 (2012), pp. 349-355

[93]

A. Carrillo-Vico, J.M. Guerrero, P.J. Lardone, R.J. Reiter

Endocrine, 27 (2005), pp. 189-200

[94]

I. Gulcin, M.E. Buyukokuroglu, O.I. Kufrevioglu

J. Pineal Res., 34 (2003), pp. 278-281

[95]

J. Banerjee, D. Maitra, M.P. Diamond, H.M. Abu-Soud

J. Pineal Res., 53 (2012), pp. 122-128

[96]

T. Ogino, T.A. Than, M. Hosako, M. Ozaki, M. Omori, S. Okada

Adv. Exp. Med. Biol., 643 (2009), pp. 451-461

[97]

D.S. Rudra, U. Pal, N.C. Maiti, R.J. Reiter, S. Swarnakar

J. Pineal Res., 54 (2013), pp. 398-405

[98]

B. Solun, Y. Shoenfeld

Med. Drug Discov., 7 (2020)

[99]

C.K. Sun, F.Y. Lee, Y.H. Kao, H.J. Chiang, P.H. Sung, T.H. Tsai, Y.C. Lin, S. Leu, Y.C. Wu, H.I. Lu, Y.L. Chen, S.Y. Chung, H.L. Su, H.K. Yip

J. Pineal Res., 58 (2015), pp. 137-150

[\[100\]](#)

P.J. Stover

Curr. Opin. Clin. Nutr. Metab. Care, 13 (2010), pp. 24-27

[\[101\]](#)

M. Ikeda, M. Asai, T. Moriya, M. Sagara, S. Inoue, S. Shibata

Brain Res., 795 (1998), pp. 98-104

[\[102\]](#)

R. Hardeland

J. Pineal Res., 65 (2018), Article e12525

[\[103\]](#)

G. Anderson, R.J. Reiter

Rev. Med. Virol., 30 (2020)

[\[104\]](#)

E. Koyuncu, H.G. Budayeva, Y.V. Miteva, D.P. Ricci, T.J. Silhavy, T. Shenk, I.M. Cristea

mBio, 5 (2014)

[\[105\]](#)

A.M. Pedrosa, R. Weinlich, G.P. Mognol, B.K. Robbs, J.P. Viola, A. Campa, G.P. Amarante-Mendes

J. Immunol., 184 (2010), pp. 3487-3494

[\[106\]](#)

Y. Shang, S.P. Xu, Y. Wu, Y.X. Jiang, Z.Y. Wu, S.Y. Yuan, S.L. Yao

Chin. Med. J., 122 (2009), pp. 1388-1393

[\[107\]](#)

S.C. Miller, S.R. Pandi-Perumal, A.I. Esquifino, D.P. Cardinali, G.J. Maestroni

Int. J. Exp. Pathol., 87 (2006), pp. 81-87

[\[108\]](#)

H.M. Abu-Soud, S.L. Hazen

Biochemistry, 40 (2001), pp. 10747-10755

[\[109\]](#)

I. Sliskovic, I. Abdulhamid, M. Sharma, H.M. Abu-Soud

Free Radic. Biol. Med., 47 (2009), pp. 1005-1013

© 2021 Elsevier Inc. All rights reserved.