

# Zinc Ionophore Activity of Quercetin and Epigallocatechin-gallate: From Hepa 1-6 Cells to a Liposome Model

Husam Dabbagh-Bazarbachi,<sup>†,⊥</sup> Gael Clergeaud,<sup>‡,⊥</sup> Isabel M. Quesada,<sup>§</sup> Mayreli Ortiz,<sup>‡</sup> Ciara K. O'Sullivan,<sup>\*,‡,||</sup> and Juan B. Fernández-Larrea<sup>\*,†</sup>

<sup>†</sup>Nutrigenomics Research Group, Department of Biochemistry and Biotechnology, and <sup>‡</sup>Nanobiotechnology & Bioanalysis Group, Department of Chemical Engineering, Universitat Rovira i Virgili, 43007 Tarragona, Spain

<sup>§</sup>Vascular Biology Laboratory, IMBECU-CONICET, Facultad de Ciencias Médicas, Universidad Nacional de Cuyo, 5500 Mendoza, Argentina

<sup>||</sup>Institució Catalana de Recerca i Estudis Avançats, 08010 Barcelona, Spain

**ABSTRACT:** Labile zinc, a tiny fraction of total intracellular zinc that is loosely bound to proteins and easily interchangeable, modulates the activity of numerous signaling and metabolic pathways. Dietary plant polyphenols such as the flavonoids quercetin (QCT) and epigallocatechin-gallate act as antioxidants and as signaling molecules. Remarkably, the activities of numerous enzymes that are targeted by polyphenols are dependent on zinc. We have previously shown that these polyphenols chelate zinc cations and hypothesized that these flavonoids might be also acting as zinc ionophores, transporting zinc cations through the plasma membrane. To prove this hypothesis, herein, we have demonstrated the capacity of QCT and epigallocatechin-gallate to rapidly increase labile zinc in mouse hepatocarcinoma Hepa 1-6 cells as well as, for the first time, in liposomes. In order to confirm that the polyphenols transport zinc cations across the plasma membrane independently of plasma membrane zinc transporters, QCT, epigallocatechin-gallate, or clioquinol (CQ), alone and combined with zinc, were added to unilamellar dipalmitoylphosphocholine/cholesterol liposomes loaded with membrane-impermeant FluoZin-3. Only the combinations of the chelators with zinc triggered a rapid increase of FluoZin-3 fluorescence within the liposomes, thus demonstrating the ionophore action of QCT, epigallocatechin-gallate, and CQ on lipid membrane systems. The ionophore activity of dietary polyphenols may underlay the raising of labile zinc levels triggered in cells by polyphenols and thus many of their biological actions.

**KEYWORDS:** clioquinol, epigallocatechin-gallate, flavonoids, liposomes, quercetin, zinc ionophores

## 1. INTRODUCTION

Quercetin (QCT), a water-insoluble flavonoid present in onions, nuts, and many other vegetables, and epigallocatechin-3-gallate (EGCG), a water-soluble flavonoid present in green tea, are among the most consumed and most studied polyphenols present in the human diet.<sup>1</sup> Flavonoids are considered bioactive micronutrients whose regular consumption, either as food components, or as dietary supplements and nutraceuticals,<sup>2</sup> entails benefits for human health, including prevention and amelioration of cancers,<sup>3</sup> diabetes, and cardiovascular<sup>4</sup> and neurodegenerative<sup>5</sup> diseases. Many of the health benefits of flavonoids have historically been ascribed to their antioxidant activity, which they exert directly by scavenging reactive oxygen species (ROS) and by chelating the redox-active transition metals iron and copper, which may act as ROS generators in biological systems.<sup>6</sup> Flavonoids also act as antioxidants indirectly by inhibiting redox-sensitive transcription factors and pro-oxidant enzymes as well as through induction of phase II and antioxidant enzymes.<sup>7</sup> However, it is currently believed that the levels of polyphenols achieved through ingestion are not enough to justify their wide array of biological actions. Beyond their antioxidant actions, flavonoids are also known to act as signaling molecules that, either directly or indirectly, interact with proteins and nucleic acids, thus modulating multiple cell signaling pathways, gene

transcription, metabolic fluxes, and cell fate including apoptosis.<sup>8,9</sup>

Diverse polyphenols have been shown able to form complexes with the redox-inactive transition metal zinc.<sup>10</sup> Zinc is an essential micronutrient for humans, the deficiency of which causes multiple dysfunctions, including alterations of glucidic and lipidic metabolisms.<sup>11</sup> Within cells, the vast majority of zinc cations (in concentrations usually ranging from 100 to 300  $\mu\text{M}$  for most cells) are tightly bound to proteins, functioning as a catalytic or structural component of an estimated 3000 mammalian proteins involved in virtually all cellular processes.<sup>12</sup> A minor fraction of intracellular zinc, termed labile zinc, exists in its free ionic form (picomolar concentrations) or loosely bound to proteins (in nanomolar concentrations). This pool of zinc is detectable by specific fluorophores with very high affinities for zinc cations at neutral pH such as Zinquin and FluoZin-3. Zinc ionophores such as pyrithione and clioquinol (CQ) have been used to increment labile zinc within cells and determine the fundamental roles that this zinc pool plays in cellular biology. Thus, free and labile zinc acts as second messenger molecule, which modulates the

Received: April 1, 2014

Revised: July 22, 2014

Accepted: July 22, 2014

Published: July 22, 2014

activity of many enzymes and thus signaling and metabolic pathways and cellular processes, including cell fate and apoptosis.<sup>13,14</sup> While many enzymes are inhibited by small elevations of zinc concentrations, others are activated.<sup>15,16</sup> Mammalian cells tightly control the subcellular distribution of zinc cations and the levels of labile zinc through the coordinated action of dedicated transmembrane zinc transporters and zinc-chelating metallothioneins.

Zinc from the extracellular milieu and from intracellular compartments enters the cytoplasm through 14 specialized transmembrane proteins of the ZIP/SLC39 family, whereas cytoplasmic extrusion of zinc toward organelles or the extracellular environment is performed by 10 transporters of the ZnT/SLC30 family, being ZnT1, located at the plasma membrane, the main regulator of cellular zinc efflux and export of excess zinc in most cells.<sup>17</sup> Within the cytoplasm, zinc may bind to metal free apo-metallothionein (apo-MT) to generate Zn–MT complexes. The apo-MT/Zn–MT ratio controls free and labile zinc concentrations. MT also serves as a ROS scavenger and heavy metal chelator, and the transcription of MT responds, in addition to zinc, to stress stimuli such as ROS, heavy metals, and proinflammatory cytokines.<sup>18</sup> In response to elevations of intracellular zinc, the zinc-sensor transcription factor MTF1 coordinately upregulates the expression of MT and ZnT1,<sup>19</sup> thus keeping zinc levels within a functional range. Excessively high levels of labile zinc are associated with cellular death through apoptosis.<sup>20</sup> Dysfunctions of MT and zinc transporters are promoting factors in cardiovascular diseases,<sup>21</sup> diabetes,<sup>22</sup> Alzheimer's disease,<sup>23</sup> and cancer.<sup>24</sup>

Several studies have shown that flavonoids affect zinc metabolism. For instance, rats fed during long periods with bicalain and rutin showed reduced hepatic levels of total zinc, as well as iron and copper, implying that flavonoids may sequester these metals and render them unavailable for absorption in a similar way as phytate.<sup>25</sup> Consistent with this, feeding obese rats with proanthocyanidins reverse dyslipidemia and lower protein levels of ZnT1 in the liver, reflecting lower levels of hepatic zinc.<sup>26</sup> Early *in vitro* studies showed that, in human intestinal Caco-2 cells, genistein enhanced the expression of MT, here regarded as an antioxidant enzyme,<sup>27</sup> while QCT increased the copper-induced expression of MT.<sup>28</sup> More recently, QCT was shown to enhance zinc uptake by Caco-2 cells, increasing total zinc accumulation and MT expression.<sup>29</sup> In contrast, grape seed flavonoids, produced a reduction in apical zinc uptake in Caco-2 cells, similar to that produced by phytate, whereas EGCG did not alter zinc absorption.<sup>30</sup> In prostate cancer cells, EGCG accelerated the accumulation of total zinc in the cytosol and mitochondria.<sup>31</sup> Two reports have shown that polyphenols may produce an increase of intracellular labile zinc. A water-soluble glycoside of the isoflavone genistin enhanced MT expression in human hepatocarcinoma HepG2 cells concomitantly increasing labile zinc and cellular death.<sup>32</sup> Furthermore, the stilbene resveratrol was shown to enhance total and labile zinc in normal human prostate epithelial cells, while not significantly affecting MT expression, and this was accompanied by increased cellular death. These authors suggested that the increment of labile zinc elicited by resveratrol might be due to the uptake of resveratrol–zinc complexes, followed by the dissociation of the complexes in the cytoplasm.<sup>33</sup>

Conversely, zinc may affect the bioactivity of flavonoids, as detailed in a few reports, including one that outlines the stimulating effect of zinc on the apoptotic effect of genistein in

osteoclastic cells.<sup>34</sup> Zinc also yields EGCG effective in protecting cultured rat hepatocytes against hepatotoxin-induced cell injury<sup>35</sup> and enhances the antiproliferative, proapoptotic effects of EGCG on various lines of prostate cancer cells.<sup>36</sup> Zinc was also shown to affect the uptake of EGCG by prostate carcinoma cells, where Zn–EGCG chelates were less internalized by cells than EGCG alone, while mixtures of EGCG with zinc enhanced the transport of EGCG into the cells. These authors also showed that zinc enhances the incorporation of EGCG into liposomes.<sup>37</sup>

We have previously reported that the water-soluble flavonoid EGCG and the water-insoluble flavonoid QCT profoundly alter zinc homeostasis in cultured human and mouse hepatocarcinoma cells. Whereas EGCG reduced the levels of total intracellular zinc and the expression of MT and ZnT1,<sup>38</sup> QCT enhanced total zinc accumulation as well as MT and ZnT1 expression (M. Bustos, personal communication, 2011, Universitat Rovira i Virgili). However, both QCT and EGCG dose-dependently prevented zinc-induced toxicity, suggesting that most zinc cations in the culture medium are rendered unavailable to cells due to their chelation by flavonoids and the formation of flavonoid–zinc concatemers, as shown for iron and copper complexed with diverse polyphenols.<sup>39</sup> In addition, both polyphenols enhanced cytoplasmic levels of Zinquin-detectable labile zinc, suggesting that a fraction of the flavonoid molecules in the culture medium formed complexes with zinc that cross the plasma membrane; that is, the flavonoids may also act as zinc ionophores, transporting zinc cations across the plasma membrane independently from zinc transporters.

The aim of this work was to evaluate the capacity of QCT and EGCG to act as zinc ionophores. Clioquinol (CQ) was also tested in this study as it is a synthetic antitumor drug recently reported to induce apoptosis in diverse cells lines by enhancing intracellular labile zinc and therefore inferred to act as a water-soluble zinc ionophore.<sup>40</sup> We evaluated the ability of QCT, EGCG, and CQ to chelate zinc cations and the subsequent formation of a complex with FluoZin-3, a fluorophore that displays a very high affinity for zinc cations ( $K_d = 15 \text{ nM}$ ).<sup>41</sup> The uptake of zinc by mouse hepatocarcinoma cells was measured fluorescently using FluoZin-3 in the presence and absence of QCT, EGCG, and CQ. This study was repeated using unilamellar liposomes with encapsulated FluoZin-3 to investigate whether the transport of the zinc cations across the cytoplasmic membrane to form a complex with FluoZin-3 was indeed enhanced by the presence of QCT, EGCG, or CQ or was simply due to the activity of cellular zinc transporters.

## 2. MATERIALS AND METHODS

**2.1. Chemicals.** The lipid 1,2-dipalmitoyl-*sn*-glycero-3-phosphocholine (DPPC), cholesterol, zinc chloride ( $\text{ZnCl}_2$ ), QCT, EGCG, CQ, dimethyl sulfoxide (DMSO), ethanol, and phosphate-buffered saline (0.01 M PBS, pH 7.4) were bought from Sigma–Aldrich. Cell-impermeant FluoZin-3 tetrapotassium salt and cell permeant FluoZin-3 AM were bought from Molecular Probes. A Simplicity 185 Millipore water system was used to obtain Milli-Q water ( $18.2 \text{ m}\Omega \cdot \text{cm}^{-1}$ ) for the preparation of buffers and liposomes. The compounds QCT, EGCG, and CQ were dissolved as 100  $\mu\text{M}$  solutions in 100% DMSO, aliquoted, and stored at  $-20^\circ\text{C}$ .  $\text{ZnCl}_2$  was stored as 1 M solution in ethanol/PBS (50%/50% v/v). FluoZin-3 indicators were used at 10  $\mu\text{M}$  in 100% DMSO.

**2.2. Cell Cultures and Treatments.** The mouse hepatoma cell line Hepa 1-6 was obtained from the European Collection of Cell Cultures (BW7756 ECACC) and propagated in Dulbecco's Modified Eagle medium (DMEM; BioWittaker) supplemented with 10% fetal

bovine serum (BioWittaker), 2 mM glutamine in 0.85% NaCl, 1000 U/mL penicillin/streptomycin, and 1.25 M HEPES. This medium contains  $4.9 \pm 0.2 \mu\text{M}$  zinc, as determined by flame atomic absorption spectroscopy (FAAS). Cells were cultured at  $37^\circ\text{C}$  in a humidified, 5%  $\text{CO}_2$ -enriched atmosphere and routinely split every 3–4 days at a 1:5 ratio upon reaching approximately 80% confluence. For treatments, cells at 80% confluence were detached with Accutase (Sigma–Aldrich) and resuspended at a density of  $5 \times 10^5$  cells/mL; 500  $\mu\text{L}$  of this cell suspension ( $25 \times 10^4$  cells) was then seeded per well in 24-well plates (Orange Scientific). Twenty four hours after plating, medium was removed, and the cells were treated by adding 100  $\mu\text{L}$  of fresh medium containing either 50  $\mu\text{M}$   $\text{ZnCl}_2$ , 100  $\mu\text{M}$  QCT, 100  $\mu\text{M}$  EGCG, 100  $\mu\text{M}$  CQ, or the combination of 50  $\mu\text{M}$   $\text{ZnCl}_2$  with each chelator for 1 and 4 h, respectively. As a control experiment, untreated cells were incubated just with medium and vehicle (final 0.1% DMSO and 0.05% ethanol).

**2.3. Measurements of Cytoplasmic Labile Zinc in Hepa 1-6 Cells.** The intracellular levels of free and labile zinc cations were measured as the fluorescence emission of cells upon loading them with the membrane-permeant zinc specific detector FluoZin-3, using fluorescence microscopy as described.<sup>42</sup> Briefly, following cell treatment (Section 2.2), culture media were replaced with a fresh one containing 1.5  $\mu\text{M}$  FluoZin-3 (AM, cell permeant) and incubated for 30 min at  $37^\circ\text{C}$ . This medium was then removed, and the cells were washed three times with PBS, and the zinc-dependent FluoZin-3 fluorescence within cells was measured using a Nikon Eclipse TE2000-S microscope, with excitation set at 494 nm and emission at 516 nm. Fluorescent intensities were quantified using the NIS-Elements AR software (Nikon Instruments) and the software ImageJ, a Java-based image processing program developed at the NIH (National Institutes of Health).<sup>43</sup>


**2.4. Liposomes as Cell Membrane Models.** Homogeneous populations of liposomes were prepared using a previously reported method.<sup>44</sup> FluoZin-3 in a final concentration of 3  $\mu\text{M}$  was mixed with 5 mL of PBS (0.01 M, pH 7.4) in a glass reactor protected from light-induced degradation, under stirring conditions and a blanket of argon gas. After 15 min, a mixture of DPPC and cholesterol (9:1 molar ratio) was added and maintained under stirring conditions and argon at  $25^\circ\text{C}$  for another 15 min. The homogeneous mixture was then treated with a rapid pH jump from pH 7.4 to pH 11 and then back to pH 7.4 within a 3 s frame, followed by an equilibration step of 25 min where lipid clusters curl into FluoZin-3 encapsulating liposomes. The resulting FluoZin-3-loaded liposomes were purified using a Sephadex G-100 size-separation column and used immediately.

Liposomes with encapsulated FluoZin-3 were separately incubated with 10  $\mu\text{M}$  QCT, EGCG, or CQ, in the presence and absence of 10  $\mu\text{M}$   $\text{ZnCl}_2$ . All the solutions were allowed to incubate at  $25^\circ\text{C}$  for 30 min before measuring their fluorescence. For the kinetic experiment, liposomes loaded with FluoZin-3 were added to three different cuvettes, the fluorescence was measured for 15 min, followed by addition of  $\text{ZnCl}_2$  (final 10  $\mu\text{M}$ ) to each cuvette, and the fluorescence emission was measured for another 15 min. Finally, QCT, EGCG, or CQ was added (final 10  $\mu\text{M}$ ) to each sample, and fluorescence emission was monitored over the duration of 1 h until the fluorescent intensity reached a plateau.

### 3. RESULTS

**3.1. QCT, EGCG, and CQ Increase the Cytoplasmic Labile Zinc in Hepa 1-6 Cells.** The increase of cytoplasmic labile zinc is modulated by the cellular zinc transporters, where the zinc ions are transported to the cytoplasm through specific channels of the ZIP family, bound to ionophore molecules that independently cross the lipid bilayer, or liberated from zinc-binding proteins such as metallothioneins (Figure 1).

In order to assess the effect of QCT, EGCG, CQ and zinc on cytoplasmic labile zinc, Hepa 1-6 cells were treated for 1 and 4 h with the chelators and supplemental zinc, and variations in the intracellular levels of labile zinc were measured as changes




**Figure 1.** Schematic representation of zinc homeostasis. Intracellular labile zinc is modulated by the coordinated activity of a large family of zinc transporters (ZnT and ZIP) and zinc-binding proteins, such as metallothionein or ionophore molecules.

in the FluoZin-3 fluorescence intensity as described in Section 2.3. Figure 2 shows the fluorescent images of the Hepa 1-6 cells after 1 and 4 h of treatment. No significant increase in fluorescence was obtained when QCT or EGCG were added to the culture medium without additional zinc. Only CQ insignificantly enhanced the fluorescence in these conditions, that is, with basal zinc concentration in the culture medium, which is roughly 5  $\mu\text{M}$ .


However, when 50  $\mu\text{M}$   $\text{ZnCl}_2$  was added, both QCT and EGCG doubled the amount of FluoZin-3-detectable zinc after 1 h, and CQ increased this pool of zinc 10-fold with respect to the control (50  $\mu\text{M}$   $\text{ZnCl}_2$  in the absence of any of QCT, EGCG, or CQ) (Figure 3), suggesting a slower ionophore action of the flavonoids as compared to CQ. After 4 h treatment with additional 50  $\mu\text{M}$   $\text{ZnCl}_2$ , all treatments triggered a significant increase in fluorescence intensity. In the case of the control, the increase of the cytoplasmic labile zinc is associated with the plasma membrane ZIP transporters, whereby zinc ions are transported into the cell. QCT doubled the amount of labile zinc attained with only 50  $\mu\text{M}$  zinc, EGCG quadrupled this value, and CQ increased it 7-fold.

A closer view of intracellular distribution of FluoZin-3 fluorescence (Figure 4) after 4 h of treatment shows a similar punctuated pattern of labile zinc for CQ, EGCG, and QCT, suggesting similar ways of action for the three compounds.

**3.2. Zinc Ionophore Activity of QCT, EGCG, and CQ Using Liposomes as Membrane Models.** Increases of cytoplasmic labile zinc levels triggered by CQ and pyrithione in a variety of cell lines have been attributed to their ionophore activity, that is, to the capacity of CQ–zinc and pyrithione–zinc complexes to cross the plasma membrane. To our knowledge, however, the classification of CQ and pyrithione as zinc ionophores is based on their functional effect in cells, that is, the rapid increase in Zinquin-detectable or FluoZin-3-detectable intracellular zinc, but no direct biochemical assay has



**Figure 2.** Effect of QCT, EGCG, and CQ on the cytoplasmic pool of labile zinc in Hepa 1-6 cells. Hepa 1-6 cells were first treated with 100  $\mu\text{M}$  QCT, EGCG, or CQ, in the presence or absence of 50  $\mu\text{M}$   $\text{ZnCl}_2$  for 1 and 4 h. The medium was then removed, and 3  $\mu\text{M}$  FluoZin-3 (AM, cell permeant) was added. After 30 min incubation, cells were washed and examined using a confocal fluorescence microscope. Control cells were treated with vehicle (final 0.05% ethanol, 0.1% DMSO). Scale bars are 50  $\mu\text{m}$ .




**Figure 3.** Intensity of FluoZin-3 fluorescence signal from images in Figure 2 was quantified using quanta program and considering an equal number of cells in each field. All values are mean  $\pm$  SEM of three independent experiments. Significant differences between treatments were determined using one-way ANOVA (Tukey test). \* $P \leq 0.05$ ; \*\* $P \leq 0.01$ .

been performed to discard the involvement of plasma membrane zinc importers or the origin of labile zinc from intracellular components on this effect. Furthermore, there is no report confirming that polyphenols are able to transport zinc

across the plasma membrane independently of cell transport mechanisms, such as zinc transporters or endocytosis. To directly prove the ionophore action of the flavonoids and CQ, we tested their capacity to transport zinc cations across the lipid bilayer of protein-free liposomes as a model system that mimics a cell membrane devoid of protein and polysaccharide fractions. Taking advantage of our previously reported method for the rapid preparation of liposomes,<sup>44</sup> 3  $\mu\text{M}$  concentration of cell-impermeant FluoZin-3 was encapsulated within unilamellar liposomes composed of DPPC/cholesterol in a 9:1 molar ratio. The resulting FluoZin-3-loaded liposomes were purified by passing the sample through a Sephadex G-100 size-separation column to remove the unencapsulated FluoZin-3 molecules. Dynamic light scattering (DLS) and  $\zeta$  potential analysis were performed to clearly confirm the presence of stable liposomes within a size range 1–2  $\mu\text{m}$  and surface charge around zero (Table 1).

The zinc ionophore activity of polyphenols was then tested as their capacity to transport zinc cations into the liposome cavity, interacting with the encapsulated zinc-dependent FluoZin-3 and consequently increasing the fluorescence signal within the liposomes (Figure 5).  $\text{ZnCl}_2$  (10  $\mu\text{M}$ ) was added to the liposomal suspension in the absence and presence of 10  $\mu\text{M}$  QCT, EGCG, and CQ, respectively, and zinc-dependent fluorescence intensity was measured over time. Following the addition of QCT, EGCG, or CQ, the Zn–polyphenol chelation complex is formed and transported across the bilayer, followed by interaction with the encapsulated FluoZin-3 probe, resulting in a significant and immediate increase in the fluorescence intensity.

Following the addition of 10  $\mu\text{M}$   $\text{ZnCl}_2$  to the liposomes, a very small increase in the fluorescence appears due to the presence of a few free FluoZin-3 molecules that were not removed during the purification process and represent the background fluorescent signal. Even so, the fluorescent signal remains very low as zinc ions alone cannot cross the liposome membrane. The results from the kinetic experiment demon-




**Figure 4.** Subcellular localization of FluoZin-3 detectable zinc in Hepa 1-6 cells after 4 h treatment in the same samples as in Figure 2 showed at a greater magnification. Scale bars are 10  $\mu\text{m}$ .

**Table 1.** Dynamic Light Scattering and  $\zeta$  Potential Measurements of Liposomes Loaded with FluoZin-3 before and after Treatments with 10  $\mu\text{M}$  Quercetin (QCT10), 10  $\mu\text{M}$  Epigallocatechin-3-gallate (EGCG10), or 10  $\mu\text{M}$  Clioquinol (CQ10) in the Presence and Absence of 10  $\mu\text{M}$  Zinc Chloride (Zn10)<sup>a</sup>

sample	size average ( $\mu\text{m}$ )	$\zeta$ potential (mV)
FluoZin-3-loaded liposomes	1.4 $\pm$ 0.3	-4.7 $\pm$ 2.5
FluoZin-3-loaded liposomes + Zn10	1.1 $\pm$ 0.7	-5.0 $\pm$ 5.9
FluoZin-3-loaded liposomes + QCT10	1.2 $\pm$ 0.5	-1.9 $\pm$ 4.1
FluoZin-3-loaded liposomes + EGCG10	1.5 $\pm$ 0.4	-6.9 $\pm$ 7.6
FluoZin-3-loaded liposomes + CQ10	1.4 $\pm$ 0.2	-3.0 $\pm$ 3.7
FluoZin-3-loaded liposomes + Zn10 + QCT10	1.8 $\pm$ 0.2	-1.2 $\pm$ 2.5
FluoZin-3-loaded liposomes + Zn10 + EGCG10	1.6 $\pm$ 0.3	-3.5 $\pm$ 3.2
FluoZin-3-loaded liposomes + Zn10 + CQ10	1.4 $\pm$ 0.2	-9.0 $\pm$ 2.7


<sup>a</sup>Final concentrations of solvents in the samples were 0.05% ethanol and 0.1% DMSO. Standard deviations were calculated from the mean data of a series of experiments ( $n \geq 3$ ).

strate that QCT, EGCG, and CQ present different ionophore properties. CQ showed the strongest ionophore activity, producing a 35-fold increase in the zinc-dependent FluoZin-3 fluorescence intensity. Moreover, the maximum fluorescence is achieved rapidly after reaching the equilibrium in less than 15 min. QCT and EGCG also display a high ionophore activity in the system, although to a lesser extent as compared to the strong ionophore CQ, with 8- and 16-fold increases in fluorescence signal observed for QCT and EGCG, respectively. It can also be observed that both QCT and EGCG required more time (>60 min) to achieve the plateau phase, displaying slower chelation and transport kinetics (Figure 6).




**Figure 6.** Effect of QCT, EGCG, and CQ on the uptake of zinc cations by liposomes. Zinc-dependent fluorescence emission of FluoZin-3 encapsulated within liposomes treated with zinc cations, polyphenols, and CQ. The fluorescence emission ( $\lambda_{\text{ex}} = 494 \text{ nm}$ ;  $\lambda_{\text{em}} = 516 \text{ nm}$ ) of purified FluoZin-3-loaded liposomes was recorded continuously. Background fluorescence (0–15 min) was negligible (I). Upon the addition of 10  $\mu\text{M}$   $\text{ZnCl}_2$  to the liposomal suspensions (II), a small fluorescence signal was detected, presumably due to the presence of trace amounts of unencapsulated FluoZin-3 in the liposomal solutions. At time point 30 min, 10  $\mu\text{M}$  quercetin (QCT10), epigallocatechin-3-gallate (EGCG10), clioquinol (CQ10), or vehicle (control, final 0.1% DMSO) were added to the liposomal solutions, and the fluorescence was monitored for one additional hour (III).


Confocal microscopy analysis was also performed in order to visualize and corroborate that the fluorescence produced by the interaction between zinc and FluoZin-3 was attributable to fluorescence in the inner part of the liposomes. As can be seen in Figure 7, fluorescence is only observed when the combination of  $\text{ZnCl}_2$  with QCT, EGCG, or CQ is present, and the fluorescent signal comes from the inside part of the liposomes and not from the lipid membrane or the background



**Figure 5.** Schematic design of the FluoZin-3-loaded liposomes and the ionophore-like effect interpretation.



**Figure 7.** Effect of QCT, EGCG, and CQ on the uptake of zinc cations by liposomes. Three-dimensional confocal microscopy images of zinc-dependent fluorescence emission of FluoZin-3 loaded within liposomes treated with zinc cations, polyphenols, and CQ.



**Figure 8.** Effect of QCT, EGCG, and CQ on the uptake of zinc cations by liposomes. Transmission electron micrographs of liposomes with encapsulated FluoZin-3 after treatment with  $ZnCl_2$ , QCT, EGCG, and CQ.

solution. To further support this, stability studies were carried out in order to check whether the QCT, EGCG, or CQ can destabilize and break the lipid vesicles. All the liposomes were characterized using transmission electron microscopy (TEM), DLS, and  $\zeta$  potential to check that their stability was maintained following exposure.

As shown in Figure 8, TEM images of the liposomes before and after treatments clearly demonstrate that liposome stability was not affected with the morphology and mean size (1–2  $\mu\text{m}$ ) being maintained. Moreover, the DLS and  $\zeta$  potential results presented in Table 1, confirmed that the size of the liposomes and their surface charge were not significantly affected by the addition of zinc and/or the ionophores, thus demonstrating that FluoZin-3-loaded liposomes were not destabilized and their around-zero charge, due to the zwitterionic nature of main lipid component DPPC, was maintained, thus confirming that the fluorescence signal is due to transport of the Zn–QCT/EGCG/CQ complex across the lipid membrane.

#### 4. DISCUSSION

The consequences that zinc chelation by flavonoids may have on zinc availability to the cells may in principle be dual: sequestering or ionophore, as shown for other well-characterized zinc-binding compounds.<sup>45</sup> On the one hand, the formation of zinc–flavonoid complexes may render zinc unavailable for cells, as do other dietary phytochemicals such as phytates<sup>46</sup> and zinc-chelator drugs such as TPEN that induce zinc deficiency *in vitro* and *in vivo*.<sup>47</sup> Metal chelating

therapy using CQ has been proposed for neurodegenerative disorders that course with high levels of metal accumulation such as Alzheimer's and Parkinson.<sup>48</sup> On the other hand, flavonoids may form water-insoluble membrane-permeant complexes with zinc that cross the plasma membrane, and thereby act as zinc ionophores. The ionophore effect of zinc-binding compounds has been characterized for pyrithione and CQ. Both drugs chelate zinc cations and, when applied to cells, trigger a rapid increase of the intracellular pool of zinc that is detectable with different fluorophores such as Zinquin or FluoZin-3. Thereafter, it is assumed that these chelators form membrane-permeable complexes that are transported into the cell and that, once within the cell, chelator–zinc complexes dissociate into the single compounds due to the low concentration of intracellular free and labile zinc, thus providing labile zinc cations. Although FluoZin-3 has been widely accepted as a fluorophore that specifically interacts with zinc,<sup>49–52</sup> a recent report has indicated that this marker may in fact suffer from a lack of specificity.<sup>53</sup> However, this has no impact on the proof-of-concept study reported here as zinc is the only ion present and studied and thus no interfering effect from other ions will occur. We have shown here that treatment of Hepa1-6 cells with zinc together with QCT, EGCG, or CQ elicits a rapid and drastic increase in FluoZin-3-detectable intracellular zinc. The same effect was previously observed using Zinquin upon treatments of HepG2 cells with combinations of zinc and EGCG or zinc and QCT.<sup>38</sup> In these cells, the upregulation of MT and ZnT1 by zinc was

enhanced by QCT. In contrast, EGCG decreased the intracellular zinc accumulation. Similar to QCT, the stilbene resveratrol efficiently chelates zinc in solution and enhances total and Zinquin-detectable cytoplasmic zinc in cultured human prostate epithelial cells, and this correlated with the antiproliferative action of resveratrol on cells.<sup>33</sup>

While an increasing effort has been made to understand the interaction of flavonoids with lipid bilayers,<sup>54</sup> no report has been published reporting the use of liposomes to demonstrate the zinc ionophore activity of polyphenols. Liposomes have been widely used as the simplest cell membrane systems in order to study the ionophore activity of molecules across the lipidic bilayer.<sup>55</sup> We have herein used a liposomal system to prove that zinc can transverse lipid bilayers when combined with flavonoids. It is not necessary to evoke the intervention of zinc transporters in the plasma membrane or the mobilization of zinc from intracellular compartments to account for the elevation of intracellular zinc levels in cells treated with flavonoids. The flavonoid-dependent transport of zinc cations into the liposomal cavity also implies that polyphenols may cross biological membranes when conjugated with metal cations. The mechanisms by which polyphenols enter the cells are largely unknown, but our results imply that complexation with metals may increase the bioavailability of polyphenols to cells.

There are several reports that strongly suggest that the demonstrated zinc ionophore effect on polyphenols will be observed in real physiological conditions if studied in vivo. Lee et al.<sup>56</sup> in 2002 has reported that a minor part of the EGCG found in plasma conserved its native form, and it has also been reported that the polyphenol metabolites still maintain their ability to chelate and form complexes with metal ions<sup>30</sup> and furthermore that nanomolar concentrations of polyphenols or their metabolites are able to modulate some metabolic pathways,<sup>57,58</sup> as does labile zinc in a picomolar to nanomolar concentration range.<sup>14</sup> While all these reports point toward the same ionophore effect being observed under physiological conditions, in a recent report by Oyama et al.,<sup>59</sup> the authors suggest the dual effect of CQ depending on the extracellular zinc concentration, where the known ionophore effect of CQ was only observed when extracellular zinc was available, and when zinc was not available in the extracellular environment, CQ could cross the membrane and chelate the intracellular zinc ions. To this end, ongoing work is looking at extending the proof-of-concept reported and demonstrated here with both liposomes and cellular models to true physiological conditions exploring the interactions between a range of polyphenols and polyphenol metabolites and zinc in a lower concentration range of picomolar to micromolar.

In conclusion, we have demonstrated that QCT, EGCG, and CQ rapidly increase intracellular labile zinc in Hepa 1-6 cells and that they function as ionophores for zinc in a liposomal system. Thus, natural flavonoids can be added to an arsenal of drugs that may be used to modulate zinc homeostasis and regulate zinc-dependent biological pathways.

## AUTHOR INFORMATION

### Corresponding Authors

\*E-mail: ciara.osullivan@urv.cat.

\*E-mail: juanbautista.fernandez@urv.cat.

### Author Contributions

<sup>†</sup>Husam Dabbagh-Bazarbachi and Gael Clergeaud contributed equally to this work.

## Funding

This work was supported by grant AGL2008-00387 from the Spanish Ministry of Education and Science. H.D. was granted a predoctoral grant from the University Rovira i Virgili.

## Notes

The authors declare no competing financial interest.

## ABBREVIATIONS USED

DPPC, 1,2-dipalmitoyl-*sn*-glycero-3-phosphocholine; EGCG, (–)-epigallocatechin-3-gallate; FluoZin-3, 2-[2-[2-[bis-(carboxylatomethyl)amino]-5-methoxyphenoxy]ethoxy]-4-(2,7-difluoro-3-oxido-6-oxo-4a,9a-dihydroxanthren-9-yl)-anilino]acetate; MRE, metal response element; MT, metallothionein; MTF-1, MRE-binding transcription factor-1; TPEN, *N,N,N',N'*-tetrakis(2-phridylmethyl) ethylenediamine; Zinquin, ethyl (2-methyl-8-*p*-toluenesulfonamido-6-quinoloxo); ZIP, ZRT/IRT related protein (SLC39 family of zinc transporters); ZnT, zinc transporter (SLC30 family of zinc transporters)

## REFERENCES

- (1) Aron, P. M.; Kennedy, J. A. Flavan-3-ols: Nature, occurrence and biological activity. *Mol. Nutr. Food Res.* **2008**, *52*, 79–104.
- (2) Egert, S.; Rimbach, G. Which sources of flavonoids: Complex diets or dietary supplements? *Adv. Nutr.* **2011**, *2*, 8–14.
- (3) Ramos, S. Cancer chemoprevention and chemotherapy: Dietary polyphenols and signalling pathways. *Mol. Nutr. Food Res.* **2008**, *52*, 507–526.
- (4) Jagtap, S.; Meganathan, K.; Wagh, V.; Winkler, J.; Hescheler, J.; Sachinidis, A. Chemoprotective mechanism of the natural compounds, epigallocatechin-3-O-gallate, quercetin and curcumin against cancer and cardiovascular diseases. *Curr. Med. Chem.* **2009**, *16*, 1451–1462.
- (5) Williams, R. J.; Spencer, J. P. E. Flavonoids, cognition, and dementia: Actions, mechanisms, and potential therapeutic utility for Alzheimer disease. *Free Radical Biol. Med.* **2012**, *52*, 35–45.
- (6) Selvaraj, S.; Krishnaswamy, S.; Devashya, V.; Sethuraman, S.; Krishnan, U. M. Flavonoid–metal ion complexes: A novel class of therapeutic agents. *Med. Res. Rev.* **2014**, *34*, 677–702.
- (7) Na, H.-K.; Surh, Y.-J. Modulation of Nrf2-mediated antioxidant and detoxifying enzyme induction by the green tea polyphenol EGCG. *Food Chem. Toxicol.* **2008**, *46*, 1271–1278.
- (8) Williams, R. J.; Spencer, J. P. E.; Rice-Evans, C. Flavonoids: Antioxidants or signalling molecules? *Free Radical Biol. Med.* **2004**, *36*, 838–849.
- (9) Ramos, S. Effects of dietary flavonoids on apoptotic pathways related to cancer chemoprevention. *J. Nutr. Biochem.* **2007**, *18*, 427–442.
- (10) Le Nest, G.; Caille, O.; Woudstra, M.; Roche, S.; Burlat, B.; Belle, V.; Guigliarelli, B.; Lexa, D. Zn–polyphenol chelation: Complexes with quercetin, (+)-catechin, and derivatives: II Electrochemical and EPR studies. *Inorg. Chim. Acta* **2004**, *357*, 2027–2037.
- (11) Prasad, A. S. Discovery of human zinc deficiency: 50 years later. *J. Trace Elem. Med. Biol.* **2012**, *26*, 66–69.
- (12) Maret, W. Zinc biochemistry: From a single zinc enzyme to a key element of life. *Adv. Nutr.* **2013**, *4*, 82–91.
- (13) Fukada, T.; Yamasaki, S.; Nishida, K.; Murakami, M.; Hirano, T. Zinc homeostasis and signaling in health and diseases. *J. Biol. Inorg. Chem.* **2011**, *16*, 1123–1134.
- (14) Maret, W. Metals on the move: Zinc ions in cellular regulation and in the coordination dynamics of zinc proteins. *Biometals* **2011**, *24*, 411–418.
- (15) Wilson, M.; Hogstrand, C.; Maret, W. Picomolar concentrations of free zinc(II) ions regulate receptor protein-tyrosine phosphatase  $\beta$  activity. *J. Biol. Chem.* **2012**, *287*, 9322–9326.
- (16) Maret, W. Inhibitory zinc sites in enzymes. *Biometals* **2013**, *26*, 197–204.

- (17) Cousins, R. J.; Liuzzi, J. P.; Lichten, L. A. Mammalian zinc transport, trafficking, and signals. *J. Biol. Chem.* **2006**, *281*, 24085–24089.
- (18) Krężel, A.; Maret, W. Thionein/metallothionein control Zn(II) availability and the activity of enzymes. *J. Biol. Inorg. Chem.* **2008**, *13*, 401–409.
- (19) Laity, J. H.; Andrews, G. K. Understanding the mechanisms of zinc-sensing by metal-response element binding transcription factor-1 (MTF-1). *Arch. Biochem. Biophys.* **2007**, *463*, 201–210.
- (20) Truong-Tran, A. Q.; Ho, L. H.; Chai, F.; Zalewski, P. D. Cellular zinc fluxes and the regulation of apoptosis/gene-directed cell death. *J. Nutr.* **2000**, *130*, 1459S–1466S.
- (21) Little, P. J.; Bhattacharya, R.; Moreyra, A. E.; Korichneva, I. L. Zinc and cardiovascular disease. *Nutrition* **2010**, *26*, 1050–1057.
- (22) Myers, S. A.; Nield, A.; Myers, M. Zinc transporters, mechanisms of action and therapeutic utility: Implications for type 2 diabetes mellitus. *J. Nutr. Metab.* **2012**, *2012*, 173712.
- (23) Watt, N. T.; Whitehouse, I. J.; Hooper, N. M. The role of zinc in Alzheimer's disease. *Int. J. Alzheimer's Dis.* **2010**, *2011*, 971021.
- (24) Hogstrand, C.; Kille, P.; Nicholson, R. I.; Taylor, K. M. Zinc transporters and cancer: A potential role for ZIP7 as a hub for tyrosine kinase activation. *Trends Mol. Med.* **2009**, *15*, 101–111.
- (25) Gao, Z.; Xu, H.; Chen, X.; Chen, H. Antioxidant status and mineral contents in tissues of rutin and baicalin fed rats. *Life Sci.* **2003**, *73*, 1599–1607.
- (26) Baiges, I.; Palmfeldt, J.; Blade, C.; Gregersen, N.; Arola, L. Lipogenesis is decreased by grape seed proanthocyanidins according to liver proteomics of rats fed a high fat diet. *Mol. Cell Proteomics* **2010**, *9*, 1499–513.
- (27) Kuo, S.-M.; Leavitt, P. S. Genistein increases metallothionein expression in human intestinal cells, Caco-2. *Biochem. Cell Biol.* **1999**, *77*, 79–88.
- (28) Kuo, S.-M.; Huang, C.-T.; Blum, P.; Chang, C. Quercetin cumulatively enhances copper induction of metallothionein in intestinal cells. *Biol. Trace Elem. Res.* **2001**, *84*, 1–10.
- (29) Sreenivasulu, K.; Raghu, P.; Nair, K. M. Polyphenol-rich beverages enhance zinc uptake and metallothionein expression in Caco-2 cells. *J. Food Sci.* **2010**, *75*, H123–H128.
- (30) Kim, E.-Y.; Pai, T.-K.; Han, O. Effect of bioactive dietary polyphenols on zinc transport across the intestinal Caco-2 cell monolayers. *J. Agric. Food Chem.* **2011**, *59*, 3606–3612.
- (31) Yang, J.; Yu, H.; Sun, S.; Zhang, L.; Das, U.; Ruan, H.; He, G.; Shen, S. Mechanism of free Zn(2+) enhancing inhibitory effects of EGCG on the growth of PC-3 cells: Interactions with mitochondria. *Biol. Trace Elem. Res.* **2009**, *131*, 298–310.
- (32) Chung, M. J.; Kang, A.-Y.; Lee, K. M.; Oh, E.; Jun, H.-J.; Kim, S.-Y.; Auh, J. H.; Moon, T.-W.; Lee, S.-J.; Park, K.-H. Water-soluble genistin glycoside isoflavones Up-regulate antioxidant metallothionein expression and scavenge free radicals. *J. Agric. Food Chem.* **2006**, *54*, 3819–3826.
- (33) Zhang, J. J.; Wu, M.; Schoene, N. W.; Cheng, W.-H.; Wang, T. T. Y.; Alshatwi, A. A.; Alsaif, M.; Lei, K. Y. Effect of resveratrol and zinc on intracellular zinc status in normal human prostate epithelial cells. *Am. J. Physiol.: Cell Physiol.* **2009**, *297*, C632–C644.
- (34) Uchiyama, S.; Yamaguchi, M. Genistein and zinc synergistically stimulate apoptotic cell death and suppress RANKL signaling-related gene expression in osteoclastic cells. *J. Cell. Biochem.* **2007**, *101*, 529–542.
- (35) Kagaya, N.; Kawase, M.; Maeda, H.; Tagawa, Y.-i.; Nagashima, H.; Ohmori, H.; Yagi, K. Enhancing effect of zinc on hepatoprotectivity of epigallocatechin gallate in isolated rat hepatocytes. *Biol. Pharm. Bull.* **2002**, *25*, 1156–1160.
- (36) Sun, S.-l.; He, G.-q.; Yu, H.-n.; Yang, J.-g.; Borthakur, D.; Zhang, L.-c.; Shen, S.-r.; Das, U. N. Free Zn<sup>2+</sup> enhances inhibitory effects of EGCG on the growth of PC-3 cells. *Mol. Nutr. Food Res.* **2008**, *52*, 465–471.
- (37) Yang, J. G.; Yu, H. N.; Sun, S. L.; Zhang, L. C.; He, G. Q.; Das, U. N.; Ruan, H.; Shen, S. R. Epigallocatechin-3-gallate affects the growth of LNCaP cells via membrane fluidity and distribution of cellular zinc. *J. Zhejiang Univ., Sci., B* **2009**, *10*, 411–21.
- (38) Quesada, I. M.; Bustos, M.; Blay, M.; Pujadas, G.; Ardèvol, A.; Salvadó, M. J.; Bladé, C.; Arola, L.; Fernández-Larrea, J. Dietary catechins and procyanidins modulate zinc homeostasis in human HepG2 cells. *J. Nutr. Biochem.* **2011**, *22*, 153–163.
- (39) Hider, R. C.; Liu, Z. D.; Khodr, H. H. Metal chelation of polyphenols. In *Methods in Enzymology*; Lester, P., Ed.; Academic Press: Waltham, MA, 2001; Vol. 335, pp 190–203.
- (40) Ding, W.-Q.; Liu, B.; Vaught, J. L.; Yamauchi, H.; Lind, S. E. Anticancer activity of the antibiotic clioquinol. *Cancer Res.* **2005**, *65*, 3389–3395.
- (41) Li, Y.-M.; Shi, J.; Wu, X.; Luo, Z.-F.; Wang, F.-L.; Guo, Q.-X. Tracing of intracellular zinc(II) fluorescence flux to monitor cell apoptosis by using FluoZin-3AM. *Cell Biochem. Funct.* **2009**, *27*, 417–423.
- (42) Yu, H.; Zhou, Y.; Lind, S. E.; Ding, W.-Q. Clioquinol targets zinc to lysosomes in human cancer cells. *Biochem. J.* **2009**, *417*, 133–139.
- (43) Collins, T. J. ImageJ for microscopy. *Biotechniques* **2007**, *43*, 25–30.
- (44) Genç, R. k.; Ortiz, M.; O'Sullivan, C. K. Curvature-tuned preparation of nanoliposomes. *Langmuir* **2009**, *25*, 12604–12613.
- (45) Ding, W.-Q.; Yu, H.-J.; Lind, S. E. Zinc-binding compounds induce cancer cell death via distinct modes of action. *Cancer Lett.* **2008**, *271*, 251–259.
- (46) Wise, A. Phytate and zinc bioavailability. *Int. J. Food Sci. Nutr.* **1995**, *46*, 53–63.
- (47) Li, B.; Tan, Y.; Sun, W.; Fu, Y.; Miao, L.; Cai, L. The role of zinc in the prevention of diabetic cardiomyopathy and nephropathy. *Toxicol. Mech. Methods* **2013**, *23*, 27–33.
- (48) Bareggi, S. R.; Cornelli, U. Clioquinol: Review of its mechanisms of action and clinical uses in neurodegenerative disorders. *CNS Neurosci. Ther.* **2012**, *18*, 41–46.
- (49) Muylle, F. R.; Adriaensen, D.; Coen, W.; Timmermans, J.-P.; Blust, R. Tracing of labile zinc in live fish hepatocytes using FluoZin-3. *Biomaterials* **2006**, *19*, 437–450.
- (50) Wessels, I.; Haase, H.; Engelhardt, G.; Rink, L.; Uciechowski, P. Zinc deficiency induces production of the proinflammatory cytokines IL-1 $\beta$  and TNF $\alpha$  in promyeloid cells via epigenetic and redox-dependent mechanisms. *J. Nutr. Biochem.* **2013**, *24*, 289–297.
- (51) Kim, J.-H.; Jeon, J.; Shin, M.; Won, Y.; Lee, M.; Kwak, J.-S.; Lee, G.; Rhee, J.; Ryu, J.-H.; Chun, C.-H.; Chun, J.-S. Regulation of the catabolic cascade in osteoarthritis by the zinc-ZIP8-MTF1 axis. *Cell* **2014**, *156*, 730–743.
- (52) Chevallet, M.; Jarvis, L.; Harel, A.; Luche, S.; Degot, S.; Chapuis, V.; Boulay, G.; Rabilloud, T.; Bouron, A. Functional consequences of the over-expression of TRPC6 channels in HEK cells: Impact on the homeostasis of zinc. *Metallomics* **2014**, *6*, 1269–1276.
- (53) Landero Figueroa, J. A.; Subramanian Vignesh, K.; Deepe, G. S., Jr.; Caruso, J. Selectivity and specificity of small molecule fluorescent dyes/probes used for the detection of Zn<sup>2+</sup> and Ca<sup>2+</sup> in cells. *Metallomics* **2014**, *6*, 301–315.
- (54) Efimova, S. S.; Ostroumova, O. S. Effect of dipole modifiers on the magnitude of the dipole potential of sterol-containing bilayers. *Langmuir* **2012**, *28*, 9908–9914.
- (55) Sankaram, M. B.; Shastri, B. P.; Easwaran, K. R. K. Interaction of carrier ionophores with phospholipid vesicles. *Biochemistry* **1987**, *26*, 4936–4941.
- (56) Lee, M.-J.; Maliakal, P.; Chen, L.; Meng, X.; Bondoc, F. Y.; Prabhu, S.; Lambert, G.; Mohr, S.; Yang, C. S. Pharmacokinetics of tea catechins after ingestion of green tea and (–)-epigallocatechin-3-gallate by humans: Formation of different metabolites and individual variability. *Cancer Epidemiol., Biomarkers Prev.* **2002**, *11*, 1025–1032.
- (57) Schroeter, H.; Bahia, P.; Spencer, J. P. E.; Sheppard, O.; Rattray, M.; Cadenas, E.; Rice-Evans, C.; Williams, R. J. (–)Epicatechin stimulates ERK-dependent cyclic AMP response element activity and up-regulates GluR2 in cortical neurons. *J. Neurochem.* **2007**, *101*, 1596–1606.



(58) Vauzour, D.; Vafeiadou, K.; Rice-Evans, C.; Williams, R. J.; Spencer, J. P. E. Activation of pro-survival Akt and ERK1/2 signalling pathways underlie the anti-apoptotic effects of flavanones in cortical neurons. *J. Neurochem.* **2007**, *103*, 1355–1367.

(59) Oyama, T. M.; Ishida, S.; Okano, Y.; Seo, H.; Oyama, Y. Clioquinol-induced increase and decrease in the intracellular  $Zn^{2+}$  level in rat thymocytes. *Life Sci.* **2012**, *91*, 1216–1220.