# Crave Contro

# **Crave Control**



Distributed By: HormoneSynergy

www.hormonesynergy.com
Portland, OR



# **Clinical Applications**

- Nutritional Support for Carbohydrate, Alcohol & Drug Cravings\*
- Supports Healthy Weight (by reducing carbohydrate cravings)\*
- Improve Sense of Wellbeing and Energy\*
- Supports Healthy Serotonin Levels\*

**Crave Control** contains key amino acids to support the biosynthesis of neurotransmitters involved in appetite control, carbohydrate or fat cravings, and mood. Chromium is present to support healthy glucose metabolism and support food intake regulation.\*

All Homone Synergy™ Nutraceuticals Formulas Meet or Exceed cGMP Quality Standards

# **Discussion**

**5-Hydroxytryptophan** ("**5-HTP**") is a naturally-occurring amino acid precursor to serotonin. Numerous studies during the '90s, including those randomized, double-blind, and placebo-controlled, confirmed the safety and efficacy of 5-HTP in reducing appetite and food intake in obese healthy and non-insulin-dependent diabetic individuals. [1,2,3] A 2006 study in mice concluded, "5-HTP-induced anorexia may be mediated by facilitation of leptin secretion." [4] Vitamins B6 and C are important cofactors in the 5-HTP to serotonin pathway. Among the several serotonin receptors thus identified, the 5HT2C receptors are suspected in control of food intake. Mice without this receptor exhibit increased food intake and become obese.\* [5]

**DL-Phenylalanine (DLPA)** is a combination of the d- and the l- forms of this essential amino acid. Phenylalanine suppresses appetite by regulating the release of cholecystokinin, which in turns signals satiety in the brain. D-phenylalanine increases endorphins, while L-phenylalanine is an amphetamine-like stimulatory compound. DLPA has been found to elevate mood, curb appetite and reduce pain.\*

**L-Tyrosine**, an essential amino acid is needed for conversion into the catecholamine neurotransmitters stress depletes: dopamine, norepinephrine, and epinephrine. It is also a precursor for thyroxine. Doctors use tyrosine as a mood elevator, to increase alertness after sleep deprivation and as an appetite suppressant; although support for the latter appears anecdotal.\*<sup>[6]</sup>

**L-Glutamine**, well–recognized for gut and immune support, has also been espoused to reduce carbohydrate cravings and support alcohol withdrawal, although the mechanism of action for these benefits is not known.\*<sup>[7,8]</sup>

**Chromium**, as chromium picolinate is widely used to optimize insulin function; thereby preventing swings in blood glucose levels that may be responsible for carbohydrate cravings. The mineral in the form present was indeed demonstrated to reduce carbohydrate cravings in a double-blind, placebo-controlled study.\*[9,10]

<sup>\*</sup>These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.



# **Supplement Facts**

Serving Size: 4 Capsules Servings Per Container: 60

An	nount Per Serving	%Daily Value
Vitamin C (ascorbic acid)	500 mg	833%
Vitamin B6 (as pyridoxine HCI)	37.5 mg	1875%
Chromium (as TRAACS® chromium nicotinate glycinate ch	elate) 60 mcg	50%
DL-Phenylalanine	1 g	**
L-Tyrosine	750 mg	**
L-Glutamine	375 mg	**
5-HTP (5-hydroxytryptophan)(from <i>Griffonia simplicifolia</i> )	(seed) 75 mg	**
** Daily Value not established.		

Other Ingredients: HPMC (capsule), stearic acid, magnesium stearate, medium-chain triglyceride oil, and silica.



TRAACS and the Albion Medallion design are registered trademarks of Albion Laboratories, Inc.

## **Directions**

Take four capsules before breakfast and four capsules before 3 PM, or use as directed by your healthcare practitioner. Do not take after 3 PM.

Consult your healthcare practitioner prior to use if you have, or suspect you have, a medical condition or are taking prescription drugs for depression, migraines, Parkinson's disease, or psychiatric disorders. Not for use by children. Do not use if tamper seal is damaged.

## **Does Not Contain**

Wheat, gluten, yeast, soy, animal or dairy products, fish, shellfish, peanuts, tree nuts, egg, ingredients derived from genetically modified organisms (GMOs), artificial colors, artificial sweeteners, or artificial preservatives.

# References

- 1. Cangiano C, et al. Eating behavior and adherence to dietary prescriptions in obese adult subjects treated with 5-hydroxytryptophan. Am J Clin Nutr 1992 Nov;56(5):863-7
- 2. Cangiano C Effects of oral 5-hydroxy-tryptophan on energy intake and macronutrient selection in non-insulin dependent diabetic patients. Int J Obes Relat Metab Disord 1998 Jul;22(7):648-544
- 3. Amamoto T, Sarai K. On the tryptophan-serotonin metabolism in manic-depressive disorders. Changes in plasma 5-HT and 5-HIAA levels and urinary 5-HIAA excretion following oral loading of L-5HTP in patients with depression. *Hiroshima J Med Sci.* 1976 Sep;25(2-3):135-40 [PMID: 1088369]
- 4. Yamada J, Sugimoto Y, Ujikawa M. Involvement of leptin in hypophagia induced by the serotonin precursor 5-hydroxytryptophan (5-HTP) in mice. *Biol Pharm Bull.* 2006 Mar;29(3):557-9 [PMID: 16508167]
- 5. Rogers PJ, Blundell JE. Reanalysis of the effects of phenylalanine, alanine, and aspartame on food intake in human subjects. *Physiol Behav.* 1994 Aug;56(2):247-50 [PMID: 7938234]
- 6. http://web.indstate.edu/thcme/mwking/aminoacidderivatives.html {accessed 09 July 07}
- 7. Goodwin, F. APA Psychiatric News, Dec 5, 1986 in Atkins, R. Dr. Atkins Vita-Nutient Solution. Simon & Schuster, NY 1998 p169
- 8. Rogers,L., Pelton, R. Quarterly Journal of Studies of Alcohol, 1957;18(4):581-87 in Atkins, R. Dr. Atkins Vita-Nutient Solution. Simon & Schuster, NY 1998 p169
- 9. Broadhurst CL, Domenico P. Clinical studies on chromium picolinate supplementation in diabetes mellitus--a review. *Diabetes Technol Ther.* 2006 Dec:8(6):677-87 [PMID: 17109600]
- 10. Docherty JP, et al. A double-blind, placebo-controlled, exploratory trial of chromium picolinate in atypical depression: effect on carbohydrate craving. *J Psychiatr Pract.* 2005 Sep;11(5):302-14 [PMID: 16184071]

# **Cautions**

Do not take if you are, or suspect you are, pregnant or if you are lactating.

\*These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.