

WELLVITA VITAMIN C TIME RELEASE

PROPRIETARY NAME AND DOSAGE FORM

WELLVITA VITAMIN C Time Release (tablet)

COMPOSITION

Each time released tablet contains: Active ingredients: Vitamin C (ascorbic acid) 1 000 mg.

PHARMACOLOGICAL CLASSIFICATION

D 34.11 Vitamins

Complementary medicine: Health supplement

INDICATIONS

Wellvita™ Vitamin C Time Release tablets provide the gradual release of vitamin C over time. This essential vitamin contributes to collagen formation for the normal function of gums, teeth, skin and cartilage. It also supports normal immune system function during and after intense physical stress.

CONTRA-INDICATIONS

Do not use this medicine if you are hypersensitive (allergic) to any of the ingredients in the formulation. Not suitable for children younger than 13 years, unless under medical supervision.

WARNINGS AND SPECIAL PRECAUTIONS

If you suffer from any serious ailments (such as kidney stones) or conditions (such as blood-iron disorders or sickle cell disease); or if you are taking any prescribed medication (including warfarin), please check with your healthcare provider before taking this medicine.

- No studies on the effect on the ability to drive and use machines have been performed. Do not drive or operate machinery until you know how it affects you.
- Porphyria: Safety has not been established.

Nutritional supplementation should not replace a balanced diet. Do not exceed the recommended dose without consulting a healthcare provider.

PREGNANCY AND LACTATION

If you are pregnant or breastfeeding, please consult your doctor, pharmacist or other healthcare provider for advice before taking this medicine.

INTERACTIONS

If you are taking other medicines regularly, including complementary or traditional medicines, or start taking any additional medicines while taking this medicine, consult your healthcare provider for advice. Vitamin C might affect the metabolism of certain medicines. These can include, but are not limited to protease inhibitors, estrogen, fluphenazine, warfarin and statins. Consult your healthcare provider for more information.

DOSAGE AND DIRECTIONS FOR USE

Adults: Take one tablet daily with a meal and plenty of water, or as directed by your healthcare provider.

SIDE EFFECTS

Side effect may include heartburn, nausea, diarrhoea, headache or stomach cramps. Not all side effects reported for this medicine are included in this leaflet. Should your general health worsen or if you experience any untoward effects while taking this medicine, discontinue use immediately and consult your doctor, pharmacist or other healthcare provider for advice.

KNOWN SYMPTOMS OF OVERDOSAGE AND PARTICULARS OF ITS TREATMENT

Symptoms of overdose can include nausea, diarrhoea and stomach cramps. In case of an overdose or suspected overdose, contact your healthcare provider immediately. Treatment is symptomatic and supportive.

IDENTIFICATION

Yellow tablet.

PRESENTATION

30 or 60 Tablets are packed in a plastic container.

STORAGE INSTRUCTIONS

Store all medicines out of reach of children. Store in a cool dry place below 25 °C and keep away from direct sunlight. Keep the container tightly closed.

NAME AND BUSINESS ADDRESS OF THE APPLICANT

Alveta Healthcare (Pty) Ltd, 1 Greenwich Grove, Station Road, Rondebosch, 7700. Company registration number: 2004/021899/07. Pharmacy audit number: Y53008.

DATE OF PUBLICATION OF THE PACKAGE INSERT

1 September 2020.

WELLVITA VITAMIN C TIME RELEASE

HANDELSMERKNAAM EN DOSEERVORM

WELLVITA VITAMIN C Time Release (stadige vrystellingstabelt)

SAMESTELLING

Elke stadige vrystellingstabelt bevat: Aktiewe bestanddeel: Vitamien C (askorbiensuur) 1 000 mg.

FARMAKOLOGIESE KLASSIFIKASIE

D 34.11 Vitamienes

Komplementêre medisyne: Gesondheidsaanvulling

INDIKASIES

Wellvita™ Vitamin C Time Release tablette stel geleidelik vitamien C vry oor 'n tydperk. Hierdie essensiële vitamien dra by tot kollageenvorming vir die normale funksie van tandvleis, tande, vel en kraakbeen. Dit ondersteun ook normale immuunstelsel-funksie gedurende en na fisiese inspanning.

KONTRA-INDIKASIES

Vermyn as jy hipersensitief (allergies) is vir enige van die bestanddele in die formule. Dit is nie geskik vir kinders jonger as 13 jaar nie, behalwe onder die toesig van 'n geneesheer.

WAARSKUWINGS EN SPESIALE VOORSORGMATREËLS

As jy enige voorgeskrewe medikasie (insluitend warfarien) gebruik of aan enige ernstige siektes (soos bv. bloed-yster verstourings of sekelsel siekte) of toestande ly (soos bv. nierstene), raadpleeg jou geneesheer voordat jy hierdie medisyne begin gebruik.

- Geen studies is gedoen om die effek wat die medisyne het op bestuur en die gebruik van masjinerie vas te stel nie. Moenie bestuur of masjinerie gebruik totdat jy weet hoe die medisyne jou beïnvloed nie.
- Porfirie: Veiligheid is nie vasgestel nie.

Voedingsaanvullings moet nie 'n gebalanseerde dieet vervang nie. Moenie die aanbevole dosis oorskry sonder om 'n geneesheer te raadpleeg nie.

SWANGERSKAP EN BORSVOEDING

Raadpleeg jou geneesheer as jy swanger is of borsvoed voordat jy hierdie medisyne gebruik.

INTERAKSIES

As jy ander medisyne gereeld gebruik, insluitend aanvullende (komplementêre) of tradisionele medisyne, of as jy enige addisionele medisyne begin neem terwyl jy hierdie medisyne gebruik, raadpleeg jou geneesheer vir raad. Vitamien C kan moontlik veroorsaak dat sekere medisyne vinniger of stadiger gemetaboliseer word. Hierdie medikasie kan, onder andere, insluit: Protease inhibeerdere, estrogeen, fluphenasien, warfarien, statins, ens. Raadpleeg jou geneesheer vir meer inligting.

DOSES EN GEBRUIKSAANWYSINGS

Volwassenes: Neem een tablet daagliks met 'n ete en genoegsame water, of soos jou geneesheer voorskryf.

NEWE-EFFEKTE

Nuwe-effekte kan sooibrand, naarheid, diarree, hoofpyn en maagkrampe insluit. Sommige aangemelde nuwe-effekte is nie in hierdie voubiljet ingesluit nie. Indien jy enige onaangename nuwe-effekte ervaar of voel dat jou algemene gesondheid verswak, hou onmiddellik op om die medisyne te gebruik en kontak jou geneesheer.

BEKENDE SIMPTOME VAN OORDOSERING EN BESONDERHEDE VAN DIE BEHANDELING

Simptome van oordosering kan naarheid, diaree en maagkrampe insluit. In geval van oordosering of indien oordosering vermoed word, kontak jou geneesheer onmiddellik. Behandeling is simptomaties en ondersteunend.

IDENTIFIKASIE

Geel tablet.

AANBIEDING

30 of 60 Tablette is verpak in 'n plastiek houer.

BERGINGSANWYSINGS

Berg alle medisyne buite bereik van kinders. Berg in 'n koel, droë plek onder 25 °C en hou weg van direkte sonlig. Hou die houer dig toe.

NAAM EN BESIGHEIDSADRES VAN DIE APPLIKANT

Alveta Healthcare (Pty) Ltd, Greenwich Grove 1, Station weg, Rondebosch, 7700. Maatskappy registrasie nommer: 2004/021899/07. Apteek oudit nommer: Y53008.

DATUM VAN VOUBILJET PUBLIKASIE

1 September 2020.