

Citrus

Fungicide ♦ Bactericide ♦ Nematicide

WARHAMMER

BLACKSMITH
BIOSCIENCE



Blacksmith BioScience
25018 Spring Ridge Dr
Spring TX 77386
www.blacksmithbio.com

For More
Information
Please Contact
Your Territory
Representative

Florida & SE US

Jack Kilgore
239-707-7677 cell
g8trmanjek@comcast.net

Western US

Tim Lichatowich
503-705-0384 cell
timlich@gmail.com

Pacific NW

Tony Shepherd
503-791-8517 cell
tonyshepherd499@gmail.com



WarHammer™ is a powerful crop management tool in CITRUS trees for use against diseases and nematodes

WarHammer is a high concentration of a natural bio-polymer that produces a powerful disease and nematode suppressing action. When applied, WarHammer induces a systemic response triggering the plant's natural defence mechanisms, allowing it to fend off harmful diseases and nematodes.

Soil diseases suppressed include *Fusarium*, *Rhizoctonia*, *Pythium*, *Phytophthora*, *Sclerotinia*, *Verticillium* and other soil borne diseases.


Foliar diseases effectively suppressed: Powdery and Downy Mildew, *Botrytis*, *Erwinia amylovora*, *Sclerotinia*, *Alternaria*, *Monolinia*, *Pseudomonas sp* including *P. syringae*, *Xanthomonas axonopodis* (citrus canker) and *Xanthomonas arboricola pv. Juglandis* and other foliar diseases.

Nematodes suppressed: Root knot nematode (*Meloidogyne spp.*), Root lesion nematode (*Pratylenchus spp.*), Reniform nematode (*Rotylenchulus spp.*), Sting nematode (*Belonolaimus spp.*), Stem nematode (*Anguina spp.*) and Citrus nematode (*Tylenchulus spp.*).

WarHammer is an excellent tool for integrated pest management (IPM)

- Perfect rotational partner
- No Resistance issues
- Compatible with fertilizers
- Dissolves easily into solution
- 0-day PHI
- Non toxic

Nematicidal effect of abamectin, boron, chitosan, hydrogen peroxide and *Bacillus thuringiensis* against citrus nematode on Valencia orange trees
MAM El-Saedy, Sandy E Hammad and Sherin FA Awd Allah
Alexandria University, Alexandria, Egypt
2011



Disease Application Rates:

Soil Application:

4-12 oz/acre through drip line or other irrigation.

Foliar Spray

4-12 oz/acre as a foliar spray via airblast, backpack, aerial or other spray application
Repeat application every 1-4 weeks

Nematode Application Rates:

Initial Soil Application:

12 oz/acre through drip line or other irrigation.

Follow Up Soil Drench

4-8 oz/acre through irrigation or other soil application
Repeat application every 3 weeks



WARHAMMER
2.5 gal jug

Not registered in all states. Always read and follow label directions. Forge is a registered trademark of Blacksmith BioScience 2023

www.blacksmithbio.com