

Ingestion of bioactive collagen hydrolysates enhance facial skin moisture and elasticity and reduce facial aging signs in a randomized double-blind placebo-controlled clinical study

Naoki Inoue¹, Fumihito Sugihara¹, Xuemin Wang²

¹Peptide division, Nitta Gelatin Inc., Osaka, Japan.

²Shanghai Skin Disease Hospital, Shanghai, China

Correspondence author: Naoki Inoue

Peptide division, Nitta Gelatin Inc.

2-22 Futamata, Yao, Osaka, 581-0024, Japan.

Phone: +81-72-948-8252, FAX: +81-72-949-6014,

E-mail: na-inoue@nitta-gelatin.co.jp

Key Words: Clinical study, Collagen hydrolysate, Collagen peptide, Skin health, Bioactive peptides, Skin roughness.

ABSTRACT

BACKGROUND: Several human studies have demonstrated occurrence of two major collagen peptides, prolyl-hydroxyproline (Pro-Hyp), hydroxyprolyl-glycine (Hyp-Gly), in human peripheral blood. Some *in vitro* studies have demonstrated that Pro-Hyp and Hyp-Gly has chemotaxis to dermal fibroblast and enhance cell proliferation. Additionally, Pro-Hyp enhances the production of hyaluronic acid by dermal fibroblast. These findings suggest that contents of Pro-Hyp and Hyp-Gly in blood are important factors to exhibit the efficacy of collagen hydrolysates on skin health.

RESULTS: We conducted a randomized double-blind placebo-controlled clinical trial of ingestion of two types of collagen hydrolysates, which are composed of different contents of the bioactive dipeptides, Pro-Hyp and Hyp-Gly, to investigate their effects on skin condition improvement. Improvement in skin conditions such as skin moisture, elasticity, wrinkles, and roughness were compared with a placebo group at baseline, and four and eight weeks after the start of the trial. In addition, the safety of dietary supplementation with these peptides was evaluated by blood test. Collagen hydrolysate with a higher content of bioactive collagen peptides (H-CP) showed significant and more improvement than the collagen hydrolysate with a lower content of bioactive collagen peptides (L-CP) and the placebo, in facial skin moisture, elasticity (R2), wrinkles and roughness, compared with the placebo group. In addition, there were no adverse events during the trial.

CONCLUSION: This study demonstrated that the collagen hydrolysate with a higher content of Pro-Hyp and Hyp-Gly, showed more improvement in facial skin condition, including facial skin moisture, elasticity, wrinkles and roughness.

INTRODUCTION

There are many products that have beneficial effects on skin health available in the current health food market. Collagen hydrolysates have been developed over the past two decades as supplements or cosmeceutical products for use worldwide. Although a number of studies have demonstrated the efficacy of collagen hydrolysates on skin condition, little is known regarding what peptides derived from collagen hydrolysates function as bioactive peptides and have physiological effects, which is fundamental information for the maintenance of healthy facial skin.

Denatured collagen forms a substance called gelatin, which when treated by enzymatic hydrolysis results in what are called collagen hydrolysates. Collagen hydrolysates are soluble in water at ambient temperature due to low molecular weight, and possess no gelation ability. This high solubility of collagen hydrolysates allow for the development of products in drink- and jelly-stick-form.

Pharmacological bioavailability trials revealed that two types of collagen dipeptides, prolyl-hydroxyproline (Pro-Hyp) and hydroxyprolyl-glycine (Hyp-Gly), were available at high concentrations for several hours in the human blood stream after oral administration¹⁻⁵. It has been demonstrated that ¹⁴C-labelled Pro-Hyp reaches the skin and bone tissues rapidly after ingestion by mice⁶. Moreover, in a clinical study, Pro-Hyp was identified in the urine after collagen hydrolysate intake⁷. These findings suggest that Pro-Hyp and Hyp-Gly are stable and relatively resistant to peptidases in the blood^{4,8}, and are able to reach the skin tissues.

In addition, some *in vitro* studies demonstrated the physiological function of Pro-Hyp and Hyp-Gly in skin dermal fibroblasts. Pro-Hyp stimulated chemotaxis of dermal fibroblasts⁹ and both Pro-Hyp and Hyp-Gly enhanced cell proliferation activity^{10,11}. Additionally, it was observed that Pro-Hyp enhanced the production of hyaluronic acid in dermal fibroblasts¹¹.

Pro-Hyp and Hyp-Gly involvement in such physiological roles may be important to improve the efficacy of collagen hydrolysates on the maintenance of skin health. The current study, a randomized double-blind placebo-controlled clinical trial, was carried out to evaluate the efficacy of two types of collagen hydrolysates with differing contents of the bioactive dipeptides, Pro-Hyp and Hyp-Gly.

MATERIALS AND METHODS

Investigational products

The placebo, maltodextrin TK-16, was purchased from Matsutani Chemical Industry Co., Ltd. (Itami, Japan). Two forms of collagen hydrolysates derived from fish gelatin, which were composed of different ratios of free-formed Pro-Hyp and Hyp-Gly, were used in this study. One form of collagen hydrolysate (L-CP) had a low ratio of dipeptide-to-product content, with about 0.1 g kg⁻¹ of product. The other form of collagen hydrolysate (H-CP) had a high ratio of dipeptide-to-product content, with more than 2 g kg⁻¹ of product. These products were provided by Nitta Gelatin Inc. (Osaka, Japan), and are commercially available under the Wellnex brand. Each 5 g test sample was packed in an aluminum sachet and was indistinguishable from all subjects and investigators.

Study Design

This clinical study was conducted in the Shanghai Skin Disease Hospital (Shanghai, China), under the supervision of Dr. Xuemin Wang, MD. Randomized administration of the products was carried out in 85 Chinese female subjects who were shown to have no medical issues by blood test performed prior to the study.

The randomized double-blind placebo-controlled study consisted of three groups: Placebo, L-CP, and H-CP. Participants were randomly assigned to one of the three groups in a 1:1:1 ratio using a computer generated randomization schedule. This study was conducted from February to April in 2012. At the start of the trial, each group contained 28 or 29 subjects. Five gram samples were ingested orally in hot milk, coffee, or any other beverages, once a day after dinner for 8 weeks. Efficacy was assessed at baseline, week 4, and week 8. The amount of daily protein except for collagen peptides was not confined in the study but sustainable intake amount was continued through this trial.

This study was conducted in accordance with the ethical principles of the Declaration of Helsinki. The study protocol was approved by the ethics committee of Shanghai Skin Disease Hospital, and written informed consent was obtained from all subjects participating in the study. Selection criteria were: age between 35 to 55 years, subject conscious of their dry and rough skin, Body Mass Index less than 30, not regularly using other supplements or health foods, no treatment with sex hormones over the prior three months, and non-pregnant. The subjects were advised to avoid excessive eating, drinking, exercise, strong sunburn, change in lifestyle, and change cosmetics.

Physiological measurements of the skin

Instrumental measurements of skin condition were evaluated at three points: at baseline prior to regular ingestion (baseline), and after 4 weeks and 8 weeks of ingestion. The subjects washed off their makeup by conventional methods, and were acclimatized for 30 minutes in the waiting lounge at a constant

temperature of $20\pm 2^{\circ}\text{C}$ and humidity of $50\pm 5\%$ before facial skin evaluation.

1 Skin moisture

The change of the dielectric constant measured by an electrical capacitance method was used as an estimate of the amount of skin moisture at the cheek and canthus using a Corneometer CM820 (Courage and Khazaka, Germany). Three measurements were taken and averaged.

2 Skin elasticity

Skin elasticity was measured by the suction method using a Cutometer SEM575 (Courage and Khazaka). Decompression suction was carried out for 5 seconds with a pressure of 300mbr and a mouth diameter of 2 mm. The return rate, R2 (skin elasticity: $Ua1/Uf1$), after expansion was assessed at the cheek and canthus.

3 Skin wrinkles and roughness

Analysis of the cutaneous surface of the area from the cheek to the canthus was conducted using a VisioFace SSA (Skin Surface Analysis, Courage and Khazaka) on the following items: number of wrinkles, wrinkle area, wrinkle depth, and roughness.

Statistical analysis

Comparison of skin moisture, elasticity, and VisioFace SSA data at different time points within a group were carried out with paired Student's *t*-test. Comparison between the two experimental (H-CP and L-CP) and placebo groups was performed using one-way analysis of variance (ANOVA) with Tukey's *post hoc* test for evaluation of significance. Comparison of skin moisture and elasticity between the groups was performed using the difference of these variables before ingestion and after 4 weeks (changing rate week 4) or 8 weeks (changing rate week 8) of ingestion.

Significance was defined as $p < 0.05$ using the data analysis software SPSS Ver.13.0. Each value was expressed as the mean \pm standard deviation (SD).

RESULTS

Panel demographics

Five subjects dropped out over the course of the study, due mainly to difficulty in visiting the hospital. There was no significant difference in age between the groups (Table 1). None of the subjects involved in the study demonstrated any dietary problems.

Skin moisture

Skin moisture results are summarized in Table 2. Skin moisture at the cheek and canthus in both the L-CP and H-CP groups showed a significant increase between baseline and weeks 4 and 8 ($p < 0.05$), while the placebo group did not show such an increase. In the L-CP group, skin moisture at the canthus was significantly higher than the placebo group by week 8. On the other hand, both cheek and canthus skin moisture in the H-CP group was significantly higher by week 8 ($p < 0.05$) when compared to the placebo group.

The change of skin moisture from baseline, namely changing rate (%) week 4 and 8, at the cheek and canthus in both the L-CP and H-CP groups showed a significant increase compared to the placebo group ($p < 0.05$).

Moreover, the change in skin moisture from baseline in the H-CP group was significantly greater ($p < 0.05$) at the cheek by week 8 and at the canthus by weeks 4 and 8, when compared to the L-CP group. Additionally, the changing rate of H-CP showed a 2-fold increase in the L-CP group by week 8 in cheek moisture and by weeks 4 and 8 in canthus.

Skin Elasticity (R2)

Skin elasticity (R2) results are summarized in Table 3. The placebo group showed elasticity of the canthus decreased significantly between baseline and weeks 4 and 8. The L-CP group showed no significant improvement in facial skin elasticity between baseline and weeks 4 and 8, and no significant differences between the placebo group at weeks 4 and 8. On the other hand, in the H-CP group, elasticity of the cheek increased significantly between baseline and weeks 4 and 8, as well as the elasticity of the canthus by week 8. Skin elasticity of both the cheek and canthus in the H-CP group was significantly higher ($p < 0.05$) than in the placebo group by week 8. Moreover, improvement of elasticity from baseline in the H-CP group was significantly higher ($p < 0.05$) than the placebo group by week 4 at the canthus and by week 8 at both the cheek and canthus. Furthermore, there was a significant difference in change rate of elasticity improvement at both the cheek and canthus between the L-CP and H-CP groups by week 8 ($p < 0.05$).

Skin Surface analysis by VisioFace SSA

Skin surface analysis results by VisioFace SSA are summarized in Table 4. In the L-CP group,

wrinkle area by weeks 4 and 8 was reduced significantly ($p < 0.05$), and roughness also improved significantly ($p < 0.05$) by week 8, when compared to baseline.

On the other hand, the H-CP group showed significant improvement compared to baseline in many categories, including the number of wrinkles by week 8, and wrinkle area, wrinkle depth, and roughness by weeks 4 and 8. Moreover, comparison between the H-CP and placebo groups showed significant differences ($p < 0.05$) in the number of wrinkles by week 8, and both wrinkle depth and roughness by weeks 4 and 8.

Additionally, there were significant differences ($p < 0.05$) between the H-CP and L-CP groups, including the number of wrinkles and wrinkle depth by week 8, and roughness by weeks 4 and 8.

Blood test

Blood test analysis results are shown in Table 5. Each value at baseline and after eight weeks of ingestion was within the limits of standard values. Furthermore, no adverse effects were observed during the clinical trial.

DISCUSSION

The present study demonstrated that ingestion of H-CP, which contains a higher content of the free-formed bioactive peptides, Pro-Hyp and Hyp-Gly, resulted in significantly better improvements in facial skin conditions compared to ingestion of L-CP, which has a lower content of these bioactive peptides. These results suggest that, despite using the same raw material, it may be possible to control the effects of collagen hydrolysates on facial skin conditions by modifying the manufacturing process and thus the dipeptide content. Previous reports have demonstrated the effects of Pro-Hyp and Hyp-Gly on skin dermal fibroblasts as signal transducers, which can stimulate metabolism, migration, proliferation, and production of hyaluronic acid⁹⁻¹¹. In addition, these dipeptides are absorbed into the blood by peptide transporters of the small intestinal epithelial cells in the human digestive and absorption process¹². Taking into account the bioavailability of these oligopeptides, we hypothesize that it may be possible to enhance uptake of bioactive peptides like Pro-Hyp and Hyp-Gly by increasing the concentration of free-formed bioactive peptides in collagen hydrolysate products. Another type of collagen hydrolysate product, which we have previously reported on, contains more than 3 g kg⁻¹ of Pro-Hyp and Hyp-Gly and may have similar or improved effectiveness in enhancing facial skin moisture, elasticity (R2) and roughness, with as little as half the ingested dose (2.5 g) utilized in the present study¹³. On the other hand, we need to consider an effect of beverage co-ingested with collagen hydrolysate for better absorption of collagen bioactive peptides. In the presence study, we reflected the actual use of powder type of collagen hydrolysate by ingestion with tea, coffee, juice, milk, kind of hot soup like miso soup, etc. Further studies are needed to better understanding the optimum combinations with drink type and general food to enhance the functional effects of collagen hydrolysate.

Skin moisture and elasticity depends on the condition of the extracellular matrix, which consists of primarily collagen, hyaluronic acid, and elastin. In an *in vitro* study using human dermal fibroblast cells, Ohara *et al.*¹¹ reported that Pro-Hyp enhanced cell proliferation and hyaluronic acid synthesis with up-regulated hyaluronic synthase 2 (HAS2) mRNA levels. In addition, they demonstrated that Pro-Hyp stimulates phosphorylation of signal transducer and activator of transcription 3 (STAT3), which is a fundamental intracellular signaling factor¹⁰. Recently, we have reported the daily oral administration of Pro-Hyp + Hyp-Gly improved skin barrier dysfunction and moisture in HR-1 hairless mice¹⁴. These reports suggest that Pro-Hyp and Hyp-Gly have a crucial effect in improving the barrier function to enhance skin moisture. We hypothesize that Pro-Hyp and Hyp-Gly stimulated production of hyaluronic acid in the dermis. Hyaluronic acid has been shown to play crucial roles in skin moisture and elasticity¹⁵. Additionally, several animal studies demonstrated that oral intake of collagen hydrolysates stimulated the synthesis of type I collagen and other extracellular matrix molecules^{16, 17, 18}.

Regarding the degree of the efficacy between the cheek and the canthus, moisture and elasticity were slightly better in the canthus. In general, the elasticity and thickness of human skin depends on age and measurement site¹⁹.

In the present study, the H-CP group showed improvement in the number of wrinkles and depth of wrinkles by VisioFace SSA.

E. Proksch *et al.*²⁰ have shown that the synthesis of procollagen Type I and elastin, components of the dermal extracellular matrix, led to a pronounced, statistically significant reduction in eye wrinkle volume in a double blind clinical trial. Their data support the idea that a decline in the number of eye wrinkles and wrinkle depth around the eye area effectively improves eye wrinkles, which was similar to the results of the present study (data not shown)²¹.

Regarding the effect of collagen hydrolysate on facial spots, we have previously reported in a clinical study that collagen hydrolysates help reduce ultraviolet (UV) spots after four weeks of ingestion²². Gu *et al.*²³ reported that hyaluronan plays a beneficial role by interacting with fibroblasts to enhance epidermal morphogenesis in a co-culture system. Okawa *et al.*²⁴ suggested that induced hyaluronic acid in dermal

fibroblasts followed by oral administration of collagen hydrolysate may provide beneficial effects on maintaining epidermal and dermal homeostasis in mice. Additionally, Le Vu *et al.*²⁵ demonstrated that Pro-Hyp induced an increase in expression of Krtap and Krt genes in keratinocytes in co-culture with fibroblasts. These findings suggest that Pro-Hyp may affect signaling to change the phenotype of keratinocytes through the regulation of dermal cells.

Further studies are needed to better understand the mechanisms of the bioactive peptides, Pro-Hyp and Hyp-Gly, which may be associated with their bioavailability. The findings would contribute not only to better understanding of collagen hydrolysate but also to further the understanding of fundamental mechanisms in anti-aging.

CONCLUSIONS

The present study demonstrates that both L-CP and H-CP are effective supplements for the improvement in skin moisture and roughness in women who were conscious of their dry and rough skin. Fortified collagen hydrolysate, H-CP, demonstrated a greater improvement in skin elasticity and reducing wrinkles on facial skin.

The present study is the first of its kind to demonstrate that there is a significant difference between conventional collagen hydrolysate and new types of collagen hydrolysate with higher contents of specific bioactive dipeptides such as Pro-Hyp and Hyp-Gly for improvement of human skin condition.

REFERENCES

- Iwai K, Hasegawa T, Taguchi Y, Morimatsu F, Sato K, Nakamura Y, Higashi A, Kido Y, Nakabo Y, and Ohtsuki K, Identification of food-derived collagen peptides in human blood after oral ingestion of gelatin hydrolysates. *J Agric Food Chem* **53**:6531-6536(2005).
- Ohara H, Matsumoto H, Ito K, Iwai K and Sato K, Comparison of quantity and structures of hydroxyproline-containing peptides in human blood after oral ingestion of gelatin hydrolysates from different sources. *J Agric Food Chem* **55**:1532-1535(2007).
- Ichikawa S, Morifuji M, Ohara H, Matsumoto H, Takeuchi Y and Sato K, Hydroxyproline-containing dipeptides and tripeptides quantified at high concentration in human blood after oral administration of gelatin hydrolysate. *Int J Food Sci Nutr* **61**:52-60(2010).
- Shigemura Y, Akaba S, Kawashima E, Park YE, Nakamura Y and Sato K, Identification of a novel food-derived collagen peptide, hydroxyprolyl-glycine, in human peripheral blood by pre-column derivatization with phenyl isothiocyanate. *Food Chem* **129**:1019-1024(2011).
- Sugihara F, Inoue N, Kuwamori M and Taniguchi M, Quantification of hydroxyprolyl-glycine (Hyp-Gly) in human blood after ingestion of collagen hydrolysate. *J Biosci Bioengi* **113**:202-203(2012).
- Kawaguchi T, Nanbu NP and Kurokawa M, Distribution of prolylhydroxyproline and its metabolites after oral administration in rats. *Biol Pharm Bull* **35**:422-427(2012).
- Meilman EB, Urivetzky MM and Rapoport MC, Urinary hydroxyproline peptides. *J Clin Invest* **42**:40-50(1963).
- Liu C, Sugita K, Nihei K, Yoneyama K and Tanaka H, Absorption of hydroxyproline-containing peptides in vascularly perfused rat small intestine *in situ*. *Biosci Biotechnol Biochem* **73**:1741-1747(2009).
- Postlethwaite EA, Seyer MJ and Kang HA, Chemotactic attraction of human fibroblasts to type I, II, and III collagens and collagen-derived peptides. *Proc Natl Acad Sci USA* **75**:871-875(1978).
- Shigemura Y, Iwai K, Morimatsu F, Iwamoto T, Mori T, Oda C, Taira T, Park YE, Nakamura Y and Sato K, Effect of prolyl-hydroxyproline (Pro-Hyp), a food-derived collagen peptide in human blood, on growth of fibroblasts from mouse skin. *J Agric Food Chem* **57**:444-449(2009).
- Ohara H, Ichikawa S, Matsumoto H, Akiyama M, Fujimoto N, Kobayashi T and Tajima S, Collagen-derived dipeptide, proline-hydroxyproline, stimulates cell proliferation and hyaluronic acid synthesis in cultured human dermal fibroblasts. *J Dermatol* **37**:330-338(2010).
- Adibi AS, The oligopeptide transporter (Pept-1) in human intestine: biology and function. *Gastroenterology* **113**:332-340(1997).
- Sugihara F, Inoue N and Wang X, Clinical effects of ingesting collagen hydrolysate on facial skin properties. *Jpn Pharmacol Ther* **43**:67-70(2015).
- Shimizu J, Asami N, Kataoka A, Sugihara F, Inoue N, Kimira Y, Wada M and Mano H, Oral collagen-derived dipeptides, prolyl-hydroxyproline and hydroxyprolyl-glycine, ameliorate skin barrier dysfunction and alter gene expression profiles in the skin. *Biochem Biophys Res Commun* **456**:626-630(2015).
- Wiest L, Kerscher M, Native hyaluronic acid in dermatology - results of an expert meeting. *J Dtsch Dermatol Ges* **6**:176-80(2008).
- Matsuda N, Koyama Y, Hosaka Y, Ueda H, Watanabe T, Araya T, Irie S and Takehana K, Effects of ingestion of collagen peptide on fibrils and glycosaminoglycans in the dermis. *J Nutr Sci Vitaminol* **52**:211-215(2006).

17. Liang J, Pei X, Zhang Z, Wang N, Wang J and Li Y, The protective effects of long-term oral administration of marine collagen hydrolysate from chum salmon on collagen matrix homeostasis in the chronological aged skin of Sprague-dawley male rats. *J Food Sci* **75**:H230-238(2010).
18. Zague V, Freitas DV, Costa Rosa DM, Castro DAG, Jaeger GR and Machado-Santelli MG, Collagen hydrolysate intake increases skin collagen expression and suppresses matrix metalloproteinase 2 activity. *J Med Food* **14**:618-624(2011).
19. Takema Y, Yorimoto Y, Kawai M and Imokawa G, Age-related changes in the elastic properties and thickness of human facial skin. *Br J Dermatol* **131**:641-648(1994).
20. Proksch E, Schunck M, Zague V, Segger D, Degwert J and Oesser S, Oral intake of specific bioactive collagen peptides reduces skin wrinkles and increases dermal matrix synthesis. *Skin Pharmacol Physiol* **27**:113-119(2014).
21. Inoue N, Sugihara F, Koizumi S, Efficacy of collagen peptide ingestion on skin. A bible for measurement and evaluation on skin, *Technical Information Institute* text No.1717. Chapter 6, Section 4(2013).
22. Sugihara F, Inoue N, Clinical effects of collagen hydrolysates ingestion on UV-induced pigmented spots of human skin: A preliminary study. *Health Sciences* **28**:2,153-156(2012).
23. Gu H, Huang L, Wong PY and Burd A, HA modulation of epidermal morphogenesis in an organotypic keratinocyte-fibroblast co-culture model. *Exp Dermatol* **19**:e336-339(2010).
24. Okawa T, Yamaguchi Y, Takada S, Sakai Y, Numata N, Nakamura F, Nagashima Y, Ikezawa Z and Aihara M, Oral administration of collagen tripeptide improves dryness and pruritus in the acetone-induced dry skin model. *J Dermatol Sci* **66**:136-143(2012).
25. Le Vu P, Takatori R, Iwamoto T, Akagi Y, Satsu H, Totsuka M, Chida K, Sato K, Shimizu M, Effects of food-derived collagen peptides on the expression of keratin and keratin-associated protein genes in the Mouse Skin. *J Pharm Pharmacol* **28**:5,227-235(2015).