

BreedPlus

Hi Form BreedPlus offers essential support for weanlings, yearlings and broodmares. Includes an excellent level and balance of calcium and phosphorous. This plays a role in normal growth and maintenance of bones.

BreedPlus is a complete amino acid, vitamin and high available calcium formula with an emphasis on essential minerals. Contains trace elements and will assist in maintaining the health of pregnant and lactating mares, weanlings and yearlings.



ALL IN ONE GREAT PRODUCT THE BEST START

Our BreedPlus formula has been specifically designed for breeding stock. Broodmares, breeding stallions, foals and yearlings. This formula is designed to support the growth and development of the foetus and young horse, the mare through her pregnancy and lactation stages as well as the servicing stallion.

Research has shown that developmental orthopedic diseases such as OCD, cervical vertebral malformations, angular deformities, contracted tendons and phytitis can be caused by many factors such as genetic predisposition, management practices, diseases and dietary influences. Good nutrition may assist with these types of conditions.

DOSAGE RATES

300kg : 12.5gm (2/3 large scoop)
 500kg : 25.5gm (1.5 large scoop)
 600+ kg : 34gm (2 large scoop)
 Mix well into slightly damp feed.



MILK THISTLE

Safe to use throughout pregnancy and lactation, it has an array of actions. Mainly known for its liver protective effects, in reproducing horses it enables sufficient detoxification of waste products and toxins and increases the effectiveness of antioxidants and anti-inflammatory mediators. Especially targeted for stallions and broodmares to significantly reduce the toxin load and therefore may increase fertility. In addition, as the name suggests, Milk Thistle is a galactagogue meaning it promotes milk production, which is important for the lactating mare.



BIOEQUUS

The synergy between prebiotics and probiotics supported by the gut soothing certified organic herb extracts, helps to restore a healthy balance and benefits metabolism in the digestive system. Horses love this formula and the BioEquus helps to maintain a healthy bacterial flora and has a beneficial impact on metabolic processes. BioEquus will assist in maintaining a healthy gut microbiome.



MINERAL TISSUE SALTS

Biochemical tissue salts, or cell salts, are mineral salts that exist in the cells and play a critical role in cellular metabolism.

**Including Calcium, Phosphorus, Zinc, Vitamin D, Folate and Vitamin A.



HI FORM

BreedPlus

MINERAL TISSUE SALTS		
Tricalcium Phosphate	299000	mg / kg
Trimagnesium Phosphate	30000	mg / kg
Monopotassium Phosphate	6000	mg / kg
Potassium Chloride	6000	mg / kg
Sodium Sulphate	6000	mg / kg
Monosodium Phosphate	6000	mg / kg
Calcium Sulphate	2000	mg / kg
Iron Phosphate	7000	mg / kg
Zinc Sulphate	40000	mg / kg
Silica	2000	mg / kg
MAJOR MINERALS		
Calcium	119.0973	g / kg
Phosphorus	72.99854	g / kg
Sodium	4.38405	g / kg
Chloride	2.93256	g / kg
Potassium	14.70535	g / kg
Magnesium	6.505725	g / kg

TRACE MINERALS		
Zinc	14408.943	mg / kg
Copper	3200	mg / kg
Selenium	100	mg / kg
Iodine	246.5	mg / kg
Manganese	0.5508	mg / kg
Iron	1668.5292	mg / kg
Chromium	0.2295	mg / kg
Silica	1996	mg / kg
VITAMINS		
Vitamin A	1070352	IU / kg
Vitamin B1 (Thiamine)	11.892058	mg / kg
Vitamin B2 (Riboflavin)	3.9849	mg / kg
Vitamin B (Niacin)	33.9265	mg / kg
Vitamin B5 (Pantothenic Acid)	1.733	mg / kg
Vitamin B6 (Pyridoxine)	2.295	mg / kg
Vitamin C	14970	mg / kg
Vitamin D	204562	IU / kg
Vitamin E	40229.116	IU / kg
Folic Acid	0.918	mg / kg
Choline	7.975	mg / kg
AMINO ACIDS		
Lysine	11.621414	g / kg



DOSAGE RATES

300kg : 12.5gm (2/3 large scoop)
 500kg : 25.5gm (1.5 large scoop)
 600+ kg : 34gm (2 large scoop)
 Mix well into slightly damp feed.



References

- Wise, M. B., Ordoveza, A. L., & Barrick, E. R. (1963). Influence of variations in dietary calcium: phosphorus ratio on performance and blood constituents of calves. *Journal of Nutrition*, 79, 79-84
- Meyer, H. (1987). Nutrition of the equine athlete.
- West, K. P. (1991). Dietary vitamin A deficiency: effects on growth, infection, and mortality. *Food Nutr Bull*, 13(2), 119-31.
- P. H. Mäenpää, T. Koskinen, E. Koskinen; Serum Profiles of Vitamins A, E and D in Mares and Foals During Different Seasons, *Journal of Animal Science*, Volume 66, Issue 6, 1 June 1988, Pages 1418-1423, <https://doi.org/10.2527/jas1988.6661418x>
- Temporal folate status during lactation in mares and growth in foals. Amy L. Ordakowski-Burk, David S. Kronfeld, Carey A. Williams, Louisa S. Gay, and David J. Sklan. *American Journal of Veterinary Research*, July 2005, Vol. 66, No. 7, Pages 1214-1221
- Ott, E. A., & Asquith, R. L. (1994). Trace mineral supplementation of broodmares. *Journal of Equine Veterinary Science*, 14(2), 93-101.
- Bazzano, A. N., Hofer, R., Thibeau, S., Gillispie, V., Jacobs, M., & Theall, K. P. (2016). A Review of Herbal and Pharmaceutical Galactagogues for Breast-Feeding. *The Ochsner Journal*, 16(4), 511-524.
- Jarde, A., Lewis-Mikhael, A.-M., Moayyedi, P., Stearns, J. C., Collins, S. M., Beyene, J., & McDonald, S. D. (2018). Pregnancy outcomes in women taking probiotics or prebiotics: a systematic review and meta-analysis. *BMC Pregnancy and Childbirth*, 18, 14. <http://doi.org/10.1186/s12884-017-1629-5>
- Penagos Tabares, F., Bedoya Jaramillo, J. V., & Ruiz-Cortés, Z. T. (2014). Pharmacological overview of galactagogues. *Veterinary medicine international*, 2014.
- Piercy, R.J., K. W. Hinchcliff, and S. M. Reed. 2002. Folate deficiency during treatment with orally administered folic acid, sulphadiazine and pyrimethamine in a horse with suspected equine protozoal myeloencephalitis (EPM). *Equine Vet. J.* 34:311-316.
- De Simone, E. A., Bottini, J. M., Alvarez, E. A., & Barbará, A. C. (2013). Association Between Low Serum Zinc Concentration and Hypogammaglobulinemia in Foals of Different Age Categories. *Journal of equine veterinary science*, 33(6), 401-405.