



Disinfectant, Sanitizer, Cleanser
 Bactericidal, Virucidal*, Fungicidal
 Eliminates and Controls Odor
 Controls Bacteria and Algae
 Safe for Poultry, Cattle, Equine, Sheep, Swine,
 Livestock Drinking Water
 Human Drinking Water Disinfectant
 Multi-Purpose Effervescent Tablets
 Kills 99.999% of Bacteria in 60 Seconds

For use in Sanitation, Cleaning and Disinfection in amusement parks, breweries, beverage and food processing, schools, hospitals, nursing homes, hotels, child care centers, daycares, restaurants, spas, hot tubs, salons, veterinary clinics, zoos and aquariums, milk processing facilities, dairy farms, farms, poultry premises, poultry hatcheries, and livestock quarters, office buildings, industrial facilities, homes, camp sites, marine and recreational vehicles, R/V holding tanks, kennels, boarding facilities, laboratories, lab animal facilities, licensed care facilities.

Disinfects precleaned, hard, nonporous surfaces in 5 minutes. Effective against *Pseudomonas aeruginosa*, *Staphylococcus aureus*, *Salmonella enterica* (formerly *choleraesuis*), *Feline Calicivirus as a Surrogate Virus for Norovirus, *Canine parvovirus, *Respiratory syncytial virus, *2009-H1N1 Influenza A virus in 5 minutes.

Sanitizes precleaned, hard, non-porous surfaces in 1 minute. No Rinse Sanitizer. At 200 ppm available chlorine, this product is an effective sanitizer against *Campylobacter jejuni*, *Vibrio cholerae*, *Escherichia coli O157:H7*, *Klebsiella pneumoniae*, *Listeria monocytogenes*, *Yersinia enterocolitica*, *Shigella sonnei*, *Salmonella typhi* and *Staphylococcus aureus* in 1 minute. At 1300 ppm available chlorine, this product is effective against *Trichophyton mentagrophytes* (common cause of Athlete's Foot) in 5 minutes.

ACTIVE INGREDIENT:

Sodium Dichloro-s-Triazinetrione50%
 OTHER INGREDIENTS50%
 Available Chlorine...31.75% Total.....100%

Keep Out of Reach of Children
DANGER

Manufactured for Activon, Inc.
 123 Commercial Drive, Beaver Dam, WI 53916
 800.841.0410

EPA Reg. No. 66570-2
 EPA Est. No. 087268-WI-001

PRECAUTIONARY STATEMENTS - Hazards to Humans & Domestic Animals

DANGER: Corrosive: Causes irreversible eye damage. Harmful if swallowed, inhaled or absorbed through skin. Do not get in eyes, on skin or on clothing. Avoid breathing dust. Wear goggles or face shield. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash separately before reuse.

FIRST AID: Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also call 800.222.1222 for emergency medical treatment advice.

If in Eyes : Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If Swallowed: Immediately call a poison control center or doctor for treatment advice. Have person drink large amounts of water if able to swallow. Avoid alcohol. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If Inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

If on Skin or Clothing: Take off contaminated clothing. Immediately rinse skin with plenty of water for 15-20 minutes. Call a poison control center or doctor for further treatment advice.

Note to Physician: Probable mucosal damage may contraindicate the use of gastric lavage.

Physical or Chemical Hazards

Use only clean, dry utensils. Mix only into water. Contamination with moisture, dirt, organic matter or other chemicals (including pool chemicals) or any other foreign matter may start a chemical reaction with generation of heat, liberation of hazardous gasses and possible generation of fire and explosion. Avoid any contact with flaming or burning materials such as a lighted cigarette. Do not use this product in any chlorinating device that has been used with any inorganic or unstabilized chlorinating compounds (e.g., calcium hypochlorite). Such use may cause fire or explosion.

Environmental Hazards

This pesticide is toxic to fish and aquatic organisms. Disinfects floors, walls and other hard nonporous surfaces including walls, floors, tables,

chairs, countertops, bathroom fixtures, sinks, shelves, racks, carts, refrigerators, coolers, glazed tile, linoleum, vinyl, glazed porcelain, plastic (such as polypropylene and polyethylene), stainless steel, or glass. This product is designed for use in amusement parks, breweries, beverage and food processing plants, schools, hospitals, nursing homes, hotels, child care centers, restaurants, spas, hot tubs, veterinary clinics, zoos, milk processing facilities, dairy farms, farms, poultry premises, poultry hatcheries, and livestock quarters, office buildings, industrial facilities, homes, camp sites, marine and recreational vehicles, kennels, boarding facilities, laboratories, lab animal facilities.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

DISINFECTION

Prepare a 1,300 ppm solution (refer to dilution chart). Clean surface then apply solution with mop, cloth, sponge, brush, or coarse trigger sprayer. Allow surface to remain wet for 5 minutes. Allow to air dry. All treated equipment that will contact food, feed, or drinking water must be rinsed with potable water before reuse.

This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the blood stream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to pre-clean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection.

FOR USE THROUGHOUT FOOD AND BEVERAGE PROCESSING AND FOOD HANDLING OPERATIONS

This product is recommended for sanitizing all types of hard, non-porous equipment and utensils used in food processing and canning plants, bottling plants, breweries, fish processing plants, meat and poultry processing plants, milk handling and processing plants, restaurant and institutional dining establishments. Use a 100 ppm available chlorine solution (refer to Dilution Chart) to sanitize previously cleaned processing and packaging equipment. Allow at least a one minute contact time before draining. Allow adequate draining before contact with beverages. To control the growth of bacteria in brewery pasteurizers, clean badly fouled systems before treatment. When the system is noticeably fouled, add 3 ppm available chlorine (refer to Dilution Chart) to system water. Repeat this dosage if necessary until the free available chlorine (FAC) level is 0.5 -1.0 ppm, as determined by use of a reliable test kit. To maintain a FAC of 0.5 – 1.0 ppm, add 0.5 ppm – 1 ppm available chlorine (refer to chart) daily as needed. Add this product to the system at a point where adequate flow is maintained.

EGG PROCESSING PLANTS

Clean and destain egg shells prior to sanitizing. To clean egg shells, spray with a 90°F to 120°F 100 ppm available chlorine solution. Spray-rinse the cleaned eggs with warm potable water. Only clean, whole eggs may be sanitized. Dirty, cracked or punctured eggs may not be sanitized. To destain egg shells, immerse the eggs in a 90°F to 120°F solution containing 100 ppm available chlorine. After destaining, the eggs must be cleaned by spraying with an acceptable cleaner. Follow with a potable water rinse. To sanitize clean shell eggs intended for food or food products, spray with a solution containing 100 ppm available chlorine. The solution must be equal to or warmer than the eggs, but not to exceed 130°F. Wet eggs thoroughly and allow to drain. Eggs that have been sanitized with this chlorine compound may be broken for use in the manufacture of egg products without a prior potable water rinse. Eggs must be reasonably dry before casing or breaking. The solution must **not** be reused for sanitizing eggs. Thoroughly clean and sanitize all egg cups, breaking knives, trays and other equipment that come into contact with "off" eggs. First, clean all equipment. Before placing back in use, spray with a solution containing 100 ppm available chlorine. Allow surfaces to completely drain before contact with egg product. To sanitize egg freezers and dryers (tanks, pipelines and pumps), use the spray method of treatment (see Sanitizing Application Methods section). This procedure is generally used to sanitize large, non-porous surfaces that have already been cleaned of physical soil. Prepare a solution containing 100 ppm available chlorine. Heavily apply spray to all surfaces the eggs will touch. Thoroughly spray all treated surfaces, corners and turns. Allow at least a one minute contact time before draining. Allow equipment to drain adequately before contact with eggs.

SANITIZING HARD, NON-POROUS SURFACES, DISHES, GLASSES, FOOD PROCESSING EQUIPMENT AND UTENSILS, DAIRY AND BREWERY EQUIPMENT AND UTENSILS

This product is an effective sanitizing agent. Treatment with this product throughout food and beverage processing and food handling operations can help ensure the quality and safety of the final product.

Hand Washing of Items

1. Remove all gross food particles and soil by a preflush or prescrape and, when necessary, presoak treatment. Wash surfaces or objects with a good detergent or compatible cleaner, followed by a potable water rinse before application of the sanitizing solution. 2. Prepare a 100 ppm available chlorine sanitizing solution. 3. Place equipment, utensils, dishes, glasses, etc. in the solution or apply the use solution to surfaces using a cloth, sponge, or coarse sprayer. 4. Allow to stand at least one minute, drain the excess solution from the surface and allow to air dry. 5. Fresh sanitizing solution must be prepared at least daily or more often if the solution becomes diluted or soiled.

Machine Washing of Items

1. Remove all gross food particles and soil by a preflush or prescrape and, when necessary, presoak treatment. Wash surfaces or objects with a good detergent or compatible cleaner, followed by a potable water rinse before application of the sanitizing solution. 2. Prepare a 100 ppm available chlorine solution. 3. Add the solution to the feed tank of immersion or spray type machines that can provide at least one minute contact time for sanitizing dishes, glasses, food processing equipment or utensils. Allow to drain and air dry before use. 4. Promptly use the sanitizing solution. Prepared solutions cannot be reused for sanitizing but may be used for other purposes, such as cleaning.

ANIMAL HOUSING FACILITIES

(Including Poultry Houses, Swine Confinement Facilities, Veterinary Clinics, Zoos and Farms) The problem of odor control in poultry houses and other animal facilities is not completely solved by normal cleaning practices. The regular use of an efficient bactericide and deodorant is strongly recommended and often required by health authorities. Remove all poultry or animals and feeds from premises, trucks, vehicles, coops, crates and enclosures. Remove all litter and manure or droppings from floors, walls and surfaces of barns, pens, stalls, chutes and other facilities and fixtures occupied or traversed by animals or poultry. Empty all troughs, racks and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. To disinfect, saturate all surfaces with a 1,300 ppm available chlorine solution for a period of five minutes. Immerse all halters, ropes, and other types of equipment used in handling and restraining animals, as well as forks, shovels, and scrapers used for removing litter and manure. Ventilate buildings, cars, boats, coops and other closed spaces. Do not house livestock or poultry or employ equipment until treatment has been absorbed, set or dried. Thoroughly scrub all treated feed racks, mangers, troughs, automatic feeders, fountains and waterers with soap or detergent and rinse with potable water before reuse.

SHOE AND BOOT BATH SANITIZER (Not Applicable in California)

Shoe and Boot baths containing one inch of freshly made 100 ppm available chlorine should be placed at all entrances to buildings, hatcheries and at all the entrances to the production and packaging rooms. Scrape waterproof shoes and boots and place into the bath at least daily or sooner if solution appears diluted or dirty.

MILK HANDLING AND PROCESSING EQUIPMENT

This product can be used on dairy farms and in plants processing milk, cream, ice cream and cheese. Rinse milking machines, utensils and all equipment with cold water to remove excess milk. Clean and rinse prior to sanitizing. To sanitize, spray or rinse all precleaned surfaces with 100 ppm available chlorine solution. Allow adequate draining before contact with dairy products. It is important to clean out large deposits of milk or other organic matter before sanitizing. A sharp decline in the available chlorine content of the sanitizer following circulation through milk processing equipment is usually regarded as evidence of inadequate cleaning of the equipment and should be promptly investigated.

SANITIZING APPLICATION METHODS

Freshly prepare all sanitizing solutions. Test solutions during use to ensure the concentration does not drop below the recommended level. Keep in properly labeled containers to protect against contamination. Discard unused solutions.

Pressure Method of Sanitizing Equipment

This method is commonly used to sanitize closed systems, such as fluid milk cooling and handling equipment. It is also appropriate for sanitizing weigh tanks, coolers, short-time pasteurizers, pumps, homogenizers, fillers, sanitary piping and fittings, and bottle and can fillers. For mechanical operations, prepared solutions cannot be reused for sanitizing but may be used for other purposes, such as cleaning. For manual operations, fresh sanitizing solutions must be prepared at least daily or more often if the solution becomes diluted or soiled. First, disassemble and thoroughly clean all equipment immediately after use. Remove all gross food particles and soil by a preflush or prescrape and, when necessary, presoak treatment. Wash surfaces or objects with a good detergent or compatible cleaner, followed by a potable water rinse before application of the sanitizing solution. Then place back in operating position. Prepare a solution containing 100 ppm available chlorine in a volume equal to 110% of capacity. Pump the solution through the system until it is filled with sanitizer and air excluded. Close final drain valves and hold under pressure for one minute to ensure proper contact with all surfaces. Remove a portion of the cleaning solution from the drainvalve and test with a chlorine test kit. Repeat entire cleaning or sanitizing process if effluent contains less than 50 ppm available chlorine.

Spray Method of Sanitizing Equipment

The spray method is generally used to sanitize large, non-porous surfaces that have already been freed of physical soil. It is appropriate for batch pasteurizers, holding tanks, weigh tanks, tank trucks and cars, vats, tile walls, ceilings and floors. Clean all surfaces after use. Prepare a solution containing 100 ppm available chlorine. Use pressure spraying equipment designed to resist chlorine-containing solutions (e.g. rubber-coated, plastic or stainless steel). When using any other kind of spraying equipment, always empty and thoroughly rinse the spray equipment with potable

water immediately after treatment. Apply spray heavily to all surfaces the product will touch. Thoroughly spray all treated surfaces, corners and turns until wet. Allow at least a one minute contact time before draining. Allow excess solution to drain and air dry then place in service. Vacate area for at least two hours.

General Rinse Method

Solutions containing 100 ppm available chlorine sanitize plant floors, walls and ceilings, and also control odors in refrigerated areas and drain platforms. Generously flush or swab surfaces with the solution. After one minute contact time allow solution to drain and then air dry.

DISINFECTION OF DRINKING WATER

DRINKING WATER: Add 10 ppm available chlorine (refer to Dilution Chart) to water to be disinfected. Let the water stand seven to fifteen minutes before using. A residual of 0.2 ppm available chlorine, as measured by a reliable test kit, should be maintained in the water to insure disinfection.

EMERGENCY DRINKING WATER: This product may be used to disinfect or pre-treated (settled, coagulated and/or filtered) water supplies intended for use as drinking water for humans and domestic animals on an emergency basis as defined in 40 CFR, Part 165-179. The source of the water to be treated may be a river, lake, well, cistern or similar system. To obtain the desired disinfection results, the water to be treated should be clear and free dirt and organic debris. If the source of the water is cloudy and contains dirt and organic debris, the water should be held in holding tanks or ponds, treated with coagulating agents and filtered to remove the dirt and organic debris. Preparation of Stock Solution – Dissolve proper amount of this product to produce a 6,000 ppm available chlorine (refer to Dilution Chart) stock chlorine solution. Add 20 drops of this stock solution for each liter of water to be treated. The stock solution should be prepared fresh weekly.

EMERGENCY DISINFECTION AFTER FLOODS

DISTRIBUTION SYSTEM: Flush repaired or replaced section with water. Establish a chlorinating station and apply sufficient product until a consistent available chlorine residual of at least 10 ppm available chlorine (as measured by chlorine test kit) remains after a 24 hour retention time.

EMERGENCY DISINFECTION AFTER DROUGHT

WATER SHIPPED IN BY TANKS, TANK CARS, ETC.: Thoroughly clean all containers and equipment. Spray a 500 ppm available chlorine solution and rinse with potable water after 5 minutes. During the filling of the containers, dose with sufficient amounts of this product to provide at least a 0.2 ppm chlorine residual, as measure by a chlorine test kit.

POULTRY, SWINE, CATTLE, LIVESTOCK DRINKING WATER DISINFECTION

This product can be used as a disinfectant for the drinking water of poultry, swine, cattle and other livestock. If the water supply is badly fouled, then add 5 ppm available chlorine to the water supply. After 24 hours the addition rate can be reduced to 1 ppm available chlorine. If the microbiological content of the water is eliminated, the concentration of available chlorine can be reduced to 0.5 ppm. If the microbiological control is not adequate at 1 ppm available chlorine, then add 1.5 ppm available chlorine to the livestock drinking water. This product should be administered continuously into the drinking water from the time of placement (day one). Remove this product from the drinking water 24 hours prior to vaccination and re-administer 24 hours after vaccination.

FILTRATION DEVICES

This product is for use in filtration devices (water purification systems and its cartridges). Its purpose is to clean membranes, such as reverse osmosis membranes of fouling contaminants. One [1] dose of product is necessary to achieve reduction in fouling contaminants. Product should be used following the manufacturer's instructions.

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal. **Storage:** Keep product dry in tightly closed container when not in use. Store in a cool, dry, well-ventilated area away from heat or open flame. **Pesticide Disposal:** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your state pesticide or environmental control agency, or the hazardous waste representative at the nearest EPA Regional Office for guidance.

Container Disposal: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Thoroughly rinse empty container with water to dissolve all material before discarding in trash or recycling.

Emergency Handling: In case of contamination or decomposition do not reseal container. If possible, isolate container in open and well-ventilated area. Flood with large volumes of water. Dispose of contaminated material in an approved landfill area.

Dilution Chart {4 g Tablets}			Dilution Chart {0.33g tablets }			Dilution Chart {17.68g tablets}		
Solution (Available chlorine)	Tablets (4g or 0.14 oz. each)	Gallons of Water	Solution (Available chlorine)	Tablets (0.33g or 0.01oz each)	Gallons of Water	Solution (Available chlorine)	Tablets (17.68g or 0.62oz each)	Gallons of Water
0.5ppm	1	625	0.5ppm	1	55	0.5ppm		
	4	2,500		10	550		1	2,500
	8	5,000					2	5,000
	15	10,000				4	10,000	
1ppm	1	325	1ppm	1	25	1ppm		
	2	650		5	135			
	3	1,000		10	275		4	6,000
	30	10,000					7	10,000
3 ppm	1	100	3 ppm	11	100	3 ppm		
	5	500		25	230		1	500
	9	1,000		55	500		2	1,000
	90	10,000				20	10,000	
4 ppm	6	500	4 ppm	73	500	4 ppm	3	1,000
10 ppm	15	500	10 ppm	2	5	10 ppm	7	1,000
100 ppm	1	3	100 ppm	1	1 quart	100 ppm	1	15
	3	10		4	1		3	40
	12	40						
1,300ppm	4	1	1,300ppm	12	1 quart	1,300ppm	3	3
	39	10		47	1		11	12