

#### **EARLY SPRING:**

Apply Green-Up plus Crabgrass Preventer and MAG-I-CAL on the same day.

**LATE SPRING:** Apply Green-Up Weed & Feed.

#### **SUMMER:**

Apply Love Your Soil to feed the soil and stimulate its biology.

### Apply Winter Survival to feed the lawn.

Ask for a FREE copy of our

New American Lawn Guide or go to

www.newamericanlawn.com

A WIDE SPECTRUM OF BROADLEAF WEEDS WHILE PROVIDING BOTH FAST AND LONG FEEDING ACTION. WEEDS CONTACTED BY THIS PRODUCT GRADUALLY DIE AND DISAPPEAR FROM THE LAWN.

THIS PRODUCT IS A PREMIUM GRADE HERBICIDE AND LAWN FOOD COMBINATION WHICH CONTROLS

#### WHEN TO APPLY GREEN-UP™ WEED & FEED

**OPTIONAL APPLICATION:** 

If insects are a problem, apply Summer

Survival Lawn Food & Insect Control

anytime from June through August.

Best results are obtained when weeds are growing actively in Spring or early Fall. Avoid contact with desirable shrubs, plants, vegetables, or flowers; especially when applying with a rotary (spinner) type spreader. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. For optimum results (1) Avoid mowing 1 to 2 days before and after a polication; (2) Apply when grass is moist (after a rain, sprinkling or dew); (3) Avoid unnecessary disturbances, including watering of treated areas for 48 hours after application. Watering will wash off weed killing material; and (4) Avoid applying if rainfall is expected within 48 hours following treatment. Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse law moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur. NOTE: When applying to Southern Grasses - under conditions of inadequate moisture, high temperature and high humidity,

over-application may cause minor transitory discoloration on Centipede and St. Augustine grasses. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

#### HOW TO APPLY GREEN-UP™ WEED & FEED

This 15 pound bag should be used on 5,000 square feet or apply at 3 pounds per 1,000 square feet. Apply evenly with a lawn spreader as a one pass application. DO NOT HAND SPREAD.

## SPREADER SETTINGS (One pass application)

**BROADCAST (ROTARY) SPREADERS** The following settings are approximate for applying contents of this package at a suggested rate of 3 pounds per 1,000 square feet. Settings are based on an average walking

speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 square feet and adjust to apply at the rate of .75 pounds per 250 square feet. Sweep (do not wash) all fertilizer materials from surfaces immediately after application, because certain fertilizer ingredients can cause permanent stains Wash out spreader after each use and allow it to dry thoroughly. Treat any exposed metal spreader areas with oil to reduce corrosion.

Total Nitrogen (N) .......21% Derived F

IF INHALED • Move person to fresh air.

Sign up to receive timely lawn tips

www.jonathangreen.com

21.00% Urea Nitrogen\*

FIRST AID

# **GUARANTEED ANALYSIS: 21-0-3**

Fig. 15 on SKIN • Take off contaminated clothing.
• Rinse skin immediately with plenty of water for 15 to 20 minutes **CLOTHING** • Call a poison control center or doctor for treatment advice.

IF IN EYES • Hold eye open and rinse slowly and gently with water for 15

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840

My Pledge Of Satisfaction

We have made every effort in the formulation of this product to insure that you will be satisfied with the results. 16, for any reason, you are not satisfied, please write and describe the problem. Our lown specialists will make every effort to assure your success.

tion regarding the contents and levels of metals in this product is available on the internet at http://www.aapfoa.org/metals.htm

If person is not breathing, call 911 or an ambulance, then give artificial respiration, preterably mouth to mouth, if possible.
 Call a poison control center or doctor for further treatment advice.

 Remove contact lenses, if present, after the first 5 minutes, then Call a poison control center or doctor for treatment advice.

Urea. Sulfate of Potash.

Polymer Coated Urea

#### Jonathan Green New American Lawn New American Lawn Deluxe & Pro 14 New American Lawn Hand Held AgriFab Chapin Cyclone

EZ Republic (Ortho) Earthway (Ames) Precision (Red Devil) Scotts Rotary Spreaders Scotts EasyGreen ENVIRONMENTAL HAZARDS This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Drift and runoff may be hazardous to aquatic organisms in waters adjacent to treated areas. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the

CAUTION

13

14

3-1/2

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**USE PRECAUTIONS FOR TURF, ORNAMENTAL** 

#### DROP SPREADERS 1-1/2 Agri-Fab Central EZ Republic (Ortho) Earthway (Ames) 12 Precision (Red Devil) Scotts Turf Builder Classic, AccuGreen 1000-2000-3000,

THE NEW AMERICAN LAWN **FEED YOUR LAWN** ...AND YOUR SOIL!

Alder Artichoke

Aster, Wild Bedstraw

Beggartick Betony, Florida

Bindweed, Field Bindweed, Hedge

Bitter Wintercress Bittercress, Hairy

Bristly Oxtongue

Buckharn Buckwheat, Wild

Bullthistle, Wavyleaf

Buttercup, Bulbous

Carrot, Wild Catchweed, Bedstraw

Chamamile, Corn Chickweed Chickweed, Cammon

Chickweed, Mousean

Cinquefoil, Rough

Clematis, Western

Yellow Sweet

Clover, Strawberry

Clover, Annual

Clover, Crimson Clover, Hop

Clover, Red

Catsear, Spotted

Blackeyed Susan Bloodflower, Milkweed

Bedstraw, Pineywoods

Aster

Biden

Bindweed

Bitter weed

Box Elder

Burweed

Carpetweed

Catnip Catsear

Chicary Cinquefoil

DIRECTIONS FOR USE

Clover, White Cockle

Cacklebur, Spiny Caffeebean

Careopsis, Plain Creeping Jenny

Croton, Wooly

Cudweed Curly Indigo

Daisy, English Daisy, Oxeye

Dandelian, Fall Dandelian, False

Dock, Curly

Dogbane Dogfennel

Elderberry

Dack, Smooth

Fiddle neck Fieldcress , Austrian

Filaree, Redstem

Flea bane, Daisy

Fleabane, Hair

Fleabane, Rough

Florida Pusley FourO'-Clock, Wild Frenchweed

Galinsoga Galinsoga, Smallflowe Garlic, Wild

Geranium, Wild Goathead Goatsbeard

Geranium, Carolina

Hawkweed, Orange

Heartleaf Drymary Heathaster

Hemlock, Poison

Goldenrod

Groundse

Gumweed

Healall

Hawkweed

Flixweed

Dandelion

Dock Dack, Broadleaf

Cocklebur

Caffeeweed

Croton

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS. A minimum of 30 days between applications is

required. JONATHAN GREEN WEED AND FEED is a stable mixture of three herbicides. It is for use on all residential turf. Do not

apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled. This product may be used on Bahia, common Bermuda, Bluegrass,

This product is designed to promote lush green growth in lawns and control many broadleaf weeds such as:

Onion, Wild

Pansy, Field

Parsley-piert Parsnip Parsnip, Wild

Pearlwart Pennycress

Pennywort

Peppergrass

Pepperweed

Pepperweed, Virginia

Pigweed, Redroat Pigweed, Russian Pigweed, Smooth Pigweed,

Scarlet Pimpernel Plantain

Plantain, Blackseed

Plantain, Bracted

Plantain, Broadlea Plantain, Buckhorn

Plantain, Hoary Plantain, Narrowleaf Plantain, Slender

Plantain, Wooly Pakeweed

Poorjoe

Prickly Sida

Puncture Vine

Purslane Radish, Wild

Salsify, Western Shepherdspurse

Smartweed

Somel, Red

Sowthistle

Speedwell Spiny Amaranth

Somel, Sheep

Sowthistle, Common

Spanishneedle Spatterdock

Ragweed Rape, Wild Rose, Southern Wild

Primrose

Rush

Pigweed Pigweed, Prostrate

Oxalis

Spurge, Prostrate Spurge, Spotted Spurweed St. Johnswort Stinkweed

Strawberry, Wild

Sunflower, False Sweet Potato, Wild Tansy Ragwart

Thistle, Canada Thistle, Musk Thistle, Russian

Thistle, Scotch

Trefail, Tick

Tumbleweed Velvet Leaf

Vervain, Blue

Vetch

Vetch, Bird Vetch, Milk

Vervain, Hoary

Vervain, Prostrat Vervain, Tall

Vetch, Narrowleaf Vetch, Wild

Virginia Creeper Willow Witchweed

Woodsorrel

Womseed

Yellow Rocket

Yellowflower

Pepperweed

For best results, do not apply on a windy day.

· Swath will vary due to weather conditions,

To determine how much coverage you need multiply the length of your lawn by its width

This equals your total square footage. Deduct any non-lawn areas (house, gardens,

walkways, etc.). One bag covers 5,000 sq. ft.

If your spreader is not listed here,

or if you bought a Jonathan Green

spreader prior to 2017, please visit

www.jonathangreen.com

for the correct setting.

product used and walking speed.

Apply products only to lawn.

(RV011216)

Virginia Buttanweed

Toadflax Trailing Crownvetch

Stitchwort

Sumac Sunflower

Tarweed

Thistle Thiste, Blessed Thiste, Bull

Centipede, Fescue, Ryegrass, St. Augustine, and Zoysia grass. DO NOT USE on Dichondra and Carpet grass. DO NOT USE this product where

desirable clovers are present or allow the product to "wind drift" onto flowers, vegetables, ornamentals, shrubs, trees or other desirable plants.

Hemp Henbit

Hoary Cress

Honeysuckle Horsenettle

Horsetail

Jewelweed

Knawel

Kochia

Lespedeza

Locaweed

Lettuce, Blue Lettuce, Prickly Lettuce, Wild

Mallow, Indiana

Marigold, Wild Marshelder

Matchweed

Medic, Black

Morningglory

Mustard Mustard, Black Mustard, Garlic

Mustard, Hedge

Mustard, Tansy Mustard, Tumble

Mustard, White

Nettle, Burning Nettle, Dead Nettle, Stinging

Nettle, Bull

Nettle, Tall

Oak, Poison

Nutgrass

Morningglary, Wooly Mugwort Mullein, Common

Mallow, Venice Marigold, Roundleafed

Knapweed, Spotted

Knotweed Knotweed, Prostrate

#### USAGE TIPS For rectangular lawns

Apply two header strips across each end for a turning area. Then apply back and forth in the longest direction. For irregular-shaped lawns Apply on header strip around the entire lawn for a turning area, then apply back and forth.

Note: Each pass should be within 5 feet of each other to ensure an overlap. • Do not leave spreader open when making turns.

## STORAGE AND DISPOSAL

STORAGE: Store in original container and keep any bags containing unused product in locked storage area inaccessible to children and domestic ar feed or foodstuffs. To prevent cross-contamination, do not store near open containers of fertilizer, seed or other pesticides CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. If empty: Place in trash or affer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor (induding toilet) or outdoor

#### WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY

# PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTIORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL

OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENAITIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL, TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR RISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER. OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT. f you do not agree with or do not accept any of directions for use, the warranty disdaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

22344

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular tertilizer program by providing .63 pounds of nitrogen per 1,000 square feet and herbicide to control listed weeds. Please consult your local extension office for questions regarding a regular fertilizer program. Distributed & Guaranteed by: Jonathan Green, Inc. POB 326 Farmingdale, NJ 07727

Postemergence: Limited to 2 applications per year. Maximum of 250 pounds of product/acre per application. The maximum seasonal rate is 500 pounds of product/acre, excluding spot treatments.

treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn will help to prevent run-off to water bodies or drainage systems.

Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not

apply within 25 feet of rivers, fishponds, lakes, streams, reservoirs, marshes, estuaries, both of apply within 25 feet of rivers, fishponds, lakes, streams, reservoirs, marshes, estuaries, bosy, and oceans. Do not apply when windy. Apply this product directly to your lawn and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent product run-off, do not over water the treated area to the point of run-off or apply when raining or when rain is expected that day.

Harmful if absorbed through skin or inhaled. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Avoid breathing dust. Wear long-sleeved shirt, long pants, shoes, socks, and chemical-resistant gloves made of any waterproof material. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

(golf courses, cemeteries, parks, sports fields, turfgrass, lawns and other grass areas)

000'9





For Medical Emergencies Only, Call (877) 325-1840 (800) 454-9300 For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC PRECAUTIONARY STATEMENTS SEE BACK PANEL FOR FIRST AID AND ADDITIONAL **MOITUAD** KEEP OUT OF REACH OF CHILDREN

Code on bag indicates manufacturing sile. EPA EST. NO. 0755227-PA-12, 8378-IN-1, 9198-0H-1M, 9198-0H-28, 9198-M-00-L ENV REG, NO. 228-280-50931 #CONTAINS THE SINGLE ISOMER FORMS OF MECOPROPy and DICHLORPROPy dl\ab2100.0 ;%2f.0 . \*\*\* (+) R-2 {2,4-DicHomphenoxy)propionic Acid dl\so8200.0 ;%82.0 ... ISOMER SPECIFIC METHOD, EQUIVALENT TO:

DISSOLVE® is a Registered Trademark of Nufarm Americas Inc. Patent #5,266,553 %000'00L " OTHER INGREDIENTS: 88.939% Dine Solin of (+) R.2-(2,4-Dich lorophenoxy) propionic Acid Acid (xonahenoxy) %6ZE0 . ‡\*\* bis A sin oiqorq (yxonə rlqorol rl D-l-l yrlt 9 M-S )- S-A-(+) to tl n2 ənim nlyrl təmi (I %Z81.0 ACTIVE INGREDIENTS:

NET WEIGHT: 15 LBS. (6.80 Kg) 21-0-12



**MEEDSI** BROADLEAF CONTROLS

**OR FALL** IN SPRING **FOR USE** 

See Back Panel for Details THE NEW AMERICAN LAWN

REWEMBER TO "FEED YOUR SOIL"

THE ROOT **DOMN 10** KITTS MEEDS

**SHINOM E** FOR UP TO **LEEDS LAWNS** 



QUALITY LAWNS SINCE 1881

Jonathan Green





Easy-to-use! Resealable bag!

cut or Tear here to open



COVERS 5,000 sq. FT.

Jonathan Green