

Instructions for Applying the Dalric Extension #3

State of the Art Treatment for Angular Limb Deformities in Young Foals

FOOT PREPARATION

- Trim and balance the foot.
- Round the ground surface of the hoof to accommodate the seating of the foot in the cuff.
- Roughen the wall with the smooth side of the rasp. This provides an adequate glue contact surface.
- Clean dirt from all hoof cracks and cover potential abscesses with a small piece of tape or silicone. This prevents sealing bacteria beneath the glue.

CUFF & EXTENSION PREPARATION

- Apply the aluminum extension lateral (outside) for all toe-in (varus) deformities or medial (inside) for toe-out (valgus) deformities.
- Clamp a pair of wide-mouth vice grips across the heels of the cuff and aluminum extension. This will provide proper alignment of the cuff and aluminum extension when drilling the holes.
- Use a 5/32" drill bit to drill holes through the aluminum extension into the cuff. Be careful not to get the bit too close to the felt lining of the cuff as it will grab.
- Place the aluminum rivets through the cuff and into the aluminum extension.

SHOE APPLICATION

- If you are working with an animal that is fractious, you may have your vet administer a sedative.
- Clean the hoof wall with denatured alcohol and ensure the foot is dry.
- Remove the SuperFast[™] cartridge and cut the tip off with a pair of nippers. Squeeze a small amout of adhesive from the cartridge to ensure both sides are dispensing equally.
- Apply a mixing tip to the end of the cartridge and twist clockwise to lock it in place.
- Apply SuperFast[™] to the felt side of the shoe only.
- Apply the shoe snuggly to the clean foot.
- SuperFast[™] sets in 30 seconds, so remember to work quickly. Total cure time is 2-3 minutes. If you need a longer set time, SuperFast[™] can be stored in a refrigerator prior to use.
- To apply a second shoe, simply remove the tip, squeeze a bit of SuperFast[™] from the cartridge and apply a new tip.
- When finished, apply a cap to the SuperFast[™]. The remaining adhesive can be used at another time.

REMOVAL OF THE SHOE

• Use nippers or a rasp to separate the entire cuff from the horn wall and then pull the shoe.

RECOMMENDED SHOEING SCHEDULE

 For best results, follow this shoeing schedule as the foal's feet continue to grow. Exceeding the recommended shoeing time can alter natural foot development.

Age	Maximum Time for Shoes To Be Worn	Days Following Shoe Removal To Re-evaluate
Birth - 2 Weeks	4 Days	7 Days
2-4 Weeks	7-10 Days	7-10 Days
4-8 Weeks	10-14 Days	7-10 Days
8-12 Weeks	14 Days	7-10 Days
3 Months	21 Days	7-10 Days
8 Months +	28 Days	14 Days

PREVENTION OF SHOE LOSS

- Make sure the shoe fits snuggly and the foot is clean prior to application. Keep the foal still during the curing time.
- Do not turn out into mud.
- Do not turn out into large, open fields. Limit exercise to 1-2 acres.
- Use caution when turning out with other horses, as they will step on the extension.

SORENESS FOLLOWING APPLICATION

- Fitting the shoe too tightly will pinch the quarters resulting in slight soreness. If this occurs, pull the shoe and reapply after the foal becomes sound.
- Sealing bacteria in open areas along the white line can cause acute lameness. If this occurs, remove the shoe immediately and notify your veterinarian. The abscess will need to be opened at the white line and cared for appropriately.

