

TOP SHELF SUPER T

For Use by Licensed Professionals Only

This pre-deadener has one large advantage: the addition of benzocaine for faster onset. It contains three anesthetics: 5% Lidocaine, 1% Tetracaine and 12% Benzocaine. It is safe for use on all procedures *except eyeliner*. Care should be taken to avoid getting it in the eyes and mouth.

FOR EXTERNAL USE ONLY ON INTACT SKIN: Check with client for history of allergic reactions to topical anesthetics before applying. Keep in a cool dark place, *out of the reach of children*.

WARNING: Avoid direct contact with the eyeball. If contact should occur, irrigate with copious amounts of sterile saline, and contact an eye care physician. **DO NOT SWALLOW.** If accidentally swallowed, seek medical help immediately.

SUGGESTED APPLICATION TIMES	EYEBROWS	15- 20 minutes
	For best results, lightly exfoliate prior to application. Gently massage into skin and then apply a medium coating over the top. Cover with a plastic (occlusive) dressing. Wipe away immediately prior to tattooing with Superior Super Juice or Five-Star Vasocaine .	
	LIPS	15- 20 minutes
	For best results, lightly exfoliate prior to application. Gently massage into skin and then apply a medium coating over the top. Cover with a plastic (occlusive) dressing. Wipe away immediately prior to tattooing. <i>Do not use inside of the mouth.</i> Use Superior Super Juice during procedure to continue anesthesia, or New and Improved Blue Gel , Five-Star Vasocaine or Maximum Zone 2 to continue anesthesia and constrict blood vessels.	
	BODY SKIN	90-120 minutes
Prior to tattooing or piercing: lightly exfoliate. Gently massage into skin and then apply a medium coating over the top. To prevent absorption into clothing and retain effective anesthesia, cover with an occlusive dressing. Thicker skin requires longer time to absorb the cream than thinner skin. Use Superior Super Juice during procedure to continue anesthesia, or New and Improved Blue Gel , Five-Star Vasocaine or Maximum Zone 2 to continue anesthesia and constrict blood vessels.		
PIERCING	15- 20 minutes	
Apply to eyebrows, navels and nipples. Avoid genital areas. Thicker skin requires longer pre-treating time. Lightly exfoliate. Gently massage into skin, then apply a medium coating on top. Cover with occlusive dressing. Wipe away immediately prior to piercing. Use New and Improved Blue Gel or Maximum Zone 2 during jewelry insertion to continue anesthesia and reduce swelling.		

DERMAL SOURCE
 2319 NE Glisan Street, Portland, Oregon 97232, U.S.A.
 Phone: +1 866-568-3223, Fax: 503-595-0293
www.dermalsource.com



TOP SHELF SUPER T

For Use by Licensed Professionals Only

This pre-deadener has one large advantage: the addition of benzocaine for faster onset. It contains three anesthetics: 5% Lidocaine, 1% Tetracaine and 12% Benzocaine. It is safe for use on all procedures *except eyeliner*. Care should be taken to avoid getting it in the eyes and mouth.

FOR EXTERNAL USE ONLY ON INTACT SKIN: Check with client for history of allergic reactions to topical anesthetics before applying. Keep in a cool dark place, *out of the reach of children*.

WARNING: Avoid direct contact with the eyeball. If contact should occur, irrigate with copious amounts of sterile saline, and contact an eye care physician. **DO NOT SWALLOW.** If accidentally swallowed, seek medical help immediately.

SUGGESTED APPLICATION TIMES	EYEBROWS	15- 20 minutes
	For best results, lightly exfoliate prior to application. Gently massage into skin and then apply a medium coating over the top. Cover with a plastic (occlusive) dressing. Wipe away immediately prior to tattooing with Superior Super Juice or Five-Star Vasocaine .	
	LIPS	15- 20 minutes
	For best results, lightly exfoliate prior to application. Gently massage into skin and then apply a medium coating over the top. Cover with a plastic (occlusive) dressing. Wipe away immediately prior to tattooing. <i>Do not use inside of the mouth.</i> Use Superior Super Juice during procedure to continue anesthesia, or New and Improved Blue Gel , Five-Star Vasocaine or Maximum Zone 2 to continue anesthesia and constrict blood vessels.	
	BODY SKIN	90-120 minutes
Prior to tattooing or piercing: lightly exfoliate. Gently massage into skin and then apply a medium coating over the top. To prevent absorption into clothing and retain effective anesthesia, cover with an occlusive dressing. Thicker skin requires longer time to absorb the cream than thinner skin. Use Superior Super Juice during procedure to continue anesthesia, or New and Improved Blue Gel , Five-Star Vasocaine or Maximum Zone 2 to continue anesthesia and constrict blood vessels.		
PIERCING	15- 20 minutes	
Apply to eyebrows, navels and nipples. Avoid genital areas. Thicker skin requires longer pre-treating time. Lightly exfoliate. Gently massage into skin, then apply a medium coating on top. Cover with occlusive dressing. Wipe away immediately prior to piercing. Use New and Improved Blue Gel or Maximum Zone 2 during jewelry insertion to continue anesthesia and reduce swelling.		

DERMAL SOURCE
 2319 NE Glisan Street, Portland, Oregon 97232, U.S.A.
 Phone: +1 866-568-3223, Fax: 503-595-0293
www.dermalsource.com

