

**Last Menstrual Period:**


**Ordering Physician:**  
Precision Analytical

**DOB:** 1979-01-03  
**Age:** 37  
**Gender:** Female



**Collection Times:**  
2016-10-02 06:00AM  
2016-10-02 08:00AM  
2016-10-01 06:00PM  
2016-10-01 10:00PM  
2016-10-02 02:00AM

## Hormone Testing Summary


### Key (how to read the results):




### Sex Hormones See Pages 2 and 3 for a thorough breakdown of sex hormone metabolites



Progesterone Serum Equivalent is a calculated value based on urine pregnanediol.





### Adrenal Hormones See pages 4 and 5 for a more complete breakdown of adrenal hormones




Free cortisol best reflects tissue levels. Metabolized cortisol best reflects total cortisol production.

### Total DHEA Production

| Age   | Range     |
|-------|-----------|
| 20-39 | 1300-3000 |
| 40-60 | 750-2000  |
| >60   | 500-1200  |



cortisol  
metabolism



Please be sure to always read below for any specific lab comments. More detailed comments can be found on page 8.

=====  
Your DUTCH Complete report will include a summary (page 1), a list of all of the hormones tested and their ranges (pages 2,4,6) as well as a graphical representation of the results (pages 3,5). You will also see a steroid pathway for your reference (page 7) and provider notes. This report is not intended to treat, cure or diagnose any specific diseases. There is a series of videos in our video library at dutchtest.com that you may find useful in understanding the report. The following video (which can also be found on the website under the listed names along with others) may be particularly helpful in aiding your understanding:  
[DUTCH Complete Overview](#)



**Accession # 00280399**

Female Sample Report  
123 A Street  
Sometown , CA 90266



**Sex Hormones and Metabolites**

**Last Menstrual Period:**

**Ordering Physician:**  
Precision Analytical

**DOB:** 1979-01-03  
**Age:** 37  
**Gender:** Female


**Collection Times:**  
2016-10-02 06:00AM  
2016-10-02 08:00AM  
2016-10-01 06:00PM  
2016-10-01 10:00PM  
2016-10-02 02:00AM

| Test                                     | Result                  | Units  | Luteal*<br>Range | Postmenopausal<br>Range |
|--|-------------------------|--------|------------------|-------------------------|
| <b>Progesterone Metabolites (Urine)</b>  |                         |        |                  |                         |
| b-Pregnanediol                           | Below luteal range      | 196.0  | ng/mg 600 - 2000 | 60-200                  |
| a-Pregnanediol                           | Below luteal range      | 112.0  | ng/mg 200 - 740  | 15-50                   |
| <b>Estrogens and Metabolites (Urine)</b> |                         |        |                  |                         |
| Estrone(E1)                              | Above luteal range      | 28.2   | ng/mg 12 - 26    | 3.0-7.0                 |
| Estradiol(E2)                            | Above luteal range      | 5.1    | ng/mg 1.8 - 4.5  | 0.2-0.7                 |
| Estriol(E3)                              | Above luteal range      | 22.5   | ng/mg 5 - 18     | 0.6-4.0                 |
| 2-OH-E1                                  | Below luteal range      | 4.8    | ng/mg 5.1 - 13.1 | 0.3-2.0                 |
| 4-OH-E1                                  | Within luteal range     | 0.8    | ng/mg 0 - 1.8    | 0-0.3                   |
| 16-OH-E1                                 | Above luteal range      | 4.7    | ng/mg 0.7 - 2.6  | 0.2-0.6                 |
| 2-Methoxy-E1                             | Below luteal range      | 0.8    | ng/mg 2.5 - 6.5  | 0.3-1.4                 |
| 2-OH-E2                                  | Low end of luteal range | 0.19   | ng/mg 0 - 1.2    | 0-0.3                   |
| 4-OH-E2                                  | Low end of luteal range | 0.20   | ng/mg 0.15 - 0.5 | 0-0.1                   |
| 2-Methoxy-E2                             | Low end of luteal range | 0.3    | ng/mg 0.3 - 0.7  | 0-0.4                   |
| <b>Androgens and Metabolites (Urine)</b> |                         |        |                  |                         |
| DHEA-S                                   | Low end of range        | 26.0   | ng/mg 20 - 750   |                         |
| Androsterone                             | Above range             | 1810.0 | ng/mg 200 - 1650 |                         |
| Etiocholanolone                          | Within range            | 680.0  | ng/mg 200 - 1000 |                         |
| Testosterone                             | Below range             | 1.5    | ng/mg 2.3 - 14   |                         |
| 5a-DHT                                   | Above range             | 7.2    | ng/mg 0 - 6.6    |                         |
| 5a-Androstenediol                        | Above range             | 42.0   | ng/mg 12 - 30    |                         |
| 5b-Androstenediol                        | Within range            | 32.0   | ng/mg 20 - 75    |                         |
| Epi-Testosterone                         | Within range            | 8.8    | ng/mg 2.3 - 14   |                         |

\*the Luteal Range is the premenopausal range. When patients are taking oral progesterone this range for progesterone metabolites is not luteal and reflects the higher levels expected when patients take oral progesterone. This test is intended to be taken in the luteal phase of the menstrual cycle (days 19-22 of a 28 day cycle) for premenopausal women. The ranges in the table below may be used when samples are taken during the first few days (follicular) of the cycle, during ovulation (days 11-14) or when patients are on oral progesterone. See the following pages for age-dependent ranges for androgen metabolites.

| Additional Normal Ranges | Follicular | Ovulatory | Oral Pg (100mg) |
|--------------------------|------------|-----------|-----------------|
| b-Pregnanediol           | 100-300    | 100-300   | 2000-9000       |
| a-Pregnanediol           | 25-100     | 25-100    | 580-3000        |
| Estrone (E1)             | 4.0-12.0   | 22-68     | N/A             |
| Estradiol (E2)           | 1.0-2.0    | 4.0-12.0  | N/A             |

**Hormone metabolite results from the previous page are presented here as they are found in the steroid cascade. See the Provider Comments for more information on how to read the results.**





**Accession # 00280399**

Female Sample Report  
123 A Street  
Somertown , CA 90266



**Adrenal**





**Ordering Physician:**  
Precision Analytical

**DOB:** 1979-01-03  
**Age:** 37  
**Gender:** Female


**Last Menstrual Period:**

**Collection Times:**  
2016-10-02 06:00AM  
2016-10-02 08:00AM  
2016-10-01 06:00PM  
2016-10-01 10:00PM  
2016-10-02 02:00AM

| Category   | Test                           |                   | Result | Units | Normal Range |
|--|--------------------------------|-------------------|--------|-------|--------------|
| <b>Creatinine (Urine)</b>                        |                                |                   |        |       |              |
|  | Creatinine A (Waking)          | Within range      | 0.4    | mg/ml | 0.2 - 2      |
|  | Creatinine B (Morning)         | Within range      | 0.51   | mg/ml | 0.2 - 2      |
|  | Creatinine C (Afternoon)       | Within range      | 0.92   | mg/ml | 0.2 - 2      |
|  | Creatinine D (Night)           | Within range      | 1.01   | mg/ml | 0.2 - 2      |
| <b>Daily Free Cortisol and Cortisone (Urine)</b> |                                |                   |        |       |              |
|  | Cortisol A (Waking)            | Low end of range  | 12.0   | ng/mg | 12 - 55      |
|  | Cortisol B (Morning)           | Below range       | 28.0   | ng/mg | 38 - 155     |
|  | Cortisol C (Afternoon)         | Low end of range  | 9.4    | ng/mg | 7.3 - 30     |
|  | Cortisol D (Night)             | Low end of range  | 2.6    | ng/mg | 0 - 14       |
|  | Cortisone A (Waking)           | Below range       | 30.2   | ng/mg | 40 - 120     |
|  | Cortisone B (Morning)          | Below range       | 61.2   | ng/mg | 90 - 230     |
|  | Cortisone C (Afternoon)        | Below range       | 21.4   | ng/mg | 32 - 95      |
|  | Cortisone D (Night)            | Within range      | 16.2   | ng/mg | 0 - 55       |
|  | 24hr Free Cortisol             | Below range       | 52.0   | ng/mg | 80 - 230     |
|  | 24hr Free Cortisone            | Below range       | 129.0  | ng/mg | 220 - 450    |
| <b>Cortisol Metabolites and DHEA-S (Urine)</b>   |                                |                   |        |       |              |
|  | a-Tetrahydrocortisol (a-THF)   | Above range       | 400.0  | ng/mg | 75 - 370     |
|  | b-Tetrahydrocortisol (b-THF)   | Above range       | 2500.0 | ng/mg | 1050 - 2500  |
|  | b-Tetrahydrocortisone (b-THE)  | Within range      | 3030.0 | ng/mg | 1550 - 3800  |
|  | Metabolized Cortisol (THF+THE) | High end of range | 5930.0 | ng/mg | 2750 - 6500  |
|  | DHEA-S                         | Low end of range  | 26.0   | ng/mg | 20 - 750     |



Cortisol and Cortisone interconvert (11b-HSD)



The first value reported (Waking "A") for cortisol is intended to represent the "overnight" period. When patients sleep through the night, they collect just one sample. In this case, the patient woke during the night and collected (see the top of the report for the times collected). We call this value "A1" and the value from the sample collected at waking "A2." These values are used to create a "time-weighted average" to create the "A" value. The individual values are listed here for your use:  
**The middle-of-the-night "A1" sample registered a cortisol value of 6.0ng/mg.**  
**The waking "A2" sample registered a cortisol value of 18.0ng/mg.**  
**These two values are averaged together, taking into account the amount of time each one represents, to create the "A" value of approximately 12.0ng/mg that you will see on the report.**



**Accession # 00280399**  
 Female Sample Report  
 123 A Street  
 Sometown, CA 90266



**Organic Acid Tests (OATs)**

**Last Menstrual Period:**

**Ordering Physician:**  
 Precision Analytical

**DOB:** 1979-01-03  
**Age:** 37  
**Gender:** Female

**Collection Times:**  
 2016-10-02 06:00AM  
 2016-10-02 08:00AM  
 2016-10-01 06:00PM  
 2016-10-01 10:00PM  
 2016-10-02 02:00AM

| Category   | Test                           | Result            | Units      | Normal Range |
|--|--------------------------------|-------------------|------------|--------------|
| <b>Nutritional Organic Acids</b>   |                                |                   |            |              |
| Vitamin B12 Marker (may be deficient if high) - (Urine)                                  |                                |                   |            |              |
|  | Methylmalonate (MMA)           | Within range      | 1.2 ug/mg  | 0 - 2.8      |
| Vitamin B6 Marker (may be deficient if high) - (Urine)                                   |                                |                   |            |              |
|  | Xanthurenate                   | Above range       | 6.8 ug/mg  | 0 - 1.6      |
| Glutathione Marker (may be deficient if low or high) - (Urine)                           |                                |                   |            |              |
|  | Pyroglutamate                  | Below range       | 23.2 ug/mg | 37 - 70      |
| <b>Neurotransmitter Metabolites</b>  |                                |                   |            |              |
| Dopamine Metabolite - (Urine)  |                                |                   |            |              |
|  | Homovanillate (HVA)            | Low end of range  | 5.6 ug/mg  | 4.5 - 13     |
| Norepinephrine/Epinephrine Metabolite - (Urine)  |                                |                   |            |              |
|  | Vanilmandelate (VMA)           | Within range      | 4.8 ug/mg  | 2.7 - 6.4    |
| Serotonin Metabolite - (Urine)   |                                |                   |            |              |
|  | 5-Hydroxyindoleacetate (5HIAA) | Above range       | 16.0 ug/mg | 3 - 7.5      |
| Melatonin (*measured as 6-OH-Melatonin-Sulfate) - (Urine)                                |                                |                   |            |              |
|  | Melatonin* (Waking)            | Low end of range  | 18.2 ng/mg | 10 - 85      |
| Oxidative Stress / DNA Damage, measured as 8-Hydroxy-2-deoxyguanosine (8-OHdG) - (Urine) |                                |                   |            |              |
|  | 8-OHdG (Waking)                | High end of range | 4.3 ng/mg  | 0 - 5.2      |




# Steroid Pathways

## Find these Hormones on the DUTCH Complete






Primary hormones (in CAPS) are made by organs by taking up cholesterol and converting it locally. For example, progesterone, which is made from circulating precursors like pregnenolone. For example, taking DHEA can create testosterone and estrogen, but far less than is made by the testes or ovaries, respectively.






# Provider Notes

## How to read the DUTCH report


The graphic dutch dials in this report are intended for quick and easy evaluation of which hormones are out of range. Results below the left star are shaded yellow and are below range (left). Results between the stars and shaded green are within the reference range (middle). Results beyond the second star and shaded red are above the reference range (right). Some of these hormones also change with age, and the age-dependent ranges provided should also be considered.



For female reproductive hormones, a purple band is present on the dutch dials. This band represents the expected levels (reference range) for postmenopausal (or non-cycling) women.



In a few places on the graphical pages, you will see fan-style gauges. For sex hormones, you will see one for the balance between 5a/5b metabolism as well as methylation. For adrenal hormones, you will see one to represent the balance between cortisol and cortisone metabolites. These indexes simply look at the ratio of hormones for a preference. An average or "normal" ratio between the two metabolites (or groups of metabolites) will give a result in the middle (as shown here). If the ratio between the metabolites measured is "low" the gauge will lean to the left and similarly to the right if the ratio is higher than normal.



## Patient or Sample Comments

Throughout the provider comments you may find some comments specific to your situation or results. These comments will be found in this section or within another section as appropriate. Comments in other sections that are specific to your case will be in **bold**.

**The patient reports regular menstrual cycles.**

Note: The dates listed on the samples imply that they were older than our allowed 3 weeks when they were received. The instructions ask that patients freeze or refrigerate samples if they are to be held. If that is not the case, the free cortisol and cortisone levels may drop somewhat over time if the samples are too old. Other hormones tested are stable for more than 12 weeks at room temperature. Samples that are refrigerated or frozen are stable for months.

## Progesterone Metabolism

The primary role of progesterone is to balance the strong effects of estrogen. Progesterone metabolites are measured and reflect progesterone levels well because very little progesterone is found in urine, so b-Pregnanediol is typically used as a surrogate marker because it is the most abundant metabolite, but we also test the corresponding a-pregnanediol. The average of the two metabolites is reported for progesterone. If levels are in the lower part of the reference range compared to estrogen levels, symptoms of too much estrogen may occur.

When ordering the DUTCH Complete, you will see Progesterone Serum Equivalent on the summary page 1. The urine metabolites of progesterone have been proven to correlate strongly enough to serum progesterone to provide this value. The correlation is the strongest for values within the premenopausal luteal range. Urine metabolites can at times result in somewhat higher serum equivalent results in the postmenopausal range. For this reason the postmenopausal Serum Equivalent range is slightly higher than typical serum ranges. NOTE: If progesterone is taken orally (also with sublingual), these metabolites are elevated from gut metabolism and results do NOT accurately reflect serum levels.



## Estrogen Metabolism

When evaluating estrogen levels, it is important to assess the following:

- **The status (low, normal or high?) of estrogen production:**

Levels of the primary ovarian product, estradiol (the strongest estrogen), as well as "total estrogens" may be considered. For women not on HRT, consider the appropriate range (premenopausal or postmenopausal).

- **Phase I Metabolism:**

Estrogen is metabolized (primarily by the liver) down three phase I pathways. The 2-OH pathway is considered the safest because of the anti-cancer properties of 2-OH metabolites. Conversely, the 4-OH pathway is considered the most genotoxic as its metabolites can create reactive products that damage DNA. The third pathway, 16-OH creates the most estrogenic of the metabolites (although still considerably less estrogenic than estradiol) - 16-OH-E1. If overall estrogen levels are high, production of 16-OH-E1 may exacerbate high estrogen symptoms. Similarly, a woman with very low levels of estrogens, may have less low estrogen symptoms if 16-OH metabolism is preferred. For example Armamento-Villareal showed that a higher 2-OH-E1/16-OH-E1 ratio correlated to bone loss (a low estrogen symptom). Estriol is thought of as a safer (weaker) estrogen metabolite, but it is important to remember that estriol is actually 16-OH-E2, so generally patients that make a lot of the potentially protective/weak estriol may also make a lot of the estrogenic 16-OH-E1.

When evaluating phase I metabolism, it may be important to look at the ratios of the three metabolites to see which pathways are preferred relative to one another. It may also be important to compare these metabolites to the levels of the parent hormones (E1, E2). If the ratios of the three metabolites are favorable but overall levels of metabolites are much lower than E1 and E2, this may imply sluggish phase I clearance of estrogens, which can contribute to high levels of E1 and E2. Similarly, patients with excessive phase I metabolism may have low E1 and E2 levels because of high rates of clearance (as opposed to simply not making a lot of estrogen).

The pie chart will assist you in comparing the three pathway options of phase I metabolism compared to what is "normal." 2-OH metabolism can be increased by using products containing D.I.M. or I-3-C. These compounds are found (or created from) in cruciferous vegetables and are known for promoting this pathway.

**Patients typically metabolize a much higher percentage of their estrogens down the more protective 2-OH pathway in phase 1 detoxification. Diindolylmethane (DIM) or Indole-3-Carbinol containing products can help move estrogens more efficiently down this pathway. Be aware that this typically lowers most of the other estrogens, including E1 and E2 as well. If the patients are taking or considering hormone replacement therapy, these products may be considered but a higher dose of estrogen may be needed for the same clinical effect if taken at the same time.**

- **Methylation (part of phase II metabolism) of estrogens:**

After phase I metabolism, both 4-OH and 2-OH (not 16-OH) estrogens can be deactivated and eliminated by methylation. The methylation-activity index shows the patient's ratio of 2-Methoxy-E1 / 2-OH-E1 compared to what is expected. Low methylation can be caused by low levels of nutrients needed for methylation and/or genetic abnormalities (COMT, MTHFR). The COMT enzyme responsible for methylation requires magnesium and methyl donors. Deficiencies in folate or vitamin B6 or B12 can cause low levels of methyl donors. MTHFR genetic defects can make it more difficult for patients to make sufficient methyl donors. Genetic defects in COMT can make methylation poor even in the presence of adequate methyl donors.

## Androgen Metabolism

When evaluating androgen levels, it is important to assess the following:

- **The status (low, normal or high?) of DHEA:**

DHEA and androstenedione are made almost exclusively by the adrenal gland (although a smaller amount is made in the ovaries). These hormones appear in urine as DHEA-S (DHEA-Sulfate), androsterone and etiocholanolone. The best way to assess the total production of DHEA is to add up these three metabolites. This total can be seen on the first page of the DUTCH Complete (and DUTCH Plus). DHEA production decreases quite significantly with age. Age-dependent ranges can be seen on the graphical page of results.

**The Total DHEA Production (page 1) was about 2,516ng/mg which is within the overall range and also within the age-dependent range for this patient. This implies that the adrenal glands are producing appropriate DHEA levels.**

**Because inflammation blocks DHEA being converted to DHEAS, consider inflammation as a potential part of the overall clinical picture when DHEAS is significantly lower than the downstream metabolites of DHEA (Androsterone, Etiocholanolone) as seen in this case. A sulfur deficiency can also lead to adequate androgens but deficient DHEA-S.**

- **The status (low, normal or high?) of testosterone:**


Females make most of their DHEA in the adrenal gland and a fraction of that DHEA trickles down metabolically to testosterone. For premenopausal women, some testosterone is also made by the ovaries. Levels of testosterone do drop somewhat with age, but not to the degree that DHEA decreases.

**Testosterone levels for this patient were approximately 1.50ng/mg, which is low. This can cause symptoms of low energy, low motivation, loss of lean muscle mass and low libido. Depending on her age, consider testosterone HRT or supplements that support low testosterone symptoms include Zinc, Maca, Tribulus, and Shatavari. If the patient does not present with symptoms of low testosterone, you may want to also carefully evaluate 5a-metabolism (see below) and testosterone's downstream metabolites, 5a-androstenediol and 5b-androstenediol to confirm a low androgenic state.**

**• The metabolic preference for the 5a (5-alpha) or 5b (5-beta) pathway:**

5a-reductase converts testosterone into 5a-DHT (DHT), which is even more potent (~3x) than testosterone. High levels of DHT can lead to symptoms associated with too much testosterone. Metabolites created down the 5b-pathway are significantly less androgenic than their 5a counterparts. In the examples below, the example on the left shows a patient with 5b-metabolism preference. A patient with a pattern like the example on the right may have high androgen symptoms even though the hormones are in the normal range because of the likely preference for turning a lot of her testosterone into DHT. The fan-style gauge below the hormones shows the 5a or 5b preference based on etiocholanolone (5b) and androsterone (5a) results. Progesterone metabolites are also metabolized by 5a and 5b enzymes and the balance between its two metabolites can be useful to confirm a 5a or 5b preference.

Example of how to read fan-style gauge for 5a-reductase activity:



**While testosterone levels are not high, overall DHEA production is on the higher side and androgens are preferring the androgenic 5a pathway. Since the patient did not list significant symptoms of high androgens, these higher levels may be well tolerated by the patient. Since high insulin levels can lead to more DHEA production and 5a-metabolism, it may be worth exploring potential issues with blood sugar and/or insulin.**

It is important to consider DHEA and testosterone production, 5a-metabolism patterns as well as the patient symptoms. For example, a woman with higher levels of DHEA and testosterone will often have high androgen symptoms (facial hair, thinning scalp hair, etc.) exacerbated by 5a-metabolism. If, on the other hand, she prefers 5b-metabolism she may not express high androgen symptoms in spite of higher levels of testosterone because 5b is the less androgenic pathway. Testosterone levels may be better understood by also considering its downstream metabolites (5a-androstanediol, 5b-androstanediol). Technically, these metabolites can also be formed from DHEA metabolites without going through the testosterone pathway, but they generally tend to correlate with testosterone production. You will also see levels of epi-testosterone, which is not androgenic like testosterone. It happens to be produced in about the same concentrations as testosterone (this is an approximate relationship). This can be helpful to assess testosterone therapy and rare cases where testosterone may have other complexities.

**DUTCH Adrenal**

The HPA-Axis refers to the communication and interaction between the hypothalamus (H) and pituitary (P) in the brain down to the adrenal glands (A) that sit on top of your kidneys. When a physical or psychological stressor occurs, the hypothalamus tells the pituitary to make ACTH, a hormone. ACTH stimulates the adrenal glands to make the stress hormone, cortisol and to a lesser extent DHEA and DHEA-S. Normally, the HPA-axis production follows a daily pattern in which cortisol rises rather rapidly in the first 10-30 minutes after waking in order to help with energy, then gradually decreases throughout the day so that it is low at night for sleep. The cycle starts over the next morning. Abnormally high activity occurs in Cushing’s Disease where the HPA-axis is hyper-stimulated causing cortisol to be elevated all day. The opposite is known as Addison’s Disease, where cortisol is abnormally low because it is not made appropriately in response to ACTH’s stimulation. These two conditions are somewhat rare. Examples of more common conditions related to less severely abnormal cortisol levels include fatigue, depression, insomnia, fibromyalgia, anxiety, inflammation and more.

Only a fraction of cortisol is "free" and bioactive. This fraction of cortisol is very important, but levels of metabolized cortisol best represent overall production of cortisol therefore both should be taken into account to correctly assess adrenal function.

When evaluating cortisol levels, it is important to assess the following:

**• The overall up-and-down pattern of free cortisol throughout the day, looking for low and high levels:**

Abnormal results should be considered along with related symptoms. Remember that with urine results, the “waking” sample reflects the night’s total for free cortisol. The sample collected two hours after waking captures the cortisol awakening response, which is typically the time with the most cortisol secretion.

**• The sum of the free cortisol as an expression of the overall tissue cortisol exposure:**

This total of four free cortisol measurements is the best way to assess the total of free cortisol throughout the day, and this

result correlates reasonably well to a true 24-hour urine free cortisol. Do be aware that this measurement does not take into account transitory shifts in cortisol in the late morning or early afternoon.

• **The total level of cortisol metabolites:**

We call this calculation "Metabolized Cortisol" which is the sum of a-THF, b-THF and b-THE (the most abundant cortisol metabolites). While free cortisol is the best assessment for tissue levels of cortisol, it only represents 1-3% of the total produced. The majority of cortisol results in a urine metabolite and the total of these metabolites best represents the total glandular output of cortisol for the day. When overall production is much higher than free cortisol levels, cortisol clearance may be increased (as seen in hyperthyroidism, obesity, etc.) The most common reason for sluggish cortisol clearance (assumed when free cortisol levels are much higher than metabolized cortisol) is low thyroid.

**Free cortisol levels are low. Overall HPA-Axis activity may be somewhat higher than implied by free cortisol as levels of metabolized cortisol are within range. While cortisol production may be lower than needed to meet the body's demands, increased metabolism of cortisol is contributing to the lower levels of free cortisol. Increased clearance of cortisol may be caused by long-term stress, obesity, hyperthyroidism, and other conditions.**

• **A potential preference for cortisol or cortisone (the inactive form):**

Looking at the comparison between the total for free cortisol and free cortisone is NOT the best indication of a person's preference for cortisol or cortisone. The kidney converts cortisol to cortisone in the local tissue. This localized conversion can be seen by comparing cortisol (free) and cortisone levels. To see the patient's preference systemically, it is best to look at which *metabolite* predominates (THF or THE). This preference can be seen in the fan style gauge. This is known as the 11b-HSD index. The enzyme 11b-HSD II converts cortisol to cortisone in the kidneys, saliva gland and colon. 11b-HSD I is more active in the liver, fat cells and the periphery and is responsible for reactivating cortisone to cortisol. Both are then metabolized by 5a-reductase to become tetrahydrocortisol (THF) and tetrahydrocortisone (THE) respectively.

### Nutritional Organic Acids

The following three organic acids are functional markers for vitamin deficiency. These compounds essentially back up in human biochemistry when a key nutrient is missing. These three metabolites have fairly straightforward interpretations. When the markers are elevated, it is likely that the patient's cellular levels of the related nutrient may be insufficient.

#### Methylmalonate (MMA)

Methylmalonate (also known as methylmalonic acid or MMA) is a functional marker of vitamin B12 (also known as cobalamin) deficiency. When cellular levels of B12 are low either from deficiency or due to a B12 transporter gene mutation, levels of MMA increase. This marker is considered superior to measuring serum B12 levels directly. A 2012 publication by Miller showed that 20% of those tested had a genetic defect in the protein that transports B12 to cells. These patients may have a functional B12 deficiency even if serum levels of B12 are normal.

If levels of MMA are elevated, it may be advisable to increase B12 consumption. Common foods high in B12 include beef liver, sardines, lamb, wild caught salmon, grass-fed beef, nutritional yeast and eggs. Vitamin B12 levels can also be increased through supplementation of B12 (taken as cobalamin, methylcobalamin, hydroxycobalamin, or adenosylcobalamin). Symptoms of a vitamin B12 deficiency include: fatigue, brain fog, memory problems, muscle weakness, unsteady gait, numbness, tingling, depression, migraines/headaches and low blood pressure.

#### Xanthurenate

Xanthurenate (also known as xanthurenic acid) is a functional marker of vitamin B6 (also known as pyridoxine). Vitamin B6 is a critical co-factor to over 100 important reactions that occur in the human body and is stored in the highest concentrations in muscle tissue. Tryptophan is readily converted to NAD by the liver. One of the steps in this pathway requires B6. When there is insufficient B6, xanthurenate is made instead.

**In this case xanthurenate is elevated, which implies a vitamin B6 deficiency. Not only is xanthurenate an indicator of a lack of B6, it is also harmful to the human body. It complexes with insulin and decreases insulin sensitivity. In fact, rats fed xanthurenate will actually develop diabetes because of the effects on insulin. You may consider B6 supplementation in this case. Food high in B6 include turkey breast, grass-fed beef, pinto beans, avocado, pistachios, chicken, sesame and sunflower seeds.**

**While there is always some tryptophan going down the kynurenine pathway towards NAD (and possibly xanthurenate), this process is up-regulated by inflammation, estrogen and cortisol. If levels of estrogen or cortisol are high, it may exacerbate xanthurenate elevations and increase the need for B6.**

**Xanthurenate can also bind to iron and create a complex that increases DNA oxidative damage resulting in higher 8-OHdG levels. If both markers are elevated, there is likely an antioxidant insufficiency.**

#### Pyroglutamate

Pyroglutamate (also known as pyroglutamic acid) is a functional marker of glutathione deficiency. Pyroglutamate is a step in the production/recycling of glutathione. If the body cannot convert pyroglutamate forward, it will show up elevated in the urine. Alternatively, if there is little glutathione available to be recycled, pyroglutamate will be low, so when levels of pyroglutamate are high or low, there may be insufficient glutathione levels.

Glutathione is one of the most potent anti-oxidants in the human body. It is especially important in getting rid of toxins, including the reactive quinone species formed by 4-OH-E1 and 4-OH-E2. This reactive species can damage DNA if not detoxified by either methylation or glutathione.

### Neurotransmitter Metabolites

The neurotransmitters dopamine, norepinephrine and serotonin are important for human health. Measuring neurotransmitters directly (direct testing of serotonin, for example) is difficult because of their instability and their urinary

measurements are controversial with respect to how well they reflect the body's levels of these neuro-hormones. Each of these three neurotransmitters can be assessed indirectly by measuring their urine metabolites. While these metabolites are not a perfect reflection of what's going on in the brain, the scientific literature does affirm their use for a good representation of overall levels of these neurotransmitters.

### Homovanillate (HVA)

Homovanillate (also known as HVA) is the primary metabolite of dopamine, a brain and adrenal neurotransmitter that comes from tyrosine (with BH4 and iron as co-factors) and goes on to create norepinephrine (noradrenaline) and epinephrine (adrenaline).

Low levels of HVA can be due to low levels of dopamine or poor conversion of dopamine to HVA. The latter may be due to insufficient levels of SAM, Magnesium, FAD and NAD which are needed to metabolize dopamine. Low circulating dopamine may be due to insufficient BH4, iron or tyrosine. It may also be seen when adrenal function is generally low. Low dopamine levels may be associated with addictions, cravings and pleasure seeking (to boost levels) in addition to sleepiness, impulsivity, tremors, less motivation, fatigue and low mood.

Elevated HVA may be caused by generally increased adrenal hormone output or because of a copper or vitamin C deficiency (which are needed for dopamine conversion to norepinephrine). Elevations may also be caused by a number of medications or supplements including: MAO inhibitors, quercetin, tyrosine, DL-phenylalanine (DLPA), L-dopa, macuna, dopamine medication (Levodopa, Sinemet, Methyl dopa), SNRI medication (Wellbutrin), tricyclic antidepressants, amphetamines, appetite suppressants, and caffeine. Bananas also contain dopamine. Elevated dopamine may be associated with loss of memory, insomnia, agitation, hyperactivity, mania, hyper-focus, high stress and anxiety as well as addictions, cravings and pleasure seeking (to maintain high levels).

### Vanilmandelate (VMA)

Vanilmandelate (also known as VMA) is the primary metabolite of norepinephrine and epinephrine (adrenaline). The adrenal gland makes cortisol and DHEA as well as norepinephrine and epinephrine. When adrenal hormone output is generally low, VMA levels may be low. If HVA levels are significantly higher than VMA, there may be a conversion problem from dopamine to norepinephrine. This case can be caused by a copper or vitamin C deficiency. The enzymes COMT (methylation) and MAO are needed to make VMA from norepinephrine. If these enzymes are not working properly, VMA may be low when circulating norepinephrine and/or epinephrine are not low. Low levels of norepinephrine and epinephrine may be associated with addictions, cravings, fatigue, low blood pressure, low muscle tone, intolerance to exercise, depression, loss of alertness. When the body is under physical or psychological stress, VMA levels may increase. Because dopamine gets converted to norepinephrine and ultimately to VMA, the list of medications and supplements that increase HVA may also increase VMA. Elevated levels may be associated with feeling stressed, aggression, violence, impatience, anxiety, panic, worry, insomnia, paranoia, increased tingling/burning, loss of memory, pain sensitivity, high blood pressure and heart palpitations. If VMA and HVA are both extremely high, it may be necessary to rule out a neuroblastic tumor.

### 5-Hydroxyindoleacetate (5HIAA)

5-Hydroxyindoleacetate (also known as 5HIAA) is the primary metabolite of serotonin. Serotonin is often thought of as the "antidepressant" neurotransmitter (because common antidepressants aim to increase levels) however it is important to note that 90% is made in the gut and just 1% in the brain. In the gut, serotonin is required for gut motility and activates smooth muscle activity. In the brain, the dorsal raphe nucleus (DRN) in the brain stem contains the largest serotonergic nucleus. There is also a large portion of serotonin innervation in the forebrain (cerebrum, thalamus, hypothalamus, pituitary and limbic system). Approximately 40% of the DRN contains estrogen receptors demonstrating the tight relationship estrogen has with serotonin and the brain. The Estrogen Receptor Beta (ERb) upregulates mRNA of tryptophan hydroxylase which is the rate limiting step to making 5-HTP (a precursor to serotonin). If estrogen is low, it's possible less 5-HTP will be made.

**Elevated 5HIAA implies increased serotonin turnover or the consumption of related compounds. High levels are often seen in people using SSRI antidepressants like Prozac because it increases available serotonin, which is metabolized to 5HIAA. People taking supplements like 5-HTP will usually have very high levels. Because metabolism from 5-HTP to serotonin metabolites may happen in the gut, these high levels may not represent circulating (or brain exposure) serotonin levels. Certain foods (butternuts, black walnuts, plantain, pecans, pineapple, banana) can also contain serotonin. If these variables are ruled out, it may be necessary to rule out a carcinoid tumor as well if levels are VERY high. These types of tumors make very high levels of serotonin, but they are very rare.**

### Melatonin (measured as 6-OHMS)

Melatonin is not technically an adrenal or sex hormone however it is highly involved in the entire endocrine system. It is made in small amounts in the pineal gland in response to darkness and stimulated by Melanocyte Stimulating Hormone (MSH). A low MSH is associated with insomnia, an increased perception of pain, and mold exposure. Pineal melatonin (melatonin is also made in significant quantities in the gut) is associated with the circadian rhythm of all hormones (including female hormone release). It is also made in small amounts in the bone marrow, lymphocytes, epithelial cells and mast cells. Studies have shown that a urine sample collected upon waking has levels of 6-Hydroxymelatonin-sulfate (6-OHMS) that correlate well to the total levels of melatonin in blood samples taken continuously throughout the night. The DUTCH test uses the waking sample only to test levels of melatonin production.

Low melatonin levels may be associated with insomnia, poor immune response, constipation, weight gain or increased appetite. Elevated melatonin is usually caused by ingestion of melatonin through melatonin supplementation or eating melatonin-containing foods. Elevated melatonin production that is problematic is rare, but levels can be higher in patients with Chronic Fatigue Syndrome and may be phase shifted (peaking later) in some forms of depression.

### 8-OHdG (8-Hydroxy-2-deoxyguanosine)

8-OHdG (8-hydroxy-2-deoxyguanosine) results can be seen on page 4 of the DUTCH Complete (or DUTCH Plus) report. It is a marker for estimating DNA damage due to oxidative stress (ROS creation). 8-OHdG is considered pro-mutagenic as it is a biomarker for various cancer and degenerative disease initiation and promotion. It can be increased by chronic inflammation, increased cell turnover, chronic stress, hypertension, hyperglycemia/pre-diabetes/diabetes, kidney disease, IBD, chronic skin conditions (psoriasis/eczema), depression, atherosclerosis, chronic liver disease, Parkinson's (increasing levels with worsening stages), Diabetic neuropathy, COPD, bladder cancer, or insomnia. Studies have shown higher levels in patients with breast and prostate cancers. When levels are elevated it may be prudent to eliminate or reduce any causes and increase the consumption of antioxidant containing foods and/or supplements. The reference range for 8-OHdG is a more aggressive range for Functional Medicine that puts the range limit at the 80th percentile for each gender. A classic range (average plus two standard deviations) would result in a range of 0-6ng/mg for women and 0-10ng/mg for men. Seeking out the cause of oxidative stress may be more crucial if results exceed these limits.

### **Urine Hormone Testing - General Information**

What is actually measured in urine? In blood, most hormones are bound to binding proteins. A small fraction of the total hormone levels are "free" and unbound such that they are active hormones. These free hormones are not found readily in urine except for cortisol and cortisone (because they are much more water soluble than, for example, testosterone). As such, free cortisol and cortisone can be measured in urine and it is this measurement that nearly all urinary cortisol research is based upon. In the DUTCH Adrenal Profile the diurnal patterns of free cortisol and cortisone are measured by LC-MS/MS.

All other hormones measured (cortisol metabolites, DHEA, and all sex hormones) are excreted in urine predominately after the addition of a glucuronide or sulfate group (to increase water solubility for excretion). As an example, Tajic (Natural Sciences, 1968 publication) found that of the testosterone found in urine, 57-80% was testosterone-glucuronide, 14-42% was testosterone-sulfate, and negligible amounts (<1% for most) was free testosterone. The most likely source of free sex hormones in urine is from contamination from hormonal supplements. To eliminate this potential, we remove free hormones from conjugates (our testing can be used even if vaginal hormones have been given). The glucuronides and sulfates are then broken off of the parent hormones, and the measurement is made. These measurements reflect the bioavailable amount of hormone in most cases as it is only the free, nonprotein-bound fraction in blood/tissue that is available for phase II metabolism (glucuronidation and sulfation) and subsequent urine excretion.

Disclaimer: the filter paper used for sample collection is designed for blood collection, so it is technically considered "research only" for urine collection. Its proper use for urine collection has been thoroughly validated.

