

Accuris™ Taq Plus DNA Polymerase

Description

Ideally suited for problematic templates, Taq Plus combines the superior performance of our standard Taq with a proofreading polymerase. The resulting complex provides greater yields and better sensitivity for low copy number PCR and longer DNA templates. Products have a 3'-dA overhang, suitable for TA cloning. A proprietary 5X buffer system with PCR enhancers, MgCl₂, and a high ionic strength has been formulated specifically for this enzyme blend.

- Performs across a wide range of DNA templates and low copy assays
- Proprietary 5X reaction buffer includes enhancers for maximizing enzyme activity and reaction speed
- Improved yields for longer templates with standard or fast cycling

Storage

Upon receipt, immediately store at -20°C. Avoid excessive freeze/thaw cycles. When stored as directed, this product will retain its activity for 12 months from date of receipt. The product may also be stored at 4°C for up to one month.

Limitations of Use

For research purposes only. Not intended for therapeutic or diagnostic use.

Quality Control

Accuris enzymes and reagents are tested under general assay conditions for activity, reproducibility, efficiency, heat activation, sensitivity, and absence of nuclease contamination and nuclease activity. This product is manufactured under a comprehensive quality management system, following ISO 9001:2008 standards.

General Guidelines

1. Reaction Buffer

The supplied 5X reaction buffer has been formulated for maximum efficiency, sensitivity and successful PCR with long and difficult templates. Proprietary PCR enhancers and 15mM MgCl₂ are included in the buffer. Use of additional PCR enhancers may have a negative effect on the reaction.

2. Template

For PCR of complex genomic DNA, 5ng - 500ng of template DNA may be added per reaction. Do not add more than 100ng of DNA for cDNA or plasmid DNA

3. Primers

Primers should have a predicted melting temperature of approximately 60°C, using default Primer 3 settings (<http://frodo.wi.mit.edu/primer3>). The final primer concentration should be 0.2µM to 0.6µM.

4. Annealing Temperature

An initial annealing temperature of 57°C is recommended. If nonspecific products appear, increase the temperature in 2°C increments. Alternately, a temperature gradient may be performed.

5. Extension

The polymerase performs optimally at 72°C. Extension time is dependent upon amplicon complexity and length. Generally, 30 seconds per kb is recommended for eukaryotic genomic DNA and cDNA less than 5kb. Allow 45-60 seconds per kb for longer amplicons.

Technical Support

For trouble-shooting and tech support, contact us at info@accuris-usa.com or call 908 769-5555.

Accuris is not responsible for consequential or incidental damages, whether direct or indirect, resulting from use of this product. Accuris guarantees the performance of this product as described when used in accordance with these instructions.

Reaction setup

Prepare the reaction as follows:

Component	25 µl reaction	50 µl reaction	Final concentration
Accuris 5x Taq Plus Reaction Buffer	5 µl	10 µl	1X
100mM dNTPs (25mM each)	0.25 µl	0.5 µl	1mM
Forward Primer (10µM)	1.0 µl	2.0 µl	400 nM
Reverse Primer (10µM)	1.0 µl	2.0 µl	400 nM
Template DNA	<100ng cDNA, <500ng genomic		variable
Accuris Taq Plus (5u/µl)	0.12 µl - 0.5 µl	0.25 µl - 1 µl	variable
PCR-grade water	to final reaction volume		

For other volumes, adjust the amount of each component accordingly.

Gently mix the solution. If needed, spin briefly in a microcentrifuge to bring reaction mixture to the bottom of the tube. Transfer samples to a thermal cycler and begin cycling. Be sure to include an initial activation/denaturation step of 1-2 minutes.

Routine PCR Cycling

Step	Temperature	Time
Enzyme activation	95°C	1-2 minutes
	95°C	15 seconds
25-40 cycles	55°C to 67°C**	15 seconds
	72°C	30 seconds per Kb*

*See "General Guidelines" for amplification of longer templates

Package contents and reordering

Accuris Taq Plus Polymerase is supplied at a concentration of 5 units/µl and is available in 250 and 500 unit packages. Supplied with 5X Taq Plus Buffer.

Accuris Taq Plus, Sample Pack

Catalog number PR1000-TP-S
Includes 10 µl of enzyme and 5X buffer.

Accuris Taq Plus, 250 units

Catalog number PR1000-TP-250
Includes 50 µl of enzyme and 2ml of 5X buffer in 1ml aliquots.

Accuris Taq Plus, 500 units

Catalog number PR1000-TP-500
Includes 100 µl of enzyme in 50µl aliquots, and 4ml of 5X buffer in 1ml aliquots.

Accuris offers a full line of PCR enzymes and master mixes. Visit www.accuris-usa.com for details. 09/21 slj



Accuris™
Taq Plus
PR1000-TP

5 units/µl, supplied with 5X Buffer

Package contains:

■ 250 units Taq Plus Polymerase (1x50µl)
2ml 5X Accuris Taq Plus Buffer (2x1ml)

■ 500 units Taq Plus Polymerase (2x50µl)
4ml 5X Accuris Taq Plus Buffer (4x1ml)

Store at -20°C upon receipt

PH: 908.769.5555 EM: Info@accuris-usa.com