


PENIS **EXTENDER**

INCLUDES:



A. FOAM RING (x1)



Screw Nuts

B. BASE (x1)



C. CRADLE (x1)



D. THREADED EXTENSION ROD 1 (x2)



E. THREADED EXTENSION ROD 2 (x2)



F. SILICONE RODS -FASTENING DEVICE (**x2**)




G. SILICONE STRAP -FASTENING DEVICE (x1)

ASSEMBLY INSTRUCTIONS:


Customize your experience!

Other fastening devices, foam ring, and extension rods can be to mix and match based on your preferences.




Your product arrives with four threaded extension rods for a customizable length. Simply screw the desired extension rods end to end to attach. Then, bend the base down for use.

The cradle piece is designed to rest on top of the screw nuts, furthest away from the base ring. To attach the cradle, simply slide the threaded extension rods through each of the holes on the side of the cradle. To adjust the palcement of the cradle, twist the screw nuts up or down the threaded extension rods.



Your cradle is designed with slits and holes to be used with your desired fastening device.



3b

To use the silicone strap, simply slide one end of the strap through one of the slits with the notches facing outward. Once the strap is secured into the slit, slide the other end of the strap through the opposing slit and tighten as desired.

To use the silicone rods, simply slide one end of the rod through one of these holes and secure the end of the rod to the fastener on the bottom of the cradle. Once the rod is secured into the hole, slide the other end of the rod through the opposing hole, tighten as desired, and secure to the opposing fastener.

This non-medical device is intended as a sexual enhancing aid only. Not intended for contraceptive/birth control or treatment of erection conditions. Use no longer than 30 minutes at a time. Allow 60 minutes between additional uses. Do not fall asleep under the influence of alcohol or drugs while using this product. Prolonged use (without removal) may result in bruising at the base or permanent injury to the penis. Consult your physician should any complication occur and discontinue further use immediately.