



UNISCIENCE
Corporation

For research purpose only. Not for use in diagnostic procedures for clinical purposes. For IN VITRO USE ONLY.

UniSafe Dye Nucleic Acid Staining Solution (20,000x)

Cat. No. UNI-R01031 1 ml

DESCRIPTION

UniSafe Dye Nucleic Acid Staining Solution (20,000x) is a new and safe nucleic acid stain, an alternative to the traditional ethidium bromide (EtBr) stain for detecting nucleic acid in agarose gels. It emits green fluorescence when bound to DNA or RNA. This new stain has two fluorescence excitation maxima when bound to nucleic acid, one centered at 309 nm and another at 419 nm. In addition, it has one visible excitation at 514 nm. The fluorescence emission of UniSafe Dye bound to DNA is centered at 537 nm. UniSafe Dye Nucleic Acid Staining Solution (20,000x) is as sensitive as EtBr. The staining protocol for UniSafe Dye Nucleic Acid Staining Solution (20,000x) is similar to that for EtBr. UniSafe Dye Nucleic Acid Staining Solution (20,000x) had a negative result in mouse marrow chromophilous erythrocyte micronucleus test and also in mouse spermary spermatocyte chromosomal aberration test. So it is wise to choose UniSafe Dye Nucleic acid Staing Solution (20,000x) instead of EtBr for detecting nucleic acids in agarose gels.

CHARACTERISTICS

- Used for detecting double-strand DNA and single-stranded RNA
- Alternative to the ethidium bromide staining
- As sensitive as EtBr or more sensitive than that
- Non-toxic, non-mutagenic and non-carcinogenic
- No hazard waste

CONTENTS

- UniSafe Dye Nucleic Acid Staining Solution (20,000x) 1 ml

STORAGE CONDITION

- Store at room temperature and stable for 12 months. For more stable use, should store at 4 °C (Stable for 24 months).

APPLICATION

- Visualization of DNA and RNA bands during agarose gel electrophoresis
- Isolation of DNA fragments for subcloning without introducing mutations normally caused by EtBr.

CONSIDERATION BEFORE USE

- UniSafe Dye Nucleic Acid Staing Solution (20,000x) is non-carcinogenic but may cause skin and eye irritations. Please wear gloves when working with the product.

PROTOCOL


1. Prepare a 100 ml of agarose gel solution (concentration from 0.8~3 %) in a 250 ml flask and mix it thoroughly. Place the flask in the microwave, heat in until the solution is completely clear (about 2~3 minutes).
Note: The thickness of gel should be less than 0.5 cm since thick gels may decrease sensitivity.
2. Add 5 μ l of UniSafe Dye Nucleic Acid Staining Solution (20,000x) to the agarose solution. Swirl the flask gently to mix the solution and avoid forming bubbles.
3. While the agarose solution cools, pour it into the gel tray until the comb teeth are immersed about 1/4~1/2 into the agarose.
Note: Repeated melting of gels containing UniSafe Dye Nucleic Acid Staining Solution (20,000x) may result in low sensitivity.
4. Allow the agarose gel to cool until solidified. Load samples on the gel and perform electrophoresis.
5. Detect the bands under UV illumination.
Note: UniSafe Dye Nucleic Acid Staining Solution (20,000x) allows visualization of DNA(>50 ng) in the agarose gel under visible light. This eliminates the need for exposure to UV light, which may nick and damage DNA. The intact DNA fragments purified from agarose gel can increase the efficiency of subsequent molecular biology manipulations such as cloning, transformation and transcription.



EXPERIMENTAL INFORMATION

• Spectrum

1. Excitation Wavelength (EX) Spectrum




EM : 540.0 nm
 Data Mode : Fluorescence
 Scan Speed : 2400 nm / min
 PMT Voltage : 400 V
 Slit (EX/EM) : 5.0 nm / 5.0 nm
 Response : Auto

NO	Wavelength (nm)	Peak
1	308.8	38.17
2	419.2	35.93
3	513.8	87.06

Fig. 1. Measurement of fluorescence excitation wavelength
 UniSafe Dye Nucleic Acid Staining Solution (20,000x) has two fluorescence excitation maxima, one centered at 309 nm and another at 419 nm. In addition, it has one visible excitation at 514 nm.

2. Emission wavelength (EM) Spectrum



EX : 416.0 nm
 Data Mode : Fluorescence
 Scan Speed : 2400 nm/min
 PMT Voltage : 400 V
 Slit (EX/EM) : 5.0 nm / 5.0 nm
 Response : Auto

NO	Wavelength (nm)	Peak
1	537.2	36.26

Fig. 2. Measurement of fluorescence emission wavelength
 The fluorescence emission of UniSafe Dye Nucleic Acid Staining Solution (20,000x) bound to DNA is centered at 537 nm.

• Sensitivity

1. DNA

Sensitivity of DNA detection of UniSafe Dye Nucleic Acid Staining Solution (20,000x) under UV transmission




Fig. 3. Gel analysis of serially diluted genomic DNA using UniSafe Dye Nucleic Acid Staining Solution (20,000x) and EtBr
 Genomic DNA was extracted from SNU-1 cells using G-DEX™ IIc Genomic DNA Extraction Kit (Cell/Tissue)(Cat. No. 17231)
Lane1, 5 ng of gDNA; **lane2**, 10 ng of gDNA; **lane3**, 20 ng of gDNA; **lane4**, 30 ng of gDNA; **lane 5**, 40 ng of gDNA; **lane 6**, 50 ng of gDNA; **lane 7**, 60 ng of gDNA; **lane 8**, 70 ng of gDNA.




Fig. 4. Gel analysis of diluted 100bp Ladder Molecular Weight DNA Marker (Cat. No. 24012) and 1kb Ladder Molecular Weight DNA Marker (Cat. No. 24022) using UniSafe Dye Nucleic Acid Staining Solution (20,000x) and EtBr

100bp Ladder DNA marker and 1kb ladder DNA marker were serially diluted from 2^0 to 2^{-5} . **Lane 1**, 800 ng of 100bp ladder DNA marker; **lane 2**, 400 ng of 100bp ladder DNA marker; **lane 3**, 200 ng of 100bp ladder DNA marker; **lane 4**, 100 ng of 100bp ladder DNA marker; **lane 5**, 50 ng of 100bp ladder DNA marker; **lane 6**, 25 ng of 100bp ladder DNA marker; **lane 7**, 800 ng of 1kb ladder DNA marker; **lane 8**, 400 ng of 1kb ladder DNA marker; **lane 9**, 200 ng of 1kb ladder DNA marker; **lane 10**, 100 ng of 1kb ladder DNA marker; **lane 11**, 50 ng of 1kb ladder DNA marker; **lane 12**, 25 ng of 1kb ladder DNA marker.




Fig. 5. Gel analysis of different size of PCR products using UniSafe Dye Nucleic Acid Staining Solution (20,000x) and EtBr

PCR products were synthesized using *Maxime™* PCR PreMix (*i-Taq*)(Cat. No. 25025), *Maxime™* PCR PreMix (*i-StarTaq*)(Cat. No. 25165) and *Maxime™* PCR PreMix (*i-pfu*) (Cat. No. 25185).

Lane M1, 100bp ladder DNA marker; **lane 1**, 161bp size of dsDNA; **lane 2**, 218bp size of dsDNA; **lane 3**, 375bp size of dsDNA; **lane 4**, 575bp size of dsDNA; **lane 5**, 1kb size of dsDNA; **lane 6**, 4.5kb size of dsDNA; **lane 7**, 9kb size of dsDNA; **lane M2**, 1kb ladder DNA marker.

2. RNA

Sensitivity of RNA detection of UniSafe Dye Nucleic Acid Staining Solution (20,000x) under UV transmission




Fig. 6. Gel analysis of diluted RNA using UniSafe Dye Nucleic Acid Staining Solution (20,000x) and EtBr

Total RNA was isolated from K562 cells using easy-BLUE™ Total RNA Extraction Kit (Cat. No. 17061).

Lane M, 1kb ladder DNA marker; **lane 1**, 25 ng of RNA; **lane 2**, 50 ng of RNA; **lane 3**, 100 ng of RNA; **lane 4**, 200 ng of RNA; **lane 5**, 400 ng of RNA.