



## Core SHRED Product References

Lee J.O., Kim N., Lee H.J., Moon J.W., Lee S.K., Kim S.J., Kim J.K., Park S.H., Kim H.S. [6]-Gingerol affects glucose metabolism by dual regulation via the AMPK $\alpha$ 2-mediated AS160-Rab5 pathway and AMPK-mediated insulin sensitizing effects. *J. Cell. Biochem.* 2015;116:1401–1410.

Li Y., Tran V.H., Koolaji N., Duke C., Roufogalis B.D. (S)-[6]-Gingerol enhances glucose uptake in L6 myotubes by activation of AMPK in response to [Ca<sup>2+</sup>]<sub>i</sub>. *J. Pharm. Pharm. Sci.* 2013;16:304–312.

Son M.J., Miura Y., Yagasaki K. Mechanisms for antidiabetic effect of gingerol in cultured cells and obese diabetic model mice. *Cytotechnology.* 2015;67:641–652.

Chien-Kei Wei, Yi-Hong Tsai, Michal Korinek, et al. 6-Paradol and 6-Shogaol, the Pungent Compounds of Ginger, Promote Glucose Utilization in Adipocytes and Myotubes, and 6-Paradol Reduces Blood Glucose in High-Fat Diet-Fed Mice. *Int J Mol Sci.* 2017 Jan; 18(1): 168.

Alasbahi RH, Melzig MF. Forskolin and derivatives as tools for studying the role of cAMP. *Pharmazie.* 2012 Jan;67(1):5-13.

Godard MP, et al. Body composition and hormonal adaptations associated with forskolin consumption in overweight and obese men. *Obes Res.* 2005 Aug;13(8):1335-43.

Henderson S, et al. Effects of coelus forskohlii supplementation on body composition and hematological profiles in mildly overweight women. *J Int Soc Sports Nutr.* 2005 Dec 9;2:54-62.

Tamboli ET ,et al. Metabolic diversity in Coleus forskohlii Briq. of Indian subcontinent. *Nat Prod Res.* 2013;27(19):1737-42.

Greenway FL, et al. Topical fat reduction. *Obes Res.* 1995 Nov;3 Suppl 4:561S-568S.  
Shivaprasad HN, et al. Effect of Coleus forskohlii extract on cafeteria diet induced obesity in rats. *Pharmacognosy Res.* 2014 Jan;6(1):42-5.

Alarcon-Aguilara F. J., Roman-Ramos R., Perez-Gutierrez S., Aguilar-Contreras A., Contreras-Weber C. C., Flores-Saenz J. L. Study of the anti-hyperglycemic effect of plants used as antidiabetics. *Journal of Ethnopharmacology.* 1998;61(2):101–110.

**Crush It**

[www.corenutrionals.com](http://www.corenutrionals.com)

Amel B. Traditional treatment of high blood pressure and diabetes in Souk Ahras district. *Journal of Pharmacognosy and Phytotherapy*. 2013;5(1):12–20.

Al-Azzawie H. F., Alhamdani M.-S. S. Hypoglycemic and antioxidant effect of oleuropein in alloxan-diabetic rabbits. *Life Sciences*. 2006;78(12):1371–1377.

Sato H., Genet C., Strehle A., et al. Anti-hyperglycemic activity of a TGR5 agonist isolated from *Olea europaea*. *Biochemical and Biophysical Research Communications*. 2007;362(4):793–798.

Jemai H., Feki A. E. L., Sayadi S. Antidiabetic and antioxidant effects of hydroxytyrosol and oleuropein from olive leaves in alloxan-diabetic rats. *Journal of Agricultural and Food Chemistry*. 2009;57(19):8798–8804.

Eidi A., Eidi M., Darzi R. Antidiabetic effect of *Olea europaea* L. in normal and diabetic rats. *Phytotherapy Research*. 2009;23(3):347–350.

Cumaoglu A., Rackova L., Stefek M., Kartal M., Maechler P., Karasu Ç. Effects of olive leaf polyphenols against H<sub>2</sub>O<sub>2</sub> toxicity in insulin secreting  $\beta$ -cells. *Acta Biochimica Polonica*. 2011;58(1).

Barbaro B., Toietta G., Maggio R., et al. Effects of the olive-derived polyphenol oleuropein on human health. *International Journal of Molecular Sciences*. 2014;15(10):18508–18524.

Dubey V. K., Patil C. R., Kamble S. M., et al. Oleanolic acid prevents progression of streptozotocin induced diabetic nephropathy and protects renal microstructures in Sprague Dawley rats. *Journal of Pharmacology & Pharmacotherapeutics*. 2013;4(1):47–52.

Al-Qarawi A. A., Al-Damegh M. A., ElMougy S. A. Effect of freeze dried extract of *Olea europaea* on the pituitary-thyroid axis in rats. *Phytotherapy Research*. 2002;16(3):286–287.

Reinbach HC, et al. *Clin Nutr*. 2009; 28(3): 260–265.

Hachiya S, et al. *Biosci Biotechnol Biochem*. 2007; 71(3): 671–676.

Iida T, et al. *Neuropharmacology*. 2003; 44: 958–967.

Kobata K, et al. *J Agric Food Chem*. 1998; 46: 1695–1697.

Spriet et al. (2014) *Sports Med* 44 (Suppl 1): S87-96.