

Core PUDDN Product References



Madzima, T. A., Melanson, J. T., Black, J. R., & Nepocatych, S. (2018). Pre-Sleep Consumption of Casein and Whey Protein: Effects on Morning Metabolism and Resistance Exercise Performance in Active Women. *Nutrients*, *10*(9), 1273. <u>https://doi.org/10.3390/nu10091273</u>

Joy, J.M., Vogel, R.M., Shane Broughton, K. *et al.* Daytime and nighttime casein supplements similarly increase muscle size and strength in response to resistance training earlier in the day: a preliminary investigation. *J Int Soc Sports Nutr* **15**, 24 (2018). <u>https://doi.org/10.1186/s12970-018-0228-9</u>

Snijders, T., Trommelen, J., Kouw, I., Holwerda, A. M., Verdijk, L. B., & van Loon, L. (2019). The Impact of Pre-sleep Protein Ingestion on the Skeletal Muscle Adaptive Response to Exercise in Humans: An Update. *Frontiers in nutrition*, *6*, 17. <u>https://doi.org/10.3389/fnut.2019.00017</u>

Babayan, V. K. (1987) Medium-chain triglycerides and structured lipids. Lipids 22:417-420.

Bach, A. C. & Babayan, V. K. (1982) Medium-chain triglycerides: an update. Am. J. Clin. Nutr. 36:950-962.

Scalfi, L, Coltorti, A. & Contaldo, F. (1991) Postprandial thermogenesis in lean and obese subjects after meals supplemented with medium-chain and long-chain triglycerides. Am. J. Clin. Nutr. 53:1130-1133.

Seaton, T. B., Welle, S. L, Warenko, M. K. & Campbell, R. G. (1986) Thermic effect of medium-chain and longchain triglycerides in man. Am. J. Clin. Nutr. 44:630-634.

Dulloo, A. G., Fathi, M., Mensi, N. & Girardier, L. (1996) Twenty-four-hour energy expenditure and urinary catecholamines of humans consuming low-to-moderate amounts of medium-chain triglycerides: a dose-response study in human respiratory chamber. Eur. J. Clin. Nutr. 50:152-155.



CORE NUTRITIONALS 1-888-978-2332

820 N. Pollard St., RTL#3 Arlington, VA 22203 info@corenutritionals.com www.corenutritionals.com



Flatt, J. P., Ravussin, E., Acheson, K. J. & Jequier, E. (1985) Effects of dietary fat on postprandial substrate oxidation and on carbohydrate and fat balances. J. Clin. Investig. 76:1019-1024.

Hill, J. O., Peters, J. C., Yang, D., Sharp, T., Kaler, M., Abumrad, N. N. & Greene, H. L. (1989) Thermogenesis in humans during overfeeding with medium-chain triglycerides. Metabolism 38:641-648

White, M. D., Papamandjaris, A. A. & Jones, P.J.H. (1999) Enhanced postprandial energy expenditure with medium-chain fatty acid feeding is attenuated after 14 d in premenopausal women. Am. J. Clin. Nutr. 69:883-889.

Astbury NM, et al. Dose-response effect of a whey protein preload on within-day energy intake in lean subjects. Br J Nutr. (2010)

Frestedt JL, et al. A whey-protein supplement increases fat loss and spares lean muscle in obese subjects: a randomized human clinical study. Nutr Metab (Lond). (2008)

Zhu K, et al. The effects of a two-year randomized, controlled trial of whey protein supplementation on bone structure, IGF-1, and urinary calcium excretion in older postmenopausal women. J Bone Miner Res. (2011)

Rennie MJ, et al. Control of the size of the human muscle mass. Annu Rev Physiol. (2004)

Anabolic signalling deficits underlie amino acid resistance of wasting, aging muscle

Nutrient signalling in the regulation of human muscle protein synthesis

Leucine supplementation improves muscle protein synthesis in elderly men independently of hyperaminoacidaemia

Tang JE, et al. Ingestion of whey hydrolysate, casein, or soy protein isolate: effects on mixed muscle protein synthesis at rest and following resistance exercise in young men. J Appl Physiol. (2009)

Whey protein stimulates postprandial muscle protein accretion more effectively than do casein and casein hydrolysate in older men

Dangin M, et al. The digestion rate of protein is an independent regulating factor of postprandial protein retention. Am J Physiol Endocrinol Metab. (2001)



CORE NUTRITIONALS 1-888-978-2332

820 N. Pollard St., RTL#3 Arlington, VA 22203 info@corenutritionals.com www.corenutritionals.com



Engelen MP, et al. Casein protein results in higher prandial and exercise induced whole body protein anabolism than whey protein in Chronic Obstructive Pulmonary Disease. Metabolism. (2012)

Pal S, Ellis V, Dhaliwal S. Effects of whey protein isolate on body composition, lipids, insulin and glucose in overweight and obese individuals. Br J Nutr. (2010)

Haraguchi FK, et al. Evaluation of biological and biochemical quality of whey protein. J Med Food. (2010)

de Aguilar-Nascimento JE, Prado Silveira BR, Dock-Nascimento DB. Early enteral nutrition with whey protein or casein in elderly patients with acute ischemic stroke: a double-blind randomized trial. Nutrition. (2011)

Candow DG, et al. Effect of whey and soy protein supplementation combined with resistance training in young adults. Int J Sport Nutr Exerc Metab. (2006)



820 N. Pollard St., RTL#3 Arlington, VA 22203 info@corenutritionals.com www.corenutritionals.com