

Core HARD Product References



Kingsley M, Wadsworth D, Kilduff LP, McEneny J, Benton D; Wadsworth; Kilduff; McEneny; Benton (August 2005). "Effects of phosphatidylserine on oxidative stress following intermittent running". *Medicine and Science in Sports and Exercise* 37 (8): 1300–6.

Kingsley MI, Miller M, Kilduff LP, McEneny J, Benton D; Miller; Kilduff; McEneny; Benton (January 2006). "Effects of phosphatidylserine on exercise capacity during cycling in active males". *Medicine and Science in Sports and Exercise* 38 (1): 64–71.

Jäger R, Purpura M, Geiss K-R, Weiß M, Baumeister J, Amatulli F, Schröder L, Herwegen H; Purpura; Geiss; Weiß; Baumeister; Amatulli; Schröder; Herwegen (December 2007). "The effect of phosphatidylserine on golf performance". *International Society of Sports Nutrition* 4 (1): 23.

Starks MA, Starks SL, Kingsley M, Purpura M, Jäger R; Starks; Kingsley; Purpura; Jäger (July 2008). "The effects of phosphatidylserine on endocrine response to moderate intensity exercise". *Journal of the International Society of Sports Nutrition* 5 (1): 11.

Monteleone P, Maj M, Beinat L, Natale M, Kemali D; Maj; Beinat; Natale; Kemali (1992). "Blunting by chronic phosphatidylserine administration of the stress-induced activation of the hypothalamo-pituitary-adrenal axis in healthy men". *European Journal of Clinical Pharmacology* 42 (4): 385–8. doi:10.1007/BF00280123 (inactive 2015-02-01). PMID 1325348.

Fernholz KM, Seifert JG, Bacharach DW, Burke ER, Gazal O (2000). "The Effects of Phosphatidyl Serine on Markers of Muscular Stress in Endurance Runners [abstract]". *Medicine and Science in Sports and Exercise* 32 (4): S321.

Bhattacharya SK, Muruganandam AV. Adaptogenic activity of *Withania somnifera*: An experimental study using a rat model of chronic stress. *Pharmacol Biochem Behav* 2003;75:547-55.

Singh G, Sharma PK, Dudhe R, Singh S. Biological activities of *Withania somnifera*. *Ann Biol Res* 2010;1:56-63.

Sharma V, Sharma S, Pracheta, Paliwal R. *Withania somnifera*: A rejuvenating ayurvedic medicinal herb for the treatment of various human ailments. *Int J PharmTech Res* 2011;3:187-92.

Kulkarni SK, Dhir A. *Withania somnifera*: An Indian ginseng. *Prog Neuro-Psychopharmacol Biol Psychiatry* 2008;32:1093-05.

Bhattacharya SK, Goel RK, Kaur R, Ghosal S. Antistress activity of sitoindosides VII and VIII, new acylsterylglucosides from *Withania somnifera*. *Phytother Res* 1987;1:32-7.

Ghosal S, Lal J, Srivastava R, Bhattacharya SK, Upadhyay SN, Jaiswal AK, et al. Immunomodulatory and CNS effects of sitoindosides IX and X, two new glycowithanolides from *Withania somnifera*. *Phytother Res* 1989;3:201-6.

Park, JB. N-coumaroyldopamine and N-caffeoyldopamine increase cAMP via beta 2-adrenoceptors in myelocytic U937 cells. *FASEB J.* 2005 Apr;19(6):497-502.

St-Onge. Dietary fats, teas, dairy and nuts: potential functional food for weight control? *Am J Clin Nutr* 2005 81(1):7-15.

Kuriyama S et al. Green Tea Consumption and Mortality Due to Cardiovascular Disease, Cancer, and All Causes in Japan The Ohsaki Study. *JAMA* 2006;296:1255-1265.

Coimbra S et al. Green tea consumption improves plasma lipid profiles in adults. *Nutrition Research* 2006;26(11):604-607.

Bryan J. Psychological effects of dietary components of tea: caffeine and L-theanine. *Nutr Rev* 2008;66(2):82-90.

Nobre AC, Rao A, Owen GN. L-theanine, a natural constituent in tea, and its effect on mental state. *Asia Pac J Clin Nutr* 2008;17 Suppl 1:167-8.

Kimura K, Ozeki M, Juneja LR, Ohira H. L-Theanine reduces psychological and physiological stress responses. *Biol Psychol* 2007;74(1):39-45.

L-theanine. Monograph. *Altern Med Rev* 2005;10(2):136-8.

Dulloo AG, Duret C, Rohrer D, Girardier L, Mensi N, Fathi M, Chantre P, Vandermander J. Efficacy of a green tea extract rich in catechin polyphenols and caffeine in increasing 24-h energy expenditure and fat oxidation in humans. *Am J Clin Nutr* 1999;70(6):1040-5.

Hertog MGL, Feskens EJM, Hollman PCH, et al. Dietary antioxidant flavonoids and risk of coronary disease: the Zutphen Elderly Study. *Lancet* 1993;342:1007-11.

Keli SO, Hertog MGL, Feskens EJM, Kromhout D. Dietary flavonoids, antioxidant vitamins, and incidence of stroke. *Arch Intern Med* 1996;156:637-42.

Duffy SJ, Keaney JF Jr, Holbrook M, Gokce N, Swerdloff PL, Frei B, Vita JA. Short- and long-term black tea consumption reverses endothelial dysfunction in patients with coronary artery disease. *Circulation* 2001;104:151-6.

Isemura M, Saeki K, Kimura T, Hayakawa S, Minami T, Sazuka M. Tea catechins and related polyphenols as anti-cancer agents. *Biofactors.* 2000;13(1-4):81-5.



Hakim IA, Alsaif MA, Alduwaihy M, Al-Rubeaan K, Al-Nuaim AR, Al-Attas OS. Tea consumption and the prevalence of coronary heart disease in Saudi adults: results from a Saudi national study. *Prev Med* 2003;36(1):64-70.

Sesso HD, Gaziano JM, Buring JE, Hennekens CH. Coffee and tea intake and the risk of myocardial infarction. *Am J Epidemiol* 1999;149:162-7.

Meilahn EN, De Stavola B, Allen DS, Fentiman I, Bradlow HL, Sepkovic DW, Kuller LH. Do urinary estrogen metabolites predict breast cancer? Follow up of the Guernsey III cohort. *Br J Cancer*. 1998;78:1250-1255.

Schneider J, Huh MM, Bradlow HL, Fishman J. Antiestrogen action of 2-hydroxyestrone on MCF-7 human breast cancer cells. *J Biol Chem*. 1984;259:4840-4845.

Telang NT, Suto A, Wong GY, Osborne MP, Bradlow HL. Induction by estrogen metabolite 16 alpha-hydroxyestrone of genotoxic damage and aberrant proliferation in mouse mammary epithelial cells. *J Natl Cancer Inst*. 1992;84:634-638.

Ho GH, Luo XW, Ji CY, Foo SC, Ng EH. Urinary 2/16alpha-hydroxyestrone ratio: correlation with serum insulin-like growth factor binding protein-3 and a potential biomarker of breast cancer risk. *Ann Acad Med Singap*. 1998;27:294-299.

Zheng W, Dunning L, Jin F, Holtzman J. Urinary estrogen metabolites and breast cancer: a case-control study. *Cancer Epidemiol Biomark Prev*. 1997;6:505-509.

Ursin G, London S, Stanczk FZ, Gentschein E, Paganini-Hill A, Ross RK, Pike MC. Urinary 2-hydroxyestrone/16alpha-hydroxyestrone ratio and risk of breast cancer in postmenopausal women. *J Natl Cancer Inst*. 1999;91:1067-1072.

Muti P, Bradlow HL, Micheli A, Krogh V, Freudenheim JL, Schunemann HJ, Stanulla M, Yang J, Sepkovic DW, Trevisan M, Berrino F. Metabolism and risk of breast cancer: a prospective analysis of 2:16 hydroxyestrone ratio and risk of breast cancer in premenopausal and postmenopausal women. *Cancer Epidemiol*. 2000;11:635-640.

Tiwari RK, Guo L, Bradlow HL, Telang NT, Osborne MP. Selective responsiveness of human breast cancer cells to indole-3-carbinol, a chemopreventive agent. *J Natl Cancer Inst*. 1994;86:126-131.

Sepkovic DW, Bradlow HL, Bell M. Quantitative determination of 3,3'-diindolylmethane in the urine of individuals receiving indole-3-carbinol. *Nutr Cancer*. 2002;41:57-63.

Li Y, Wang Z, Kong D, Murthy S, Dou QP, Sheng S, Reddy GP, Sarkar FH. Regulation of FOXO3a/ β -catenin/GSK-3 β signaling by 3,3'-diindolylmethane contributes to inhibition of cell proliferation and induction of apoptosis in prostate cancer cells. *J Biol Chem*. 2007;282:21542-21550.

Rahman KM, Ali S, Aboukameel A, Sarkar SH, Wang Z, Philip PA, Sakr WA, Raz A. Inactivation of NF- κ B by 3,3'-diindolylmethane contributes to increased apoptosis induced by chemotherapeutic agent in breast cancer cells. *Mol Cancer Ther*. 2007;6:2757-2765.





Core Nutritionals, LLC
820 N. Pollard St., RTL#3
Arlington, VA 22203
1-888-978-2332

Martínez-Pinilla, E., Oñatibia-Astibia, A., & Franco, R. (2015). The relevance of theobromine for the beneficial effects of cocoa consumption. *Frontiers in pharmacology*, 6, 30. <https://doi.org/10.3389/fphar.2015.00030>

Cocoabuterol®. (n.d.). Retrieved June 25, 2020, from <http://cocoabuterol.com/advantage.php>



CORE NUTRITIONALS
1-888-978-2332

820 N. Pollard St., RTL#3
Arlington, VA 22203

info@corenutritionals.com
www.corenutritionals.com