



# Introducing highly allergenic foods to infants and children

**Author:** David M Fleischer, MD

**Section Editor:** Scott H Sicherer, MD, FAACAP

**Deputy Editor:** Elizabeth TePas, MD, MS

All topics are updated as new evidence becomes available and our [peer review process](#) is complete.

**Literature review current through:** Nov 2019. | **This topic last updated:** Apr 05, 2019.

---

## INTRODUCTION

Studies support the existence of a critical time early in infancy during which the genetically predisposed atopic infant is at higher risk for developing allergic sensitization [1]. Thus, dietary interventions in the first years of life have been analyzed for their effects on the prevalence of allergic disease including food allergy [2]. Both American and European allergy expert committee guidelines recommend that solid foods be introduced between four to six months of age in all infants [3-6]. Other organizations have also concluded that complementary foods may be safely introduced between four and six months of age [7-10], although many still recommend or prefer exclusive breastfeeding for the first six months of life [8,11-14]. Recommendations regarding when to introduce highly allergenic foods, particularly in high-risk infants, have shifted over time.

Introduction of highly allergenic foods is discussed here. The general approach to introduction of solid foods during infancy is reviewed in greater detail separately, as is use of formula in high-risk infants. (See "[Introducing solid foods and vitamin and mineral supplementation during infancy](#)" and "[Introducing formula to infants at risk for allergic disease](#)".)

Other aspects of the primary prevention of allergic disease are also discussed in greater detail separately. (See "[Primary prevention of allergic disease: Maternal diet in pregnancy and lactation](#)" and "[The impact of breastfeeding on the development of allergic disease](#)".)

---

## HIGHLY ALLERGENIC FOODS

While any food has the potential to cause allergy, certain foods are more common triggers of significant acute allergic reactions due to various factors. The most common food allergens in children in the United States and many other countries include cow's milk (CM), hen's egg, soy, wheat, peanut, tree nuts, and seafood (shellfish and fish). (See "[Pathogenesis of food allergy](#)", section on 'Factors influencing sensitization or tolerance' and "[History and physical examination in the patient with possible food allergy](#)", section on 'Common culprit foods' and "[Food allergy in children: Prevalence, natural history, and monitoring for resolution](#)", section on 'Prevalence of childhood food allergy' and "[Food allergens: Overview of clinical features and cross-reactivity](#)".)

---

## INTRODUCTION IN A HIGH-RISK POPULATION

Infants and young children with a family history of atopy are at high risk for developing allergic disease, and those with a personal history of atopy, particularly those with moderate-to-severe eczema, are also at increased risk of developing other atopic diseases including food allergies. The American Academy of Pediatrics (AAP) had previously suggested in 2000 that the introduction of certain highly allergenic foods be delayed further in high-risk children: cow's milk (CM) until age one year; eggs until age two years; and peanuts, tree nuts, and fish until age three years [15]. This recommendation was based upon early studies that suggested that delayed introduction of solid foods might help prevent some allergic diseases, particularly atopic dermatitis (AD) [16-18]. (See ["Introducing formula to infants at risk for allergic disease", section on 'Infants at high risk for developing allergy'.](#))

However, this advice was modified in 2008 with consensus that there was insufficient evidence to recommend any specific practices concerning the introduction of these foods after four to six months for the prevention of allergic disease in high-risk infants [6,7,19-21]. To the contrary, delayed introduction of solid foods may increase the risk of allergy [22,23], and early introduction of certain foods (eg, egg, peanut) between four to six months of age may decrease the risk of allergy to that specific food [24,25]. These findings suggest that the increased risk of peanut allergy seen in younger siblings of a child with peanut allergy, for example, is partly due to delayed introduction, in addition to an underlying genetic susceptibility [26]. Other risk factors, such as moderate to severe eczema, may also play a role in increasing this risk. Thus, we recommend **not** delaying the introduction of complementary foods into the diet of high-risk infants beyond what is generally recommended for all infants. The approach for specific highly allergenic foods is reviewed in greater detail below. (See ["Suggested approach"](#) below and ["Introducing solid foods and vitamin and mineral supplementation during infancy", section on 'When to initiate complementary foods'.](#))

In the first randomized trial examining early introduction of a highly allergenic food, 86 high-risk infants with moderate-to-severe eczema were randomly assigned to daily consumption of 1 teaspoon of pasteurized raw whole-egg powder (n = 49) or rice powder (n = 37) from four to eight months of age [27]. A medically supervised feeding of 2 teaspoons of whole cooked egg was performed at eight months of age, with egg introduced into the diet after that if the feeding was successful. An oral food challenge (OFC) with one-half of a whole pasteurized raw egg was performed at 12 months of age. Although these infants had no known prior direct ingestion of egg, 31 percent (15 of 49) in the egg group reacted to the egg powder, most on initial exposure, and did not continue ingestion. Of the infants who were still participating in the study at 12 months, 33 percent (14 of 42) in the egg group were diagnosed with immunoglobulin E (IgE)-mediated egg allergy (elevated egg-specific IgE and a positive raw egg challenge or history of an allergic reaction to egg) compared with 51 percent (18 of 35) in the control group, a nonsignificant difference. However, the trial was terminated early for safety reasons due to the large percentage of infants allergic at four months and significant allergic reactions at the entry OFC.

The Learning Early about Peanut Allergy (LEAP) trial was the first randomized trial to show benefit of early introduction of a major food allergen, with earlier introduction of peanut at 4 to 11 months of age associated with a decreased risk of developing peanut allergy [22]. In this study, 640 children aged 4 to <11 months (median 7.8 months) with severe eczema and/or egg allergy and a skin prick test (SPT) wheal of  $\leq 4$  mm to peanut were randomly assigned to either consumption or avoidance of peanut until 60 months of age. Randomization occurred after participants were stratified into one of two groups: sensitized (SPT wheal 1 to 4 mm; n = 98) or not sensitized (no wheal on SPT; n = 542) to peanut. Infants assigned to the consumption group first underwent an OFC to peanut to confirm lack of peanut allergy. Those with a positive OFC (one in the nonsensitized group and six in the sensitized group) were told to avoid peanut. Those children in the consumption group who had a negative OFC were fed at least 6 grams of peanut protein per week (consumed in three or more meals per week). Adherence to consumption or avoidance was assessed using a validated

food frequency questionnaire and measurements of peanut protein in home dust samples. The presence of peanut allergy was then determined at 60 months of age by OFC.

Among the 530 children evaluated for the primary outcome in the nonsensitized group at 60 months of age, 13.7 percent of the avoidance group and 1.9 percent of the consumption group were allergic to peanut (absolute risk difference 11.8, 95% CI 3.4-20.3), which is an 86 percent relative risk reduction. All 98 children in the sensitized group were evaluated at 60 months; 35.3 percent of the avoidance group and 10.6 percent of the consumption group were allergic to peanuts (absolute risk difference 24.7, 95% CI 4.9-43.3), which is a 70 percent relative risk reduction. The retention rate of the study was 98.4 percent, and adherence to the assigned treatment was also high.

In the follow-up (LEAP-On) study that included 550 of the participants from the primary trial, both groups were asked to avoid peanut for 12 months until 72 months of age [23]. Compliance with avoidance was 90.4 percent in the original avoidance group and 69.3 percent in the original consumption group. Peanut allergy continued to be significantly more common in the peanut avoidance group (18.6 percent, 52 of 280 children) than the early peanut consumption group (4.8 percent, 13 of 270 children). Three children in the consumption group who were previously tolerant developed peanut allergy during the 12-month avoidance period. In the group that continued avoidance, three children developed peanut allergy, and four had resolution of the allergy during the follow-up period. Development of peanut allergy in previously tolerant children who cease consumption is consistent with data on peanut allergy recurrence in patients who had outgrown their peanut allergy but did not go on to eat it regularly after a successful OFC. (See "[Food allergy in children: Prevalence, natural history, and monitoring for resolution](#)", [section on 'Recurrence'](#).)

In the Prevention of Egg Allergy with Tiny Amount Intake (PETIT) trial, 147 infants with eczema and no prior direct ingestion of egg or history of an allergic reaction to egg were randomly assigned to daily consumption of heated egg powder (50 mg daily from 6 to 9 months of age and 250 mg daily thereafter until 12 months of age) or placebo along with aggressive treatment of eczema [28]. Participants underwent an OFC to 7 grams of heated whole-egg powder equivalent to 32 grams of boiled, whole hen's egg at 12 months of age. The trial was stopped early due to benefit. The 121 infants who performed the egg OFC at 12 months were included in the intent-to-treat analysis. Five of 60 infants (8 percent) in the treatment group had egg allergy confirmed by OFC compared with 23 of 61 (38 percent) in the placebo group (risk ratio 0.22, 95% CI 0.09-0.54). Limitations of this study include stopping early for benefit, which tends to overestimate treatment effect, especially if the event rate is low, as in this case. In addition, baseline eczema was more severe in the placebo group compared with the treatment group based upon the mean SCORAD score (42.0 versus 27.5, respectively), and the mean egg-specific IgE was higher in the placebo group as well (4.46 versus 0.73, respectively).

Two other trials of egg introduction beginning at four to six months of age were performed in infants at risk for food allergies based upon family history of atopic disease. The first of these, the Starting Time of Egg Protein (STEP) trial, enrolled 820 infants of atopic mothers with allergic disease and a positive environmental SPT who were aged four to six months and did not have a history of eczema or known prior direct ingestion of egg [29]. Testing for egg sensitization was not performed prior to starting the treatment phase of the study. Infants were randomly assigned at four to six months of age to daily ingestion of 900 mg of pasteurized, raw whole-egg protein or placebo daily until 10 months of age. Cooked egg was introduced into the diet in both groups at 10 months of age, and a raw egg OFC was performed at 12 months of age. There was no difference in the percent of infants who reacted to the egg OFC at 12 months of age (7 percent in the treatment group compared with 10.3 percent in the placebo group). However, a higher percentage of infants stopped taking the study powder due to confirmed allergic reactions compared with the placebo group (6.1 versus 1.5 percent, respectively), suggesting that some of these infants were already allergic to egg.

The second trial, the Beating Egg Allergy Trial (BEAT), enrolled 319 infants who were SPT negative to egg but had a first-degree relative with atopic disease [30]. These infants were randomly assigned at four months of age to 350 mg pasteurized, raw whole-egg protein or placebo daily and were treated until eight months of age, at which time egg was introduced into the diet. At 12 months of age, egg sensitization was significantly lower in the treatment group (13 of 122, 10.7 percent) compared with placebo (25 of 122, 20.5 percent). However, there was a significant loss to follow-up, with only 254 infants having at least partial assessment at 12 months of age. In addition, 14 of the infants in the treatment group were lost to follow-up because they reacted to the egg powder at introduction. Presumably, there would have been no difference in the rates of sensitization had those children been evaluated at 12 months. Furthermore, there was no difference in the number of infants with probable egg allergy, a more clinically relevant outcome than sensitization, at 12 months of age.

---

## INTRODUCTION IN THE GENERAL POPULATION

A number of studies, most of which were performed in population-based, prospective birth cohorts, have not supported delaying the introduction of solid foods beyond four to six months for the prevention of allergic disease [31-42]. Similar to the high-risk population, delayed introduction of solid foods may increase the risk of allergy [32], as may restricting the diversity of complementary foods introduced in the first year of life [43], and early introduction between four to six months of age may decrease the risk of allergy [24]. Thus, complementary foods, including highly allergenic foods, may be introduced into the diet of "standard or low" risk children any time after four to six months of age. However, it is still possible that some children in this lower-risk stratum will have allergic reactions with introduction since some infants are already allergic to foods such as egg by four to six months of age [44]. Introduction of solids during infancy in the general population is reviewed in detail separately. (See "[Introducing solid foods and vitamin and mineral supplementation during infancy](#)".)

One prospective study examined the feeding history of 13,019 infants and evaluated probable adverse reactions to cow's milk (CM) by history and physical exam, milk skin prick testing (SPT), and oral food challenge (OFC) [38]. The cumulative incidence for IgE-mediated CM allergy was 0.5 percent. CM protein was introduced into the diet significantly later in the allergic group than the tolerant group (mean age at introduction 116 versus 62 days of life, respectively). Early exposure to CM formula (introduction within the first 14 days of life) was associated with a lower rate of CM allergy than those started between 105 to 194 days of age (0.05 versus 1.75 percent, respectively).

A population-based, cross-sectional study examined the prevalence of egg allergy at 12 months of age (based upon positive egg-specific IgE testing followed by confirmation of allergy with an OFC to egg or a prior history of a reaction) in 2589 infants [37]. Later introduction of egg was associated with an increased risk of egg allergy (5.6, 7.8, 10.1, and 27.6 percent of infants at 12 months of age had egg allergy with introduction at 4 to 6, 7 to 9, 10 to 12, and >12 months of age, respectively), with significant differences seen with introduction at 10 to 12 and >12 months compared with 4 to 6 months (adjusted odds ratio [OR] 1.6, 95% CI 1.0-2.6 and OR 3.4, 95% CI 1.8-6.5, respectively). A lower risk of egg allergy was seen in infants who were given lightly cooked egg first rather than egg in baked goods first (OR 0.2, 95% CI 0.06-0.71).

The Enquiring about Tolerance (EAT) trial examined whether early introduction of six highly allergenic foods (hen's egg, CM, sesame, whitefish, peanut, and wheat) beginning at three months of age versus more standard introduction at six months in breastfed infants protected against the development of food allergy [45]. Infants were recruited from the general population and, as such, were technically not at increased risk for food allergy. However, the group was self-enriched for atopy. The 1303 exclusively breastfed infants were randomly assigned to introduction of highly allergenic

foods between three and six months of age or continued exclusive breastfeeding until six months of age. Milk in the form of yogurt was always the first food introduced, and wheat was always the last, with the other foods introduced in random order.

There was no difference in the prevalence of food allergy at one and three years of age between the two groups in the intention-to-treat (ITT) analysis. However, the prevalence of any food allergy, and peanut and egg allergy in particular, was lower in the early-introduction group compared with the standard-introduction group in the per-protocol analysis (2.4 versus 7.3 percent, respectively, for any food allergy). Earlier feeding did not seem to reduce rates of breastfeeding. Adherence to the food regimen was problematic, with 43 percent of the participants in the early-introduction group adhering to the protocol (ie, able to consume all six allergenic foods in the required amount and by the required time). Although the most conservative interpretation of the data from the ITT analysis did not show that this regimen reduces the risk of food allergies. This study does support the recommendation to not withhold allergenic foods, but rather to introduce them as early as four to six months of age. The data also suggest that earlier introduction may be beneficial and may not be harmful nor effect rates of concomitant breastfeeding.

The Hen's Egg Allergy Prevention (HEAP) trial randomly assigned 383 infants four to six months of age who were recruited from the general population and had hen's egg-specific IgE  $<0.35$  kU<sub>A</sub>/L to increasing doses of raw, pasteurized egg white three times per week (0.83 g/dose during week 1, 1.67 g/dose during week 2, and 2.5 g/dose week three until 12 months of age) or placebo [44]. There was no difference between the two groups with regard to egg sensitization or egg allergy confirmed by OFC. The adjusted prevalence of egg allergy in the 406 infants screened for the study was 5.3 percent. The authors noted a high rate of anaphylaxis at entry egg introduction, which may be partially attributed to the raw pasteurized form and the relatively larger amounts of egg protein consumed compared with other early egg introduction randomized, controlled trials.

A meta-analysis that combined data on trials in both normal- and high-risk infants found that early introduction of egg or peanut at four to six months of age was associated with a decreased risk of allergy to that food specifically (egg allergy risk ratio 0.56, 95% CI 0.36-0.87 and peanut allergy risk ratio 0.29, 95% CI 0.11-0.74) [24]. Although this meta-analysis found "moderate certainty" evidence for a protective association for early introduction of peanut and egg, there are notable limitations. The results for peanut were based on two randomized, controlled trials: Learning Early about Peanut Allergy (LEAP), which again was performed in a narrowly selected, high-risk group defined by severe eczema and/or egg allergy, and EAT, which recruited subjects from the general population. While LEAP showed significant risk reduction for preventing peanut allergy in ITT analysis, EAT did not, only showing risk reduction for prevention of peanut allergy in the weaker per protocol analysis. The data for egg included five trials, four of which failed to show a protective effect for early egg introduction that used a raw pasteurized egg product, while the remaining one trial (PETIT, published only in abstract form at the time of the meta-analysis) that showed benefit of early egg introduction used heated egg, which is less allergenic than raw egg. Inclusion of the data from this abstract was strong enough to influence the overall protective effect for early egg introduction. However, more research is needed to better understand the different forms of egg and their effects on egg allergy prevention.

---

## SUGGESTED APPROACH

There are few studies examining the safest way to introduce highly allergenic foods in high-risk children. For some foods, such as peanut, the majority of reactions occur in response to what is often believed to be the initial ingestion [46]. The Learning Early about Peanut Allergy (LEAP) and follow-up LEAP-On trials reviewed above focused on a specific high-risk group and a specific allergen, with skin prick tests (SPTs) and oral food challenges (OFCs) to peanut performed

prior to introduction [22,23]. While the results from this trial may not be broadly generalizable to other foods, to infants at high risk for allergy for other reasons, or to infants at low risk for allergy, the findings do suggest that past recommendations to substantially delay introduction of highly allergenic foods were appropriately rescinded. Formal guidelines with respect to introduction of highly allergenic complementary foods are under development. An interim guideline addressing introduction of peanut based upon the LEAP trial and other studies was published in 2017 [47]. Our suggested approach outlined below is consistent with these guidelines.

We counsel parents to introduce highly allergenic foods (eg, cow's milk [CM], hen's egg, peanut, tree nuts, fish, and shellfish) in the following manner in infants who are at risk based upon family history but who have not had any significant prior allergic reactions to a food or difficult-to-control moderate-to-severe atopic dermatitis (AD) [6]:

- First, the child should be at least four months of age and have shown developmental readiness to consume complementary foods. (See ["Introducing solid foods and vitamin and mineral supplementation during infancy", section on 'Developmental skills'.](#))
- In addition, the child should have tolerated a few of the more typical, initial complementary foods (such as cereals, fruits, and vegetables).
- If these two criteria are met, then the child can be given an initial taste of one of these foods at home (rather than at daycare or at a restaurant), with an oral antihistamine available.
- If there is no apparent reaction, the food can be introduced in gradually increasing amounts.

However, an allergy evaluation, including a detailed history and possible testing, before introduction of highly allergenic foods is a reasonable option in patients with the following histories (see ["History and physical examination in the patient with possible food allergy"](#) and ["Diagnostic evaluation of food allergy"](#)):

- Recalcitrant, moderate-to-severe AD despite optimal management.
- Signs or symptoms of an immediate allergic reaction while breastfeeding or with the introduction of any food, especially one of the highly allergenic foods.

For children with the above features, evaluation for introducing peanut may follow the consensus recommendations based upon LEAP and other studies [47,48]. A suggested approach is shown in the algorithm ([algorithm 1](#)). Several cases of food protein-induced enterocolitis (FPIES) triggered by peanut have been reported in high-risk infants undergoing early feeding of peanut [49]. (See ["Food protein-induced enterocolitis syndrome \(FPIES\)"](#).)

The general guidelines for introducing solid foods to an infant's diet are discussed in detail elsewhere. (See ["Introducing solid foods and vitamin and mineral supplementation during infancy", section on 'Optimal timing'.](#))

---

## SOCIETY GUIDELINE LINKS

Links to society and government-sponsored guidelines from selected countries and regions around the world are provided separately. (See ["Society guideline links: Food allergy"](#).)

---

## INFORMATION FOR PATIENTS

UpToDate offers two types of patient education materials, "The Basics" and "Beyond the Basics." The Basics patient education pieces are written in plain language, at the 5<sup>th</sup> to 6<sup>th</sup> grade reading level, and they answer the four or five key questions a patient might have about a given condition. These articles are best for patients who want a general overview and who prefer short, easy-to-read materials. Beyond the Basics patient education pieces are longer, more sophisticated, and more detailed. These articles are written at the 10<sup>th</sup> to 12<sup>th</sup> grade reading level and are best for patients who want in-depth information and are comfortable with some medical jargon.

Here are the patient education articles that are relevant to this topic. We encourage you to print or e-mail these topics to your patients. (You can also locate patient education articles on a variety of subjects by searching on "patient info" and the keyword(s) of interest.)

- Beyond the Basics topic (see ["Patient education: Starting solid foods during infancy\\_\(Beyond the Basics\)"](#))

---

## SUMMARY AND RECOMMENDATIONS

- Human milk is the optimal source of nutrition for all term infants during the first four to six months of life, regardless of their risk for allergic disease. (See ["Infant benefits of breastfeeding"](#).)
- The most common highly allergenic foods include cow's milk (CM), hen's egg, soy, wheat, peanut, tree nuts, shellfish, and fish. (See ["Highly allergenic foods"](#) above.)
- Previous guidelines recommended delayed introduction of highly allergenic solid foods for the purpose of preventing allergic disease in high-risk infants. However, evidence suggests that this practice may increase rather than decrease the incidence of food allergies. We recommend early introduction of highly allergenic solid foods in high-risk infants ([Grade 1B](#)). These infants should be at least four months of age, be developmentally ready, and have tolerated a few less allergenic complementary foods, such as rice cereal and pureed fruits or vegetables. (See ["Introduction in a high-risk population"](#) above and ["Suggested approach"](#) above.)
- Highly allergenic foods can be gradually and carefully introduced to asymptomatic high-risk infants without prior testing. The one exception is liquid, whole CM, which should be avoided in all infants less than one year of age for reasons unrelated to allergy. (CM formula and other CM products such as those in baked goods, cheese, and yogurt are not restricted prior to age one year.)
- However, an allergy evaluation, including a detailed history and possible testing, before introduction of highly allergenic foods is a reasonable option if a patient has had an immediate allergic reaction to a food or breast milk or has difficult-to-control moderate-to-severe atopic dermatitis (AD) despite optimal management. (See ["Suggested approach"](#) above.)
- Complementary foods, including highly allergenic foods, may be introduced into the diet of low-risk children at any time after four to six months of age. All solid food should be introduced in a form that does not present a choking hazard. Once introduced, highly allergenic foods should be consumed on a regular basis. (See ["Introduction in the general population"](#) above.)

Use of UpToDate is subject to the [Subscription and License Agreement](#).

## REFERENCES

1. [Zeiger RS. Food allergen avoidance in the prevention of food allergy in infants and children. Pediatrics 2003; 111:1662.](#)
2. [Pali-Schöll I, Renz H, Jensen-Jarolim E. Update on allergies in pregnancy, lactation, and early childhood. J Allergy Clin Immunol 2009; 123:1012.](#)
3. [Greer FR, Sicherer SH, Burks AW, et al. The Effects of Early Nutritional Interventions on the Development of Atopic Disease in Infants and Children: The Role of Maternal Dietary Restriction, Breastfeeding, Hydrolyzed Formulas, and Timing of Introduction of Allergenic Complementary Foods. Pediatrics 2019; 143.](#)
4. [Muraro A, Dreborg S, Halken S, et al. Dietary prevention of allergic diseases in infants and small children. Part III: Critical review of published peer-reviewed observational and interventional studies and final recommendations. Pediatr Allergy Immunol 2004; 15:291.](#)
5. [Grimshaw KE, Allen K, Edwards CA, et al. Infant feeding and allergy prevention: a review of current knowledge and recommendations. A EuroPrevall state of the art paper. Allergy 2009; 64:1407.](#)
6. [Fleischer DM, Spergel JM, Assa'ad AH, Pongratic JA. Primary prevention of allergic disease through nutritional interventions. J Allergy Clin Immunol Pract 2013; 1:29.](#)
7. [Agostoni C, Decsi T, Fewtrell M, et al. Complementary feeding: a commentary by the ESPGHAN Committee on Nutrition. J Pediatr Gastroenterol Nutr 2008; 46:99.](#)
8. [Section on Breastfeeding. Breastfeeding and the use of human milk. Pediatrics 2012; 129:e827.](#)
9. [European Food, Safety Authority. Scientific opinion on the appropriate age for introduction of complementary feeding of infants. EFSA Journal 2009; 7:1423.](#)
10. [Netting MJ, Campbell DE, Koplin JJ, et al. An Australian Consensus on Infant Feeding Guidelines to Prevent Food Allergy: Outcomes From the Australian Infant Feeding Summit. J Allergy Clin Immunol Pract 2017; 5:1617.](#)
11. World Health Organization. Global Strategy for Infant and Young Child Feeding (2003). [www.who.int/nutrition/publications/infantfeeding/en/index.html](http://www.who.int/nutrition/publications/infantfeeding/en/index.html) (Accessed on April 10, 2009).
12. [Committee on Health Care for Underserved Women, American College of Obstetricians and Gynecologists. ACOG Committee Opinion No. 361: Breastfeeding: maternal and infant aspects. Obstet Gynecol 2007; 109:479.](#)
13. [Boland M. Exclusive breastfeeding should continue to six months. Paediatr Child Health 2005; 10:148.](#)
14. [US Preventive Services Task Force, Bibbins-Domingo K, Grossman DC, et al. Primary Care Interventions to Support Breastfeeding: US Preventive Services Task Force Recommendation Statement. JAMA 2016; 316:1688.](#)
15. [American Academy of Pediatrics. Committee on Nutrition. Hypoallergenic infant formulas. Pediatrics 2000; 106:346.](#)
16. [Fergusson DM, Horwood LJ, Beautrais AL, et al. Eczema and infant diet. Clin Allergy 1981; 11:325.](#)



17. [Fergusson DM, Horwood LJ, Shannon FT. Early solid feeding and recurrent childhood eczema: a 10-year longitudinal study. Pediatrics 1990; 86:541.](#)
18. [Fergusson DM, Horwood LJ, Shannon FT. Asthma and infant diet. Arch Dis Child 1983; 58:48.](#)
19. [Sicherer SH, Burks AW. Maternal and infant diets for prevention of allergic diseases: understanding menu changes in 2008. J Allergy Clin Immunol 2008; 122:29.](#)
20. [Greer FR, Sicherer SH, Burks AW, et al. Effects of early nutritional interventions on the development of atopic disease in infants and children: the role of maternal dietary restriction, breastfeeding, timing of introduction of complementary foods, and hydrolyzed formulas. Pediatrics 2008; 121:183.](#)
21. [Høst A, Halken S, Muraro A, et al. Dietary prevention of allergic diseases in infants and small children. Pediatr Allergy Immunol 2008; 19:1.](#)
22. [Du Toit G, Roberts G, Sayre PH, et al. Randomized trial of peanut consumption in infants at risk for peanut allergy. N Engl J Med 2015; 372:803.](#)
23. [Du Toit G, Sayre PH, Roberts G, et al. Effect of Avoidance on Peanut Allergy after Early Peanut Consumption. N Engl J Med 2016; 374:1435.](#)
24. [Ierodiakonou D, Garcia-Larsen V, Logan A, et al. Timing of Allergenic Food Introduction to the Infant Diet and Risk of Allergic or Autoimmune Disease: A Systematic Review and Meta-analysis. JAMA 2016; 316:1181.](#)
25. [Sicherer SH, Allen K, Lack G, et al. Critical Issues in Food Allergy: A National Academies Consensus Report. Pediatrics 2017; 140.](#)
26. [Abrams EM, Chan ES, Sicherer SH. Should Younger Siblings of Peanut Allergic Children Be Screened for Peanut Allergy? J Allergy Clin Immunol Pract 2018; 6:414.](#)
27. [Palmer DJ, Metcalfe J, Makrides M, et al. Early regular egg exposure in infants with eczema: A randomized controlled trial. J Allergy Clin Immunol 2013; 132:387.](#)
28. [Natsume O, Kabashima S, Nakazato J, et al. Two-step egg introduction for prevention of egg allergy in high-risk infants with eczema \(PETIT\): a randomised, double-blind, placebo-controlled trial. Lancet 2017; 389:276.](#)
29. [Palmer DJ, Sullivan TR, Gold MS, et al. Randomized controlled trial of early regular egg intake to prevent egg allergy. J Allergy Clin Immunol 2017; 139:1600.](#)
30. [Wei-Liang Tan J, Valerio C, Barnes EH, et al. A randomized trial of egg introduction from 4 months of age in infants at risk for egg allergy. J Allergy Clin Immunol 2017; 139:1621.](#)
31. [Du Toit G, Katz Y, Sasieni P, et al. Early consumption of peanuts in infancy is associated with a low prevalence of peanut allergy. J Allergy Clin Immunol 2008; 122:984.](#)
32. [Zutavern A, von Mutius E, Harris J, et al. The introduction of solids in relation to asthma and eczema. Arch Dis Child 2004; 89:303.](#)
33. [Zutavern A, Brockow I, Schaaf B, et al. Timing of solid food introduction in relation to eczema, asthma, allergic rhinitis, and food and inhalant sensitization at the age of 6 years: results from the prospective birth cohort study.](#)

[LISA. Pediatrics 2008; 121:e44.](#)


34. [Snijders BE, Thijs C, van Ree R, van den Brandt PA. Age at first introduction of cow milk products and other food products in relation to infant atopic manifestations in the first 2 years of life: the KOALA Birth Cohort Study. Pediatrics 2008; 122:e115.](#)
35. [Poole JA, Barriga K, Leung DY, et al. Timing of initial exposure to cereal grains and the risk of wheat allergy. Pediatrics 2006; 117:2175.](#)
36. [Nwaru BI, Erkkola M, Ahonen S, et al. Age at the introduction of solid foods during the first year and allergic sensitization at age 5 years. Pediatrics 2010; 125:50.](#)
37. [Koplin JJ, Osborne NJ, Wake M, et al. Can early introduction of egg prevent egg allergy in infants? A population-based study. J Allergy Clin Immunol 2010; 126:807.](#)
38. [Katz Y, Rajuan N, Goldberg MR, et al. Early exposure to cow's milk protein is protective against IgE-mediated cow's milk protein allergy. J Allergy Clin Immunol 2010; 126:77.](#)
39. [Tromp II, Kiefte-de Jong JC, Lebon A, et al. The introduction of allergenic foods and the development of reported wheezing and eczema in childhood: the Generation R study. Arch Pediatr Adolesc Med 2011; 165:933.](#)
40. [Joseph CL, Ownby DR, Havstad SL, et al. Early complementary feeding and risk of food sensitization in a birth cohort. J Allergy Clin Immunol 2011; 127:1203.](#)
41. [Grimshaw KE, Maskell J, Oliver EM, et al. Introduction of complementary foods and the relationship to food allergy. Pediatrics 2013; 132:e1529.](#)
42. [Luccioli S, Zhang Y, Verrill L, et al. Infant feeding practices and reported food allergies at 6 years of age. Pediatrics 2014; 134 Suppl 1:S21.](#)
43. [Roduit C, Frei R, Depner M, et al. Increased food diversity in the first year of life is inversely associated with allergic diseases. J Allergy Clin Immunol 2014; 133:1056.](#)
44. [Bellach J, Schwarz V, Ahrens B, et al. Randomized placebo-controlled trial of hen's egg consumption for primary prevention in infants. J Allergy Clin Immunol 2017; 139:1591.](#)
45. [Perkin MR, Logan K, Tseng A, et al. Randomized Trial of Introduction of Allergenic Foods in Breast-Fed Infants. N Engl J Med 2016; 374:1733.](#)
46. [Sicherer SH, Burks AW, Sampson HA. Clinical features of acute allergic reactions to peanut and tree nuts in children. Pediatrics 1998; 102:e6.](#)
47. [Togias A, Cooper SF, Acebal ML, et al. Addendum Guidelines for the Prevention of Peanut Allergy in the United States: Report of the National Institute of Allergy and Infectious Diseases-Sponsored Expert Panel. Pediatr Dermatol 2017; 34:e1.](#)
48. [Peters RL, Allen KJ, Dharmage SC, et al. Skin prick test responses and allergen-specific IgE levels as predictors of peanut, egg, and sesame allergy in infants. J Allergy Clin Immunol 2013; 132:874.](#)

49. [Robbins KA, Ackerman OR, Carter CA, et al. Food protein-induced enterocolitis syndrome to peanut with early introduction: a clinical dilemma. J Allergy Clin Immunol Pract 2018; 6:664.](#)

Topic 107525 Version 19.0

## GRAPHICS

### Introduction of peanut into the diet during infancy/early childhood



IgE: immunoglobulin E; PN-IgE: peanut-specific immunoglobulin E.

\* Alternatively, the allergy specialist may choose to perform PN-IgE testing, or both skin prick and PN-IgE testing, for the initial evaluation.

¶ Prior to introduction of highly allergenic foods, the infant/child should:

- Be at least 4 months of age.
- Show developmental readiness to consume complementary foods.
- Have tolerated a few of the more typical, initial complementary foods (eg, cereals, fruits, vegetables).

Δ These skin prick test cutoffs for peanut are best estimates based upon limited data from 2 non-US study cohorts (Learning Early About Peanut Allergy [LEAP] and HealthNuts). A number of factors can affect the size of the skin test reaction, including test site chosen (back versus forearm), testing device used, and age of the patient. It is important that clinicians performing in-office feedings or graded challenges have the appropriate training and equipment to manage potential severe allergic reactions regardless of the test results.

Graphic 111453 Version 4.0

## Contributor Disclosures

**David M Fleischer, MD** Grant/Research/Clinical Trial Support: DBV Technologies [Food allergies (Epicutaneous immunotherapy patch)]; Aimmune Therapeutics [Food allergies (oral immunotherapy)]. Consultant/Advisory Boards: Food Allergy & Anaphylaxis Connection Team (FAACT) [Food allergy]; FARE [Food allergy]; National Peanut Board [Food allergy]; DBV Technologies [Food allergy]; Intromune Therapeutics [Food allergy]; INSYS Therapeutics [Food allergy]; AllerGenis [Food allergy]; DOTS Technology [Food allergy]. Speaker's Bureau: Abbott [Food allergy]; Nutricia [Food allergy]. **Scott H Sicherer, MD, FAACAP** Grant/Research/Clinical Trial Support: HAL Allergy [Food allergy]; Food Allergy Research and Education [Food allergy]; NIAID [Food allergy clinical trials]. Other Financial Interest: Johns Hopkins University Press [Royalties]. **Elizabeth TePas, MD, MS** Nothing to disclose

Contributor disclosures are reviewed for conflicts of interest by the editorial group. When found, these are addressed by vetting through a multi-level review process, and through requirements for references to be provided to support the content. Appropriately referenced content is required of all authors and must conform to UpToDate standards of evidence.

[Conflict of interest policy](#).