

Juglandaceae—Walnut family

Carya Nutt.

hickory

Franklin T. Bonner

Dr. Bonner is a scientist emeritus at the USDA Forest Service's Southern Research Station, Mississippi State, Mississippi

Growth habit, occurrence, and use. Of the dozen or so species of hickories native to the United States, 9 are valuable for timber and the food they provide for wildlife (table 1). All are deciduous trees. Pecan and its many horticultural varieties and hybrids are widely cultivated for nuts in large plantations in the southern and southwestern United States, as well as in many other countries. The first known selections were made in 1846, and many cultivars were available by the late 19th century (Madden and Malstrom 1975). Budding and grafting have been the primary means of improvement, but new provenance studies (Grauke and

others 1990) and advanced research on the reproductive biology and genetics of pecan (Graves and others 1989; McCarthy and Quinn 1990; Yates and Reilly 1990; Yates and Sparks 1990) demonstrate the promise for future improvements in nut production and disease resistance. Shellbark and shagbark hickories have also been planted for nut production.

Flowering and Fruiting. Hickories are monoecious and flower in the spring (table 2). The staminate catkins develop from axils of leaves of the previous season or from inner scales of the terminal buds at the base of the current

Table 1—Carya, hickory: nomenclature and occurrence

Scientific name & synonym(s)	Common name	Occurrence
C. alba (L.) Nutt. ex Ell. <i>C. tomentosa</i> (Lam. ex Poir.) Nutt. <i>Hicoria tomentosa</i> (Lam. ex Poir.) Raf.	mockernut hickory , bullnut, white hickory, whiteheart hickory, hognut, mockernut	S New Hampshire to S Michigan, S to E Texas & N Florida Valley to Illinois
C. aquatica (Michx. f.) Nutt. <i>Hicoria aquatica</i> (Michx. f.) Britt.	water hickory , bitter pecan swamp hickory	Coastal plain from Virginia to S Florida & E Texas; N in Mississippi Valley to Illinois
C. cordiformis (Wangenh.) K. Koch. <i>Hicoria cordiformis</i> (Wagenh.) Britt.	bitternut hickory , bitternut, swamp hickory, pignut	New Hampshire to Minnesota, S to E Texas & Georgia
C. glabra (P. Mill.) Sweet <i>Hicoria glabra</i> (Mill.) Britt. <i>C. microcarpa</i> (Nutt.) Britt.	pignut hickory , sweet pignut, pignut, swamp hickory	New Hampshire to NE Kansas, S to Arkansas & NW Florida
C. illinoensis (Wangenh.) K. Koch <i>Hicoria pecan</i> (Marsh.) Britt. <i>C. oliviformis</i> (Michx. f.) Nutt. <i>C. pecan</i> (Marsh.) Engl & Graebn.	pecan , sweet pecan, <i>nuez encarcelada</i>	S Indiana to SE Iowa; S to Texas & E to Mississippi & W Tennessee; local to Ohio, Kentucky, & Alabama
C. laciniosa (Michx. f.) G. Don <i>Hicoria laciniosa</i> (Michx. f.) Sarg.	shellbark hickory , bigleaf shagbark hickory, big shellbark, kingnut, bottom shellbark, big shagbark hickory	Ohio & Mississippi Valleys; W New York to E Kansas, E to Georgia & Virginia; local in Louisiana, Alabama, & Virginia
C. myristiciformis (Michx. f.) Nutt. <i>Hicoria myristicaeformis</i> (Michx. f.) Britt.	nutmeg hickory , bitter water hickory, swamp hickory	Mississippi W to SE Oklahoma, S to E Texas & Louisiana; also E South Carolina & central Alabama
C. ovata (P. Mill.) K. Koch <i>Hicoria alba</i> Britt. p.p.; <i>H. ovata</i> (P. Mill.) Britt.	shagbark hickory , scalybark hickory, shagbark, shellbark hickory	Maine to SE Minnesota, S to E Texas & Georgia
C. pallida (Ashe) Engl. & Graebn. <i>Hicoria pallida</i> Ashe	sand hickory , pale hickory, pallid hickory	New Jersey & Illinois, S to Florida & SE Louisiana

Sources: Little (1979), Sargent (1965).

Table 2—*Carya*, hickory: phenology of flowering and fruiting

Species	Flowering	Fruit ripening	Seed dispersal
<i>C. alba</i>	Apr–May	Sept–Oct	Sept–Oct
<i>C. aquatica</i>	Mar–May	Sept–Nov	Oct–Dec
<i>C. cordiformis</i>	Apr–May	Sept–Oct	Sept–Dec
<i>C. glabra</i>	Apr–May	Sept–Oct	Sept–Oct
<i>C. illinoensis</i>	Mar–May	Sept–Oct	Sept–Oct
<i>C. laciniosa</i>	Apr–June	Sept–Nov	Sept–Oct
<i>C. myristiciformis</i>	Apr–May	Sept–Oct	Sept–Oct
<i>C. ovata</i>	Apr–June	Sept–Oct	Sept–Oct
<i>C. pallida</i>	Mar–Apr	Sept–Oct	Sept–Oct

Source: Bonner and Maisenhelder (1974).

growth. The pistillate flowers appear in short spikes on peduncles terminating in shoots of the current year. Hickory fruits are ovoid, globose, or pear-shaped nuts enclosed in husks developed from the floral involucre (figure 1). Husks are green prior to maturity and then turn brown to brownish black as they ripen (Bonner and Maisenhelder 1974). The husks become dry at maturity in the fall (table 2) and split away from the nut into 4 valves along sutures. Husks of mockernut, nutmeg, shagbark, and shellbark hickories, as well as those of pecan, split to the base at maturity, usually

Figure 1—*Carya*, hickory: nuts with husks attached and removed (the size and shape of individual nuts varies greatly within a species and may differ from the examples shown here); *C. aquatica*, water hickory (**first row left**) and *C. cordiformis*, bitternut hickory (**first row right**); *C. glabra*, pignut hickory (**second row left**) and *C. myristiciformis*, nutmeg hickory (**second row right**); *C. illinoensis*, pecan (**third row left**) and *C. laciniosa*, shellbark hickory (**third row right**); *C. ovata*, shagbark hickory (**fourth row left**) and *C. alba*, mockernut hickory (**fourth row right**).



releasing the nuts. Husks of pignut, bitternut, sand, and water hickories split only to the middle or slightly beyond and generally cling to the nuts. The nut is 4-celled at the base and 2-celled at the apex. The edible portion of the embryonic plant is mainly cotyledonary tissue (figure 2) and has a very high lipid content (Bonner 1971; Bonner 1974; Short and Epps 1976).

Collection, extraction, and storage. Hickory nuts can be collected from the ground after natural seedfall or after shaking the trees or flailing the limbs. Persistent husks may be removed by hand, by trampling, or by running the fruits through a macerator or a corn sheller. Several studies have shown that the larger nuts of pecan make larger seedlings (Adams and Thielges 1977; Herrera and Martinez 1983), so sizing of nuts may be beneficial. Shagbark and shellbark hickory trees have been known to produce 0.5 to 0.75 hl (1½ to 2 bu) and 0.75 to 1.1 hl (2 to 3 bu) of nuts, respectively (Bonner and Maisenhelder 1974). Good crops of all species are produced at intervals of 1 to 3 years (table 3). Some typical yield data are presented in table 4.


Figure 2—*Carya ovata*, shagbark hickory: longitudinal section through the embryo of a nut with husk removed.

Table 3—*Carya*, hickory: height, seed-bearing age, seedcrop frequency, and year first cultivated

Species	Height at maturity (m)	Year first cultivated	Minimum seed-bearing age (yrs)	Years between seedcrops
<i>C. alba</i>	30	1766	25	2–3
<i>C. aquatica</i>	30	1800	20	1–2
<i>C. cordiformis</i>	15–30	1689	30	3–5
<i>C. glabra</i>	24–27	1750	30	1–2
<i>C. illinoensis</i>	34–43	1766	10–20	1–2
<i>C. laciniosa</i>	37	1800	40	1–2
<i>C. myristiciformis</i>	24–30	—	30	2–3
<i>C. ovata</i>	21–30	1911	40	1–3
<i>C. pallida</i>	12–30	—	—	2–3

Source: Bonner and Maisenhelder (1974).

Table 4—*Carya*, hickory: seed data

Species	Place collected	Fruits/vol		Seed wt/fruit vol		Cleaned seeds/weight			
		/hl	/bu	kg/hl	lb/bu	Range		Average	
						/kg	/lb	/kg	/lb
<i>C. alba</i>	—	—	—	—	—	75–249	34–113	200	90
	Mississippi	5,040	1,776	57	44	71–106	32–48	79	36
<i>C. aquatica</i>	Mississippi	—	—	—	—	305–419	138–140	360	164
<i>C. cordiformis</i>	—	—	—	51	40	275–408	125–185	344	156
<i>C. glabra</i>	—	—	—	51	40	386–496	175–225	441	200
	Mississippi	10,100	3,552	—	—	—	—	143	65
<i>C. illinoensis</i>	—	—	—	—	—	121–353	55–160	220	100
	Mississippi	20,800	7,330	—	—	333–384	151–174	357	162
	Texas	—	—	—	—	—	—	311	141
<i>C. laciniosa</i>	—	—	—	—	—	55–77	25–35	66	30
<i>C. myristiciformis</i>	Mississippi & Arkansas	14,500	5,110	—	—	207–375	94–170	273	124
<i>C. ovata</i>	—	17,600	6,200	38–49	30–38	176–331	80–150	220	100
	Wisconsin	—	—	—	—	—	—	291	32
	Mississippi	12,100	4,264	—	—	—	—	207	94

Source: Bonner and Maisenhelder (1974).

Storage tests with pecan and shagbark hickory have demonstrated that the hickories are orthodox in storage behavior, that is, they should be dried to low moisture contents and refrigerated. Seedlots of nuts of both species dried to below 10% moisture and stored at 3 °C in sealed containers retained viability well for 2 years before losing half to two-thirds of their initial viability after 4 years (Bonner 1976b). The poor results after 4 years are probably due to the high lipid levels in these seeds, which places them in the sub-orthodox storage category (Bonner 1990). There are no storage data for other species of hickory, but it is reasonable to think that they can be stored in a similar fashion.

Pregermination treatments. Hickories are generally considered to exhibit embryo dormancy, although work with pecan suggests that mechanical restriction by the shell is the reason for delayed germination in that species (van Staden and Dimalla 1976). Other research with pecan has shown that there is a clinal gradient in stratification requirement. Seedlots from southern sources are practically nondormant, whereas those from northern sources require treatment for prompt germination (Madden and Malstrom 1975). The common treatment is to stratify the nuts in a moist medium at 1 to 4 °C for 30 to 150 days (table 5). Stratification of

imbibed nuts in plastic bags without medium is suitable for most species (Bonner and Maisenhelder 1974), and good results have been reported for pecans from southern sources by soaking the nuts at 20 °C for 64 hours (Goff and others 1992). There are indications that stratification should be shortened for stored nuts; this was the case in one storage test on pecan and shagbark hickory (Bonner 1976b). If cold storage facilities are not available, stratification in a pit with a covering of about 0.5 m of compost, leaves, or soil to prevent freezing will suffice. Prior to any cold stratification, nuts should be soaked in water at room temperature for 2 to 4 days with 1 or 2 water changes each day to ensure full imbibition (Eliason 1965). There is evidence that germination of pecan can be increased by treatment with gibberellins (Bonner 1976a; Dimalla and van Staden 1977), but practical applications have not been developed.

Germination tests. Official testing rules for North America (AOSA 1993) prescribe testing pecan and shagbark

hickory at alternating temperatures of 20 °C (dark) for 16 hours and 30 °C (light) for 8 hours on thick creped paper for 28 days. Stratification for 60 days as described above is also recommended. Adequate germination tests can also be made on stratified nuts in flats of sand, peat, or soil at the same temperature regime (table 5). Quick tests with tetrazolium salts can also be used with hickories (Eliason 1965).

Nursery practice. Either fall-sowing with untreated seed or spring-sowing with stratified seed may be used. Excellent results with fall-sowing have been reported for shagbark hickory, but good mulching is necessary (Heit 1942). Drilling in rows 20 to 30 cm (8 to 12 in) apart and 2 to 4 cm ($3/4$ to 1 $1/2$ in) deep with 20 to 26 nuts/m (6 to 8/ft) is recommended; about 100 seedlings/m² (10/ft²) is a good density (Williams and Hanks 1976). Mulch should remain until germination is complete. Shading is generally not necessary, but shellbark hickory may profit from shade. Protection from rodents may be required for fall-sowings.

Table 5—*Carya, hickory*: stratification period, germination test conditions, and results

Species	Cold stratification (days)	Germination test conditions				Germination		Germination %	
		Medium	Temp (°C)		Days	Rate (%)	Days	Avg (%)	Samples
			Day	Night					
<i>C. alba</i>	90–150	Sand, peat, soil	30	20	93	54	64	66	4
<i>C. aquatica</i>	30–90	Soil	27–32	21	63	76	28	92	1
<i>C. cordiformis</i>	90	Sand, peat, soil	30	20	250	40	30	55	3
	90	Soil	27	21	50	60	50	60	1
<i>C. glabra</i>	90–120	Sand, peat, soil	30	20	30–45	—	—	85	2
<i>C. illinoensis</i>	30–90	Sand, peat	30	20	45–60	—	—	50	9
	30–90	Kimpak	30*	20	60	80	33	91	6
	30	Soil	32	21	35–97	—	—	75	2
<i>C. laciniosa</i>	90–120	Sand, peat, soil	30	20	45–60	—	—	—	—
<i>C. myristiciformis</i>	60–120	Kimpak	30*	20	60	53	50	60	2
<i>C. ovata</i>	90–150	Sand, peat	30	20	45–60	75	40	80	6
	60–120	Kimpak	30*	20	60	65	35	73	2

Source: Bonner and Maisenhelder (1974).

* Daily light period was 8 hours.

References

- Adams JC, Thielges BA. 1977. Research underway of pecan timber management. *Louisiana Agriculture* 20(2): 14–15.
- AOSA [Association of Official Seed Analysts]. 1993. Rules for testing seeds. *Journal of Seed Technology* 16 (3): 1–113.
- Bonner FT. 1971. Chemical contents of southern hardwood fruits and seeds. Res. Note SO-136. New Orleans: USDA Forest Service, Southern Forest Experiment Station. 3 p.
- Bonner FT. 1974. Chemical components of some southern fruits and seeds. Res. Note SO-183. New Orleans: USDA Forest Service, Southern Forest Experiment Station. 3 p.
- Bonner FT. 1976a. Effects of gibberellin on germination of forest tree seeds with shallow dormancy. In: Hatano I, Asakawa S, Katsuta M, Sasaki S, Yokoyama T, eds. Proceedings, 2nd International Symposium on Physiology of Seed Germination; 1976 October 18–30; Fuji, Japan. Tokyo: Government Forest Experiment Station: 21–32.
- Bonner FT. 1976b. Storage and stratification recommendations for pecan and shagbark hickory. *Tree Planters' Notes* 27(4): 3–5.
- Bonner FT. 1990. Storage of seeds: potential and limitations for germplasm conservation. *Forest Ecology and Management* 35: 35–43.
- Bonner FT, Maisenhelder LC. 1974. *Carya* Nutt., hickory. In: Schopmeyer CS, tech. coord. Seeds of woody plants in the United States. Agric. Handbk. 450. Washington, DC: USDA Forest Service: 269–272.
- Dimalla GG, van Staden J. 1977. The effect of temperature on the germination and endogenous cytokinin and gibberellin levels of pecan nuts. *Zeitschrift für Pflanzenphysiologie* 82: 274–280.
- Eliason EJ. 1965. Treatment of forest tree seed to overcome dormancy prior to direct seeding. In: Proceedings, Direct Seeding in the Northeast—A Symposium. Exp. Sta. Bull. Amherst: University of Massachusetts: 87–90.

-
- Goff WD, Brasher LR, McGuire JA. 1992. Germination of unstratified pecans is affected by exposure to high temperature and by soaking. *Scientia Horticulturae* 50(1/2): 159–163.
- Grauke LJ, Payne JA, Wood BW. 1990. North American pecans: a provenance study. Northern Nut Growers' Association Annual Report 80(1989): 124–131.
- Graves CH Jr, Diehl S, Thies K. 1989. Progress in breeding pecans for disease resistance. Bull. 963. Mississippi State: Mississippi Agricultural and Forestry Experiment Station. 8 p.
- Heit CE. 1942. Field germination and propagation of various hardwoods. For. Invest. Note 43. Albany: New York State Conservation Department.
- Herrera E, Martinez D. 1983. Effect of seed size on vigor of pecan seedlings. *Pecan Quarterly* 17(1): 18–19.
- Little EL. 1979. Checklist of United States trees (native and naturalized). Agric. Handbk. 541. Washington, DC: USDA Forest Service. 375 p.
- McCarthy BC, Quinn JA. 1990. Reproductive ecology of *Carya* (Juglandaceae): phenology, pollination, and breeding system of two sympatric tree species. *American Journal of Botany* 77: 261–273.
- Madden GD, Malstrom HL. 1975. Pecans and hickories. In: Janick J, Moore JN, eds. *Advances in fruit breeding*. West Lafayette, IN: Purdue University Press: 420–438.
- Sargent CS. 1965. *Manual of the trees of North America (exclusive of Mexico)*. 2nd ed., corrected and reprinted. New York: Dover Publications. 934 p.
- Short HL, Epps EA Jr. 1976. Nutrient quality and digestibility of seeds and fruits from southern forests. *Journal of Wildlife Management* 40: 283–289.
- van Staden J, Dimalla GG. 1976. Regulation of germination of pecan, *Carya illinoensis*. *Zeitschrift für Pflanzenphysiologie* 78: 66–75.
- Williams RD, Hanks SH. 1976. *Hardwood nurseryman's guide*. Agric. Handbk. 473. Washington, DC: USDA Forest Service. 78 p.
- Yates IE, Reilly CC. 1990. Somatic embryogenesis and plant development in eight cultivars of pecan. *HortScience* 25: 573–576.
- Yates IE, Sparks D. 1990. Three-year-old pecan pollen retains fertility. *Journal of the American Society for Horticultural Science* 115: 359–363.