# VISUAL GUIDE Winter Wheat

DEVELOPMENT AND GROWTH STAGING



### **Contents**

#### Introduction

**Basic wheat anatomy** 

Feekes scale for cereal growth stages

Zadoks scale for cereal growth stages

	Germination		
	Feekes <b>1</b>		1
ين	Feekes <b>2</b>		
9	Feekes <b>3</b>		
$\leq$	Feekes <b>4</b>	3.4	
STA	Feekes <b>5</b>		Design to
	Feekes <b>6</b> -	jointing / S	
工	Feekes <b>7</b>	A CP CONTRACTOR	
<u> </u>	Feekes <b>8</b>		
$\geq$	Feekes <b>9</b>		1
$\leq$	Feekes <b>10</b> -	<b>I</b>	
$\mathcal{L}$	Feekes <b>10.1</b>		
GROWTH	Feekes <b>10.2</b>		
	Feekes <b>10.3</b>		
	Feekes <b>10.4</b>	booting	
STRATE	Feekes <b>10.5</b>		_
	Feekes <b>10.5.1</b>		
$\gtrsim$	Feekes <b>10.5.2</b> -	<b>&gt;</b>	1
	Feekes <b>10.5.3</b>	\\0)	
5	Feekes <b>10.5.4</b>		
	Feekes <b>11.1</b>		
	Feekes <b>11.2</b>		
	Feekes <b>11.3</b>	flowering 🧗	
	Feekes <b>11.4</b>	-	٢
ner ce	reals		

#### Other cereals

Barley	
Oats	
Rye	
Triticale	

### Introduction

Understanding the growth stages of cereals crops and how to identify them is key to successful cropping and pest management decisions.

Although there are several growth staging methods, this guide is based on the Feekes scale, which is a popular tool used in the field. It has eleven development stages with some stages having more detailed subdivisions.

The Zadoks scale is the standard scale used in research and has ten development stages, each stage having ten subdivisions. Both scales are useful to know, so this guide cross-references the Zadoks equivalents to the Feekes.

This guide uses winter wheat as an example. However, the methods generally apply to other cereals as well and at the back of the guide are sections that showcase barley, oats, rye and triticale.

#### A few notes on growth staging plants:


- · Select plants that represent at least 50% of the field
- Dig plants (if possible), so you can assess the entire plant
- · Start at the base of plant and work your way upward
- · Use a knife to split the stems and sheaths
- · Look and feel for nodes

#### **References:**

Large, E. C. (1954). *Growth Stages in Cereals Illustration of the Feekes Scale*. Plant Pathology, 3: 128–129. doi:10.1111/j.1365-3059.1954.tb00716.x

Feekes, Willem (1941). *De Tarwe en Haar Milieu* [*Wheat and Its Environment*]. Verslagen van de Technische Tarwe Commissie. (in Dutch (English summary)). 17: 523–888.

J.C. Zadoks, T.T. Chang, C.F. Konza. *A Decimal Code for the Growth Stages of Cereals.* Weed Research 1974 14:415-421.





Auricles blunt and hairy; leaf sheath and blade always hairy; ligule medium length; leaf blades twist clockwise


## Feekes scale for cereal growth stages

SEEDLING GROWTH	1	One shoot, first leaf through coleoptile	
	2	Tillering begins; main shoot and one tiller	
TILLERING	3	Tillers formed; leaves often twisted In some varieties, plant may be prostrate in appearance	
員	4	Leaf sheaths lengthen; beginning pseudostem erection	
	5	Leaf sheaths fully elongated to form strongly erect pseudostem	
	6	First node of stem visible at base of shoot; jointing	
STEM EXTENSION	7	Second node of stem formed; next-to-last leaf just visible	
EXTE	8	Flag leaf visible but still rolled up	
E	9	Ligule of flag leaf just visible	
rs S	10	Flag leaf sheath completely grown out; booting	
	10.1	First awns of head just visible	
و	10.2	1/4 of heading process complete	
ADII	10.3	1/2 of heading process complete	
쁖	10.4	3/4 of heading process complete	
	10.5	All heads out of sheath	
U	10.5.1	Beginning of flowering	
FLOWERING	10.5.2	Flowering complete to top of head	
LOW	10.5.3	Flowering complete at base of head	
ш.	10.5.4	Flowering complete; kernel watery ripe	
	11.1	Kernel milky ripe; milk stage	
RIPENING	11.2	Kernel mealy ripe; soft but dry consistency; soft dough stage	
RIPE	11.3	Kernel hard; difficult to divide with thumbnail; hard dough stage	
	11.4	Kernel harvest ready; straw dead	

## Zadoks scale for cereal growth stages


	00	Dry seed				
on	01	Start of imbibition				
Germination	03	Imbibition complete				
ï	05	Radicle emerged from seed		Fee	Feekes	
99	07	Coleoptile emerged from seed		sc	scale	
	09	Leaf just at coleoptile	tip	equi	valent	
	10	First leaf through cole	optile		1	
	11	First leaf unfolded				
	12	2 leaves unfolded				
wth	13	3 leaves unfolded	A leaf is unfolded			
gro	14	4 leaves unfolded	when its ligule is			
ü	15	5 leaves unfolded	visible, or the tip			
Seedling growth	16	6 leaves unfolded	is visible			
	17	7 leaves unfolded				
	18	8 leaves unfolded				
	19	9 or more leaves unfo	lded	ی		
	20	Main shoot only		TILLERING		
	21	Main shoot and 1 tille	r		2	
	22	Main shoot and 2 tille	rs			
_	23	Main shoot and 3 tille	rs			
Tillering	24	Main shoot and 4 tille	rs			
E	25	Main shoot and 5 tille	rs			
	26	Main shoot and 6 tille	rs		3	
	27	Main shoot and 7 tille	rs			
	28	Main shoot and 8 tille	rs			
	29	Main shoot and 9 or n	nore tillers			
	30	Pseudostem erection			4-5	
	31	1st node detectable			6	
on	32	2 <sup>nd</sup> node detectable			7	
gati	33	3 <sup>rd</sup> node detectable		SIO		
Stem elongation	34	4 <sup>th</sup> node detectable		STEM EXTENSION		
em (	35	5 <sup>th</sup> node detectable		X		
St	36	6 <sup>th</sup> node detectable				
	37	Flag leaf just visible		ν	8	
	39	Flag leaf ligule/collar	just visible		9	


	40		Z	
6	41	Flag leaf sheath extending	STEM EXTENSION	
Booting	45	Boot just visibly swollen	X	10
B0	47	Flag leaf sheath opening		
	49	First awns visible	STE	
	50	First spikelet of inflorescence visible		10.1
ence	53	1/4 of inflorescence emerged	وِ	10.2
nflorescenc emergence	55	1/2 of inflorescence emerged	HEADING	10.3
Inflorescence	57	3/4 of inflorescence emerged	里	10.4
_	59	Emergence of inflorescence completed		10.5
sis	60	Beginning of anthesis		10.5.1
Anthesis	65	Anthesis half-way	وِ	10.5.2
Ar	69	Anthesis completed	FLOWERING	10.5.3
	70		MO	
nent	71	Kernel watery ripe	교	10.5.4
Milk lopr	73	Early milk		
Milk development	75	Medium milk		11.1
	77	Late milk		
int	80			
ngh	83	Early dough	S <sub>G</sub>	
Dough evelopment	85	Soft dough	RIPENING	11.2
de	87	Hard dough	불	
	90			
	91	Kernel hard (difficult to divide with thumbnail)		11.3
	92	Kernel hard (no longer dented with thumbnail)		11.4
	93	Kernel loosening in daytime		
Ripening	94	Overripe, straw dead and collapsing		
Ripe	95	Seed dormant		
	96	Viable seed giving 50% germination		
	97	Seed not dormant		
	98	Secondary dormancy induced		
	99	Secondary dormancy lost		














EMERGENCE | Feekes 1 | One shoot formed; first leaf through coleoptile









## Are tillers important?

## Tillers are absolutely necessary for high yields

Feekes

can produce planted seed

4-5

are also called axillary or side shoots; not all tillers will complete development and produce grain

The total numbers of tillers
a plant produces is determined by both
environmental conditions and genetic potential

#### **KEY YIELD COMPONENT**

A **tiller** is capable of forming a single head (spike)
The **head** is made up of spikelets
Each **spikelet** contains individual florets
Individual **florets** can produce a single **kernel** 



In Wisconsin, the recommended planting date range for optimal tiller development in winter wheat is **September 20** to **October 10** 





#### FEEKES 3 CAN OCCUR IN FALL OR SPRING

because winter wheat development is dependent on both temperature and planting date

Feekes



Tillering completes in fall, winter dormancy occurs



Tillering begins in the fall, winter dormancy occurs, tillering completes in the spring

# What happens during winter dormancy?

## Vernalization!



## differentiation

the growing point changes from vegetative or leaf producing to reproductive or spikelet producing

The growing point is at the **double ridge stage** and is still protected in the crown below the soil surface

The number of florets initiated during this stage will determine the **potential** number of kernels per head


**KEY YIELD COMPONENT** 



Dig plants as soon as the soil thaws, bring inside and place in a warm (preferably moist) area for a few days, then check for root regrowth



eeke 3





- Good snow cover acts as insulator; keeps soil temperature from going below critical levels
- Cyclic freezing and thawing increases injury from ice crystal growth in tissue
- Mid-winter thaw and rain cause flooding at the base of the plants; crowns can die at warmer temperatures
- Ice encasement traps carbon dioxide and suffocates plant by inhibiting respiration
- Frost heaving can push root system out of ground, leaving plants vulnerable and weak

GREEN UP

## 4 STEPS

## TO ASSESS STANDS IN EARLY SPRING

- 1 Venture out and get a general overview of the fields — vibrant green patches may be interspersed with drab brown areas, but brown does not always indicate winter-killed plants
- 2 Check for winter survival identify several representative plants and 1) dig plants and bring inside to check for root regrowth or 2) wait a week and revisit to check for regrowth in the field
- 3 Do a plant count
  - below 12 live plants per square foot is an automatic replant; 12-15 live plants per square foot requires more consideration for a replant decision; 15-22 live plants per square foot may recover and reach maximum yield potential; over 22 live plants per square foot means you're good to go!



Consider a nitrogen application — the optimal time to apply nitrogen to winter wheat in Wisconsin is during green up; for recommendations and rates, consult UW-Extension publication A2809 Nutrient Application Guidelines for Field, Vegetable and Fruit Crops in Wisconsin

In Wisconsin, the growth stage at green up can be **Feekes 3** or **Feekes 4** depending on planting date and environmental conditions

GREEN UP



#### How to do a plant count

Count the number of plants in a 3-foot length
Do this for at least 3 areas
Take the average of the counts
Multiply that number by 4
Then divide by the row width (inches)

#### **EXAMPLE**

The 3 counted areas have 40, 35 and 45 plants Add 40, 35 and 45 and then divide by 3, the average = 40 Multiply 40 x 4 = 160 Divide 160 by 7.5 inches = 21 plants/square foot





TILLERING | Feekes 4 | Leaf sheaths lengthen, pseudostem erection begins

## This is an important time for weed control and/or nitrogen applications

Wheat plants have a *pseudostem*, which is a false stem composed of concentric rolled leaf sheaths that surround the growing point (the developing head)

During this stage, these leaf sheaths lengthen, making the plants stand more upright

Feekes **4** 






TILLERING | Feekes 5 | Leaf sheaths fully elongated, pseudostem strongly erect





STEM EXTENSION | Feekes 6 | First node of the stem visible at the base of the shoot; jointing










STEM EXTENSION | Feekes 7 | Two nodes visible above the soil line









**Kernel size** is determined by crop health and water/nutrient availability beginning now and continuing through grain fill





STEM EXTENSION | Feekes 9 | Flag leaf fully emerged from the whorl; ligule just visible





completely grown out; head visible in the leaf sheath; booting






When determining the growth stage of a field, 50% of the plants must be at that stage or above Feekes 10.1 Feekes 10.2




HEADING | Feekes 10.2 | 1/4 of the head emerged from the leaf sheath











HEADING | Feekes 10.4 | 3/4 of the head emerged from the leaf sheath






HEADING | Feekes 10.5 | Head completely emerged from the leaf sheath




Feekes 10.5.1 Starting now and continuing 5-7 days after this stage is the optimum time for fungicide application to suppress Fusarium head blight (FHB), also called head scab

Flowering begins slightly above the middle portion of the head and continues towards the top



## The number of **flowers pollinated** determines the number of **kernels that will develop**




### anther

The male flower part that produces and releases pollen

### pollen

The powder-like grains that enable fertilization



FLOWERING | Feekes 10.5.2 | Flowering complete to the top of the head





FLOWERING | Feekes 10.5.3 | Flowering complete at the base of the head




This stage signals the end of pollination

Feekes 10.5.3



floret outer structure removed to show developing kernel

This is the beginning of the **grain filling stages**; kernel length is established during this stage






When squeezed, **clear fluid** is released from the kernel



Kernel size increases but not dry matter accumulation





RIPENING | Feekes 11.1 | Kernel milky ripe; milk stage





RIPENING | Feekes 11.2 | Kernel mealy ripe; soft but dry consistency; soft dough stage




Green color of the kernel, glume and peduncle begins to fade

### Starch, nutrients and dry matter

accumulate rapidly in the kernel



The kernel's content is a soft-doughy material







RIPENING | Feekes 11.3 | Kernel hard; difficult to divide with a thumbnail; hard dough stage



# Kernels reach their **maximum dry weight** and are **physiologically mature**





RIPENING | Feekes 11.4 | Kernel harvest ready; straw dead





# Barley

Auricles long, slender and hairless; leaf sheath and blade usually hairless (scattered hairs on some varieties); liqule medium length; leaf blades twist clockwise





Oats

Auricles absent; leaf sheath and blade hairless (scattered hair on some varieties); ligule medium length; leaf blades twist counter-clockwise





Rye

Auricles very short and hairless; leaf sheath and blade have an inconsistent degree of hairiness; ligule short; leaf blades twist clockwise





each other during the vegetative stage

An alternative method is to remove a seedling from the soil and check the grain shell; triticale shells are oblong in shape and dark in color, while wheat grain shells are oval and lighter



Auricles blunt and hairy, leaf sheath and blade hairy; liqule of medium length; leaf blades twist clockwise



#### **Authors**

Mimi Broeske, Nutrient and Pest Management Program; Shawn Conley, John Gaska and Adam Roth, Department of Agronomy; University of Wisconsin-Madison, College of Agricultural and Life Sciences and Division of Extension

### **Acknowledgements**

The authors would like to thank our external reviewers: Mike Cerny, Walworth County farmer Dr. Chad Lee, University of Kentucky Dr. David Marburger, Oklahoma State University



This publication is available from the **Nutrient and Pest Management Program:** web (ipcm.wisc.edu); phone (608) 265-2660; email (npm@hort.wisc.edu)

